

The Usage of the Term “Formefruste” in Ophthalmology

Editorial

In this editorial, I aimed to clarify the term “*formefruste*” and to take care to avoid its unnecessary use. Formefruste is a term defining “an incomplete or partial or arrested or in apparent or atypical or aborted or attenuated phenotypic expression or manifestation of a clinical entity or disease or syndrome or pathological condition. It often defines that disease is subclinical and that disease has not the main diagnostic criteria and the expected and common course of the disease [1]. In ophthalmological terminology, the term “*formefruste*” has been used to call “the occult form of an ocular disease”. To literature, the ophthalmological diseases in which the term has been used are keratoconus (KK), anterior segment dysgenesis, central serous chorioretinopathy, retinopathy of prematurity, hemorrhagic occlusive retinal vasculitis, juvenile idiopathic arthritis-associated uveitis, sympathetic uveitis, focal choroidal excavation in serpiginous choroiditis [2-11]. Additionally, it has been also reported for defining of atypical clinical forms of some neuro-ophthalmological diseases such as multiple sclerosis, tuberosis sclerosis, Vogt-Koyanagi Syndrome and idiopathic intracranial hypertension, trilateral retinoblastoma and acne rosacea as dermatological disease [12-17]. Actually, any ocular finding may be a sign of a formefruste of any ocular, extraocular or systemic disease. Additionally, the term “*formefruste*” has been not been choiced in ophthalmological literature except only for the nomenclature of the frank, early stage or subclinical or preclinical or symptomless manifestation of KK for a long time [18,19]. Formefruste KK has mild corneal topographic characteristics suggestive of an early subclinical KK [2,18,19]. To my opinion, the usage of a term “*occult*” or “*atypical*” seems more appropriate for incomplete forms of the ocular diseases instead of the terms of “*formefruste*” and “*subclinical disease*”.

Conflict of Interest

The author declares that there is no conflict of interest regarding the publication of this paper.

Acknowledgment

None.

References

1. Forme_fruste
2. Saad A, Gatinel D (2010) Topographic and tomographic properties of forme fruste keratoconus corneas. Invest Ophthalmol Vis Sci 51(11): 5546-5555.
3. Banitt MR, Romano A, Iragavarapu S, Budenz DL, Lee RK (2011) Forme fruste anterior segment dysgenesis. Br J Ophthalmol 95(12): 1756-1757.
4. Spaide RF (2005) Central Serous Chorioretinopathy. In: Holz FG & Spaide RF (Eds.), Medical Retina. Essentials in Ophthalmology. Springer, Berlin, Heidelberg, Germany, p. 6-93.
5. Silverman WA (2001) ROP--formefruste. J Perinatol 21(6): 393-394.
6. Baum JD (1971) Retinal photography in premature infants: forme frusta retrolental fibroplasia. Proc R Soc Med 64(7): 777-779.
7. Goldberg RA, Crawford C, Heier JS (2016) Postoperative Hemorrhagic Occlusive Retinal Vasculitis: A Forme Fruste Variant? Retin Cases Brief Rep.
8. Dashti SR, Robinson S, Rodgers M, Cohen AR (2005) Pineal region giant cell astrocytoma associated with tuberous sclerosis: case report. J Neurosurg 102(3 Suppl): 322-325.
9. Shields JA, Eagle RC, Shields CL, Marr BP (2004) Aggressive Retinal Astrocytomas in Four Patients with Tuberous Sclerosis Complex. Trans Am Ophthalmol Soc 102: 139-148.
10. Kotaniemi K, Savolainen A, Aho K (2003) Severe childhood uveitis without overt arthritis. Clin Exp Rheumatol 21(3): 395-398.
11. Roy R, Saurabh K, Vyas C, Sharma P, Bansal A (2017) Forme fruste focal choroidal excavation in a case of serpiginous choroiditis. Clin Exp Optom.
12. Burton EV, Greenberg BM, Frohman EM (2011) Optic neuritis: A mechanistic view. Pathophysiology 18(1): 81-92.
13. Yanoff M, Sassani JW (2015) Granulomatous inflammation. In: Myron Yanoff & Joseph W Sassani (Eds.), Ocular Pathology E-Book: Expert Consult. (7th edn), Elsevier-Saunders, London, p. 67.
14. O'Keefe GA, Rao NA (2017) Vogt-Koyanagi-Harada disease. Surv Ophthalmol 62(1):1-25.
15. Bahall M, Reyes AJ, Ramcharan K, Hosein N, Seegobin K, et al. (2016) Neuro-Ophthalmological Manifestations after Intramuscular Medroxyprogesterone: A FormeFruste of Idiopathic Intracranial Hypertension? Neurol Int 8(3): 6132.
16. Whittle IR, McClellan K, Martin FJ, Johnston IH (1987) Concurrent pineoblastoma and unilateral retinoblastoma: a formefruste of trilateral retinoblastoma? Neurosurgery 17(3): 500-505.
17. Blaustein BH, Gurwood AS (2001) Recurrent phlyctenular keratoconjunctivitis: a forme fruste manifestation of rosacea. Optometry 72(3): 179-184.
18. Amsler M (1961) [The “formefruste” of keratoconus] (Article in German)]. Wien Klin Wochenschr 73: 842-843.
19. Sinjab MM (2012) Classifications and patterns of keratoconus and keratectasia. In: Quick Guide to the Management of Keratoconus. Springer, Berlin, Heidelberg, Germany, p. 13-58.

Editorial

Volume 7 Issue 5 - 2017

Burak Turgut*

Department of Ophthalmology, Yuksekiltisas University, Faculty of Medicine, Turkey

*Corresponding author: Burak Turgut, Department of Ophthalmology, Faculty of Medicine, Full Professor of Ophthalmology, Yuksekiltisas University, 06520, Ankara, Turkey, Tel: +90 312 2803601; Fax: +90 3122803605; Email: burakturgut@yiu.edu.tr

Received: October 12, 2017 | Published: October 12, 2017