

# Arbuscular Mycorrhiza Fungi, NPK (15-15-15) and Cow Dung Interaction in Sustainable Cassava Production and Food Security

## Abstract

Increasing human population coupled with the depletion and degradation of soil resources constitute a threat to food security in sub-Saharan Africa. Sequel to this, the growth, performance and yield of cassava (*Manihot esculenta* L.) were assessed using pure culture of arbuscular mycorrhiza fungus - *Glomus deserticola*, NPK (15:15:15) and Cow dung singly and in combination with each other at four treatment levels under field conditions. Control experiment was also set up without any treatment. The experiment was factorial, in completely randomized block design and replicated four times. Cassava stem cuttings, 18 cm length were planted in well tilled soil at a distance of 1 m apart and allowed to grow for six months. During the study, plant height, stem girth, leaf area, leaf chlorophyll content and yield were assessed. At harvest, fresh and dry tuber weights were measured. Data collected were subjected to one way analysis of variance and means separated using Tukey's HSD test ( $p \leq 0.05$ ). The study showed that *Glomus deserticola* in combination with NPK produced cassava plants with significant differences ( $p < 0.05$ ) in height, stem girth, leaf area, chlorophyll content and yield. Arbuscular mycorrhiza fungus enhanced nutrient uptake of cassava plants. The balanced fertilization and amendment of the experimental soil with adequate nutrients supply provided useful agronomic information on the performance and yield of cassava.

**Keywords:** Soil fertilization; Amendment; *Manihot esculenta*; Growth performance; Food security

## Research Article

Volume 7 Issue 4 - 2017

**Chukwuka KS<sup>1\*</sup>, Okechukwu RU<sup>2</sup>, Umukoro BO<sup>1</sup> and Obiakara MC<sup>1</sup>**

<sup>1</sup>Department of Botany, University of Ibadan, Nigeria

<sup>2</sup>Cassava Transformation Agenda Project, International Institute of Tropical Agriculture, Nigeria

\*Corresponding author: Chukwuka KS, Department of Botany, University of Ibadan, Ibadan-Nigeria, Email: kanayodrchukwuka97@gmail.com

Received: July 24, 2017 | Published: August 22, 2017

## Introduction

Cassava (*Manihot esculenta* Crantz) is a woody shrub that is widely cultivated in many tropical and subtropical regions of the world. It is propagated from stem cuttings and produces edible energy-rich tubers. Dry cassava tubers contain up to 90% carbohydrates, of which starch is dominant Montagnac et al. [1]. The importance of this crop in food security is reflected in its remarkably increasing production over the last decade. In 2014, about 270 million tonnes constituting about 54% of global production were produced in Africa FAOSTAT [2]. This crop is reputed for its remarkable tolerance to abiotic stresses due to its biochemical and physiological adaptations, intermediate C<sub>3</sub>-C<sub>4</sub> photosynthetic pathway, reduction in leaf area index coupled with a relatively high stomatal sensitivity in response to water shortages and unaltered photosynthetic rates in nutrient-deficient soils El-Sharkawy [3]. These characteristics make it suitable for both small and large-scale cultivation with moderate agricultural inputs. However, sustainable cassava production especially in the face of both increasing reliance on fertilizers and recurring environmental issues associated with their long-term use requires exploring alternative approaches that could help meet the food demands of rapidly expanding populations.

The use of organic residues in combination with mineral fertilizers is a widely accepted soil fertility management strategy Edmeades [4], Diacono and Montemurro [5], Simon and Czako

[6]. This practice is known to improve biological, chemical and physical properties of agricultural soils as opposed to the sole application of inorganic fertilizers Kotschi [7]; Mulvaney et al. [8]. In fact, consistently lower cassava yields have been recorded under sole inorganic fertilizer application compared to when organic manures are combined with inorganic fertilizers Ayoola and Adeniyani [9]; Ojeniyi et al. [10]. For example, Ojeniyi et al. [10], reported that combination of 2.5 t/ha of poultry manure and one quarter less the recommended amount of inorganic fertilizer (NPK 15-15-15; 600 kg/ha) led to twofold increase in yield of cassava compared to sole application of the same inorganic fertilizer at 600 kg/ha after 6 months of cultivation. Similarly, Islami et al. [11], reported that the application of chemical fertilizers in monoculture cassava was inadequate to maintain sustained yields; but combining farm yield manure (FYM) with chemical fertilizers increased soil fertility and cassava production. Arbuscular Mycorrhizal Fungi (AMF) also referred to as Vesicular Arbuscular Mycorrhiza (VAM) are widespread in terrestrial ecosystems and form mutually beneficial associations with nearly 80% of higher plants Smith et al. [12]. They are characterized by their intercellular and intracellular growth forms in plant roots, which are referred to as vesicles and arbuscules Böhm and Hock [13]. Mycorrhizal symbioses are known to mitigate the problem of efficient uptake of immobile nutrients by plants Bolan [14]; Smith and Read [15]. Previous studies have shown that AMF are associated with salt Carretero et al. [16], and drought tolerance

Qiangsheng et al. [17]; Ruiz-Lozano et al. [18] and cassava is known to thrive under these conditions. The potentials of AMF in sustainable crop production have been demonstrated by many scholars Li et al. [19]; Cely et al. [20]; Tchabi et al. [21]. However its applicability has not been fully addressed.

Cassava tolerates low soil nutrient levels. However, this crop needs substantial fertilization to attain high yields like many other food crops Howeler [22]. Perhaps, cassava would not have been a successful crop under both fertilized and un-fertilized conditions without its dependency on mycorrhizal fungi for nutrient uptake, especially phosphate uptake Habte and Byappanahalli [23]. These authors reported cassava mycorrhizal dependency (that is, the change in cassava growth due to arbuscular mycorrhizal colonization) of 60% in contrast to the much lower values (44-46%) obtained in many crop species Tawaraya [24]. Recently, Burns et al. [25], showed that mycorrhizal dependency of cassava could reach up to 93%. This dependency is indicative of the wide variety of AMF associated with cassava roots as *Glomus*, *Gigaspora* and *Acaulospora* which are the most commonly reported genera Straker et al. [26]; Bi Voko et al. [27]; dos Santos Heberle et al. [28]; Begoude et al. [29]. Several species in the genus *Glomus* including *G. manihotis* Sieverding et al. [30]; *G. fasciculatum*, *G. clarum*, *G. mosseae* Oyetunji and Osonubi [31]; *G. deserticola* Okon et al., [32]; *G. intraradices* Carretero et al. [33]; *G. aggregatum* [23] and *G. etunicatum* Salami et al. [34] have been tested in field studies with or without fertilization. Obviously, cassava's poor root system architecture, a factor that is crucial in nutrient uptake and subsequent productivity in many crops Smith and De Smet [35] is a major cause of its dependency on mycorrhiza for nutrient acquisition.

The beneficial effects of AMF association with cassava have been the subject of several studies. Howeler et al. [35] reported a significant increase in growth and dry matter content of mycorrhiza-inoculated cassava cuttings compared to non-inoculated plants, which were P-deficient even at high P soil levels. Sieverding and Toro [30] found that N-P-K concentration ratios in cassava shoots and roots were more balanced in mycorrhiza-inoculated plants than in non-inoculated plants. In the same vein, Ceballos et al. [36] showed that not only 20% increase in cassava yield was obtained due to addition of *Rhizophagus irregularis* but also a 50% reduction in phosphate fertilizer. Other beneficial effects have been reported including enhanced plantlet survival, shoot, root and tuber formation Azcón-Aguilar et al. [37]; salt damage alleviation Carretero et al. [33] and resistance to transplant stress Carretero et al. [16]. Given the growing body of evidence on the beneficial effects of AMF on cassava, the present study tested effects of AMF and its combination with organic and inorganic fertilizers to improve cassava productivity in a sustainable manner. Therefore this study hypothesized that soil inoculated with *Glomus deserticola* will enhance cassava growth.

## Materials and Methods

### Study area

Smith et al. [12], This study was carried out at the Botany Research Farm, Department of Botany, University of Ibadan

(7°26.44' N, 3°53.76' E) between 18 July 2016 and 18 January 2017. The University has mean annual rainfall and temperature of 1316 mm and 27.6°C respectively

### Soil and cow dung analyses

Soil samples were randomly taken from 0-20 cm depth before planting, bulked, air-dried and sieved using 2 mm sieve for analysis. The particle size analysis was done by pipette method Gee and Bauders [38]. Soil pH in water was determined using soil: water ratio of 1:2 with a glass electrode pH meter. Organic carbon was determined using Walkley and Black method Nelson and Sommers [39]. Total nitrogen (N) in the soil was determined by Kjeldahl digestion Bremner [40]. Exchangeable bases in the samples were extracted in 1M NH<sub>4</sub>OAc at pH 7.0. Calcium (Ca) and magnesium (Mg) in the extract were read by atomic absorption spectrophotometer (AAS). Sodium (Na) and potassium (K) were analyzed by flame photometry. Available phosphorus (P) was determined by Bray-1 extraction and determined colourimetrically by the molybdenum blue procedure Bray and Kurtz [41].

Cow dung samples were air-dried and ground to powder and analysed with wet digestion method using 5:1:1 ml of HNO<sub>3</sub>: H<sub>2</sub>SO<sub>4</sub>:HClO<sub>4</sub> acid. Total N was determined by micro-Kjeldahl method Jackson [42]. For P, K, Ca and Mg, samples (0.5 g) were ashed, dissolved in 10% hydrogen chloride (HCl) and diluted to 50 ml. Phosphorous was determined using vanadomolybdate colorimetric. Calcium and Magnesium were determined by EDTA titration while Na and K by flame photometry. The physico-chemical properties of both soil and cow dung used were analyzed at the Department of Agronomy, University of Ibadan.

### Experimental design and treatment application

**The experiment involved seven treatments:** AMF (*Glomus deserticola*) 20, 30, 40 and 50g), NPK 15:15:15 40, 60, 80 and 100g; cow dung 200, 300, 400 and 500g; AMF and NPK 20 + 40, 30 + 60, 40 + 80 and 50 + 100g, AMF and cow dung 20 + 200, 30 + 300, 40 + 400, 50 + 500g, cow dung and NPK 40 + 200, 60 + 300, 80 + 400, 100 + 500g laid out in a completely randomized design (CRD) with four replicates. The soil was tilled before planting. Cassava stem cuttings (TME 419) of about 15 cm long were planted horizontally and buried completely at 5 cm depth in heaps of soil at a planting distance of 1m apart in a plot size of 14 m × 14 m. The treatments were applied two weeks after establishment. Application of NPK 15:15:15, cow dung and *Glomus deserticola* was done using the methods of Ojeniyi et al. [10], Mathias and Kabambe [43] and Okon et al. [44]. Respectively, four different levels of each *G. deserticola* (20g, 30g, 40g and 50g); NPK 15:15:15 (40g, 60g, 80g and 100g) and cow dung (200g, 300g, 400g and 500g) were applied both singly and in combination around the growing stem cuttings.

### Data collection

Data collection commenced two weeks after application of treatments and subsequently forth nightly for five months. The following growth parameters: plant height, stem girth and leaf area were determined forth nightly. Plant height (cm) and leaf

area (cm<sup>2</sup>) were determined using a Measuring Tape and Portable Electronic Area Metre Model Li-3000 respectively. Stem girth was measured at 5 cm above heap level using Mitutoyo Digimatic Electronic calliper (MDEC) Model CD-8" P. The leaf chlorophyll content was determined by using the central leaf from freshly excised leaves of the same age. The excised leaf was weighed, cut into smaller pieces and stored in the dark for 24 hrs in a mixture of 10 ml of 95% ethanol and 99.5% acetone (1:1 v/v). One millilitre of the concentrated leaf extract was added to a cuvette and adjusted to 5 ml with the ethanol-acetone mixture. The Absorbance of the mixture was measured spectrophotometrically at 652 nm using a Unico spectrophotometer. This value was taken as reference by resetting the absorbance to zero and measuring that of the diluted leaf extract. Chlorophyll concentration (ml/g) was estimated using equation (1)

$$\text{Chlorophyll Concentration} = \frac{A_{652}}{34.5} \times \frac{V}{1000W}$$

Where A<sub>652</sub> = Sample OD value (Absorbance at 652 nm)

V = Volume of the sample (ml)

W = Weight of the sample (g)

The yield data was collected six months after planting and these include fresh and dry weight of tubers. The plants were harvested and separated into roots and stems. Dry weight was determined after oven drying at 100°C for 48 hours using Ohaus Sensitive Electronic Digital Weighing Balance Model SPX2202.

### Data analysis

Data were analysed using Minitab 16 Statistical Software (2010). A one way Analysis of Variance (ANOVA) was carried out to test the effects of the treatments on cassava growth and development. Means were separated using Tukey's HSD test (p≤0.05).

### Results

Table 1 shows the values of the soil physicochemical properties of the experimental site and the nutrient composition of cow dung used in the study. The soil was silt loamy and slightly acidic. Total nitrogen and Organic carbon were higher than 0.11% and 2% respectively, which are critical values for Nigerian soils Adepetu [45]. The soil was poor in Phosphorus (Available P < 10 mg/Kg) while potassium was detected in adequate amounts (Exchangeable K > 0.2 C mol (+)/Kg. Effective CEC (sum of equivalent charge concentrations of cations (Ca<sup>2+</sup>, Mg<sup>2+</sup>, K<sup>+</sup> Na<sup>+</sup> and Al<sup>3+</sup>) was low (i.e between 5-15 C mol (+) Kg<sup>-1</sup>).

### Cassava plant height (cm)

The study showed significant differences in plant height from 8 weeks after establishment (WAE). Between 8 to 12WAE, three main effects were distinguished. AMF and NPK combination (30 g of AMF and 60 g of NPK) significantly enhanced plant height (133 ± 8, 145 ± 10 and 149 ± 13 cm respectively). On the other hand, sole application of cow dung did not produce plants with appreciable increase in height (57 ± 9, 71 ± 8 and 70 ± 14 cm respectively). Other treatments produced plant heights similar

to the control. On the other hand, from 14WAE to 20WAE, the plants showed decreasing performance in height in the order AMFNPKL2 (30 + 60) g > NPK 100 g and CDNPKL4 (500 + 100) g > control > CD (400 g) > CD (200g) (Table 2).

**Table 1:** Physico-chemical characteristics of the soil at the experimental site, Botany Research Farm, University of Ibadan and chemical composition of cow dung.

Parameter	Soil	Cow Dung
pH	5.86	-
Nitrate	4.66 g/Kg	1.81%
Organic carbon	54.75 g/Kg (H)	-
Phosphate	-	0.53%
Average phosphorus	6.52 mg/Kg (L)	-
Exchangeable aluminium	0.5 C mol (+)/Kg	-
Calcium	8.86 C mol (+)/Kg	0.28%
Magnesium	0.69 C mol (+)/Kg	0.14%
Potassium	0.55 C mol (+)/Kg	1.04%
Sodium	0.18 C mol (+)/Kg	0.22%
Manganese	17.9 mg(+)/Kg	2160 mg(+)/Kg
Iron	224 mg(+)/Kg	17.5 mg(+)/Kg
Copper	1.34 mg(+)/Kg	15.05 mg(+)/Kg
Zinc	7.83 mg(+)/Kg	1625 mg(+)/Kg
Silt	1560 g/Kg	-
Clay	116 g/Kg	-
Sand	728 g/Kg	-

### Cassava leaf area (cm<sup>2</sup>)

There was a general increase in leaf area from 8 -12WAE. However, from 14 - 20WAE, the leaf area showed general decrease in size. On the other hand, there were significant differences at 8 and 10WAE among treatments for plants with AMF and NPK combination (30 + 60 g) having the widest leaf areas while plants treated with cow dung (400 g) showed the least leaf area. Although at 12WAE and 16 -20WAE, no significant differences among treatments were observed. It is important to note that the interaction of AMF with any combination of treatment produced plants with big leaf areas (Table 3).

### Cassava stems girth (mm)

Significant differences in plant stem girth were observed from 10-20 WAE. The combination of AMF and inorganic fertilizer (30 g of AMF and 60 g of NPK) produced plants with the largest stem girth while plants treated with cow dung (200 g) produced the least girth. Stem girth showed similar growth pattern throughout the period of study (Table 4).

### Cassava leaf chlorophyll content (ml/g)

The effects of different treatments on the leaf chlorophyll content at 10 weeks after planting and establishment of the plants are represented in Figure 1. The mean chlorophyll concentration was highest in the combined application of inorganic fertilizer and *G. deserticola* (0.1825 ± 0.0007 ml/g) and lowest with cow dung in combination with *G. deserticola*. Chlorophyll in plants treated with sole NPK was next (0.15497 ± 0.000246 ml/g) after the combined application of *G. deserticola* and NPK. Cow dung, *G. deserticola* and the control treatments had similar effect on leaf chlorophyll content with concentrations of 0.0909 ± 0.0006, 0.0922 ± 0.0005 and 0.0925 ± 0.0004 ml/g respectively (Figure 1).

### Yield

Figure 2 represents the fresh cassava weight obtained from the different treatments. Significant differences were observed among the treatments. As in other growth parameters, the highest yield was obtained in cassava treated with the mixture of 30 g of *G. deserticola* and 60 g of NPK (0.87 ± 0.07 Kg) while the lowest yield was observed in all the cow dung treatments. The control (0.23 ± 0.05 Kg) was comparable to all other treatments. The figure also show that dry weight of the tubers follow the same pattern as the fresh weight with the combined AMF-NPK (30 g+60 g) producing the highest dry weight (0.17 ± 0.01 Kg) while cow dung (300 g) alone produced the lowest tuber dry weight (0.008 ± 0.006 Kg).

**Table 2:** Effects of treatments on cassava plant height (cm).

Treatment	8WAE	10WAE	12WAE	14WAE	16WAE	18WAE	20 WAE
Control	97.33±19.66 <sup>ab</sup>	100.10±16.42 <sup>ab</sup>	122.42±5.47 <sup>ab</sup>	128.98±2.37 <sup>abc</sup>	128.55±1.61 <sup>ab</sup>	128.62±1.40 <sup>abc</sup>	131.15 ± 2.38 <sup>abc</sup>
AMF 20 g	85.55±16.66 <sup>ab</sup>	94.70±15.66 <sup>ab</sup>	100.72±17.79 <sup>ab</sup>	108.35±16.38 <sup>abc</sup>	114.18±14.93 <sup>ab</sup>	114.25±13.64 <sup>abc</sup>	115.08 ± 12.35 <sup>abc</sup>
AMF 30 g	64.98 ± 5.46 <sup>ab</sup>	75.08±6.74 <sup>ab</sup>	82.70±6.28 <sup>dab</sup>	91.78±5.26 <sup>abc</sup>	89.43±5.76 <sup>ab</sup>	86.45±6.35 <sup>abc</sup>	86.58 ± 6.21 <sup>abc</sup>
AMF 40 g	83.90±7.18 <sup>ab</sup>	91.65±6.08 <sup>ab</sup>	93.25±7.13 <sup>ab</sup>	93.13±8.05 <sup>abc</sup>	93.70±7.18 <sup>ab</sup>	93.60±5.71 <sup>abc</sup>	98.90 ± 7.02 <sup>abc</sup>
AMF 50 g	70.10±16.37 <sup>ab</sup>	78.38±14.60 <sup>ab</sup>	91.13±11.52 <sup>ab</sup>	97.50±9.30 <sup>abc</sup>	90.78±6.00 <sup>ab</sup>	99.30±9.55 <sup>abc</sup>	107.22 ± 11.06 <sup>abc</sup>
NPK 40 g	103.12±10.08 <sup>ab</sup>	113.32±10.78 <sup>ab</sup>	130.33±8.26 <sup>ab</sup>	132.77±7.48 <sup>abc</sup>	134.90±18.46 <sup>ab</sup>	136.67±8.05 <sup>abc</sup>	137.90 ± 7.60 <sup>abc</sup>
NPK 60 g	113.38± 8.94 <sup>ab</sup>	90.80±8.78 <sup>ab</sup>	121.57±5.48 <sup>ab</sup>	120.03±6.07 <sup>abc</sup>	122.23±6.49 <sup>ab</sup>	124.07±7.34 <sup>abc</sup>	124.47±4.30 <sup>abc</sup>
NPK 80 g	91.75±14.52 <sup>ab</sup>	115.92±10.7 b	106.78±14.33 <sup>ab</sup>	108.50±13.27 <sup>abc</sup>	99.10±6.49 <sup>ab</sup>	99.03±7.34 <sup>abc</sup>	115.12±13.68 <sup>abc</sup>
NPK 100 g	120.85±1.74 <sup>ab</sup>	130.52±20.18 <sup>ab</sup>	133.22±22.37 <sup>ab</sup>	138.45±20.42 <sup>ab</sup>	136.95±19.88 <sup>ab</sup>	137.70±17.62 <sup>ab</sup>	143.05± 20.88 <sup>ab</sup>
CD 200 g	57.35±8.90 <sup>b</sup>	70.90±7.55 <sup>b</sup>	69.75±14.08 <sup>b</sup>	65.15±20.68 <sup>c</sup>	66.60±19.64 <sup>b</sup>	65.85±18.86 <sup>c</sup>	63.70 ± 18.96 <sup>c</sup>
CD 300 g	63.50±14.82 <sup>ab</sup>	73.08±15.03 <sup>b</sup>	78.60±12.95 <sup>ab</sup>	88.35±8.99 <sup>abc</sup>	88.18±6.10 <sup>ab</sup>	87.32±5.19 <sup>abc</sup>	88.73 ± 7.25 <sup>abc</sup>
CD 400 g	64.58±6.31 <sup>ab</sup>	69.98±7.58 <sup>b</sup>	71.63±8.99 <sup>b</sup>	84.40±8.40 <sup>b</sup>	89.68±10.38 <sup>ab</sup>	74.33±5.79 <sup>b</sup>	72.08 ± 7.43 <sup>b</sup>
CD 500 g	79.60±6.42 <sup>ab</sup>	90.25±9.31 <sup>ab</sup>	98.45±14.98 <sup>ab</sup>	102.15±16.03 <sup>abc</sup>	105.68±16.75 <sup>ab</sup>	107.32±16.47 <sup>abc</sup>	110.65 ± 16.54 <sup>abc</sup>
AMFNKPL1 (20+40)g	97.15 ± 13.21 <sup>ab</sup>	97.78±13.24 <sup>ab</sup>	108.93±14.98 <sup>ab</sup>	114.47±10.19 <sup>abc</sup>	116.43±12.93 <sup>ab</sup>	118.97±13.43 <sup>abc</sup>	124.13 ± 13.81 <sup>abc</sup>
AMFNKPL2 (30+60)g	133.22 ± 7.95 <sup>a</sup>	145.42±9.62 <sup>a</sup>	148.92±12.81 <sup>a</sup>	159.60±13.09 <sup>a</sup>	158.87±12.95 <sup>a</sup>	149.70±8.24 <sup>a</sup>	165.73 ± 11.96 <sup>a</sup>
AMFNKPL3 (40+80)g	75.90 ± 7.87 <sup>ab</sup>	106.10±9.38 <sup>ab</sup>	118.70±7.81 <sup>ab</sup>	123.77±7.44 <sup>abc</sup>	121.70±7.83 <sup>ab</sup>	120.45±8.18 <sup>abc</sup>	121.12 ± 9.86 <sup>abc</sup>
AMFNKPL4(50+100)g	97.38 ± 12.54 <sup>ab</sup>	110.45±13.04 <sup>ab</sup>	118.72±12.94 <sup>ab</sup>	124.45±11.46 <sup>abc</sup>	104.42±18.63 <sup>ab</sup>	118.35±11.84 <sup>abc</sup>	110.28 ± 13.41 <sup>abc</sup>
AMFCDL1 (20+200)g	81.73 ± 15.11 <sup>ab</sup>	91.15±13.24 <sup>ab</sup>	99.33±13.77 <sup>ab</sup>	102.55±13.54 <sup>abc</sup>	109.95±14.60 <sup>ab</sup>	111.62±14.23 <sup>abc</sup>	112.72 ± 14.61 <sup>abc</sup>
AMFCDL2(30+300)g	91.05 ± 12.50 <sup>ab</sup>	102.48±12.14 <sup>ab</sup>	108.82±16.98 <sup>ab</sup>	107.12±10.78 <sup>abc</sup>	95.43±4.69 <sup>ab</sup>	100.09±6.01 <sup>abc</sup>	96.28 ± 9.37 <sup>abc</sup>
AMFCDL3 (40+400)g	102.85±15.11 <sup>ab</sup>	112.18±12.87 <sup>ab</sup>	106.48±17.65 <sup>ab</sup>	115.55±18.14 <sup>abc</sup>	118.70±14.52 <sup>ab</sup>	122.50±16.97 <sup>abc</sup>	130.30 ± 24.30 <sup>abc</sup>
AMFCDL4 (50+500) g	66.40 ± 15.33 <sup>ab</sup>	76.23±15.36 <sup>ab</sup>	89.45±11.83 <sup>ab</sup>	91.90± 9.18 <sup>abc</sup>	90.48±8.55 <sup>ab</sup>	89.00 ± 7.56 <sup>abc</sup>	92.83 ± 10.37 <sup>abc</sup>
CDNPKL1 (200+ 40) g	95.23 ± 8.75 <sup>ab</sup>	104.73±9.77 <sup>ab</sup>	113.22±14.85 <sup>ab</sup>	121.00±16.82 <sup>abc</sup>	116.08±20.04 <sup>ab</sup>	118.50±19.56 <sup>abc</sup>	122.00 ± 21.29 <sup>abc</sup>
CDNPKL2 (300+ 60) g	99.33 ± 12.54 <sup>ab</sup>	111.02±12.71 <sup>ab</sup>	126.52±13.31 <sup>ab</sup>	120.40±12.09 <sup>abc</sup>	119.43±9.71 <sup>ab</sup>	123.40±11.44 <sup>abc</sup>	128.10 ± 12.59 <sup>abc</sup>
CDNPKL3 (400+ 80) g	83.45± 22.92 <sup>ab</sup>	99.95±22.53 <sup>ab</sup>	110.88±23.01 <sup>ab</sup>	126.88±14.89 <sup>abc</sup>	123.38±21.13 <sup>ab</sup>	121.80±1.01 <sup>abc</sup>	117.40 ± 21.66 <sup>abc</sup>
CDNPKL4(500+100)g	110.93±14.44 <sup>ab</sup>	122.83±10.99 <sup>ab</sup>	136.37±6.00 <sup>ab</sup>	140.80±4.32 <sup>ab</sup>	115.50±3.88 <sup>ab</sup>	115.37±3.86 <sup>abc</sup>	122.67 ± 1.68 <sup>abc</sup>

Means and standard error of the treatments separated using Tukey's HSD (p < .05). Means with the same letter along the column are not significantly different.

AMF: Arbuscular mycorrhiza fungi;

CD: Cow dung;

WAE: Weeks after Establishment.



**Table 3:** Effects of treatments on cassava leaf area (cm<sup>2</sup>).

Treatment	8WAE	10WAE	12WAE	14WAE	16WAE	18WAE	20WAE
Control	150.23 ± 9.99 <sup>ab</sup>	253.65±62.12 <sup>ab</sup>	280.65 ±67.30 <sup>a</sup>	198.10 ±26.53 <sup>ab</sup>	90.16±5.67 <sup>a</sup>	75.25 ± 17.12 <sup>a</sup>	71.45 ± 7.58 <sup>a</sup>
AMF 20 g	158.90±10.96 <sup>ab</sup>	277.86±39.99 <sup>ab</sup>	146.90±33.29 <sup>a</sup>	164.88±20.30 <sup>ab</sup>	99.39 ± 5.49 <sup>a</sup>	98.79 ± 11.30 <sup>a</sup>	57.99 ± 11.01 <sup>a</sup>
AMF 30 g	118.91±14.45 <sup>ab</sup>	164.61±29.83 <sup>ab</sup>	146.96±25.50 <sup>a</sup>	135.01 ± 5.23 <sup>ab</sup>	84.34 ± 7.13 <sup>a</sup>	65.27 ± 13.68 <sup>a</sup>	48.41 ± 17.48 <sup>a</sup>
AMF 40 g	126.06 ± 6.70 <sup>ab</sup>	240.28±15.85 <sup>ab</sup>	170.95±25.77 <sup>a</sup>	119.51±13.69 <sup>b</sup>	73.40±4.60 <sup>a</sup>	46.49 ± 6.35 <sup>a</sup>	74.43 ±7.66 <sup>a</sup>
AMF 50 g	132.01±14.00 <sup>ab</sup>	215.10±28.02 <sup>ab</sup>	181.17±10.42 <sup>a</sup>	157.34 ± 8.30 <sup>ab</sup>	95.58 ± 9.99 <sup>a</sup>	68.06 ± 16.36 <sup>a</sup>	74.77 ± 8.82 <sup>a</sup>
NPK 40 g	152.34±23.37 <sup>ab</sup>	323.27±41.12 <sup>ab</sup>	283.57±56.37 <sup>a</sup>	178.63±54.82 <sup>ab</sup>	109.57 ± 5.26 <sup>a</sup>	80.21 ± 18.31 <sup>a</sup>	85.42 ± 3.12 <sup>a</sup>
NPK 60 g	174.94±26.22 <sup>ab</sup>	334.26±9.38 <sup>ab</sup>	225.37±46.52 <sup>a</sup>	140.24±35.05 <sup>ab</sup>	87.65 ± 4.53 <sup>a</sup>	95.16±4.99 <sup>a</sup>	74.11 ± 4.50 <sup>a</sup>
NPK 80 g	140.68 ± 6.68 <sup>ab</sup>	276.43±61.43 <sup>ab</sup>	245.76±69.22 <sup>a</sup>	208.81±49.33 <sup>ab</sup>	69.34± 11.65 <sup>a</sup>	45.72±17.23 <sup>a</sup>	43.62 ± 12.44 <sup>a</sup>
NPK 100 g	180.17±20.54 <sup>ab</sup>	341.05±16.60 <sup>ab</sup>	307.98±49.49 <sup>a</sup>	245.45±40.19 <sup>ab</sup>	85.77 ± 9.87 <sup>a</sup>	71.65 ± 23.80 <sup>a</sup>	72.75 ± 23.57 <sup>a</sup>
CD 200 g	120.87±16.42 <sup>ab</sup>	166.22±28.63 <sup>ab</sup>	108.36±32.44 <sup>a</sup>	125.38±24.52 <sup>ab</sup>	96.77 ± 15.70 <sup>a</sup>	68.66 ± 9.95 <sup>a</sup>	88.69 ± 15.94 <sup>a</sup>
CD 300 g	138.59±28.29 <sup>ab</sup>	217.11±50.51 <sup>ab</sup>	193.75±26.94 <sup>a</sup>	149.18±11.41 <sup>ab</sup>	96.88 ± 5.02 <sup>a</sup>	85.11 ± 14.20 <sup>a</sup>	48.91 ± 11.60 <sup>a</sup>
CD 400 g	110.31± 10.26 <sup>b</sup>	163.34±27.45 <sup>b</sup>	127.84±16.86 <sup>a</sup>	131.16 ±10.94 <sup>ab</sup>	88.70 ± 11.65 <sup>a</sup>	66.74 ± 14.41 <sup>a</sup>	63.44 ± 19.18 <sup>a</sup>
CD 500 g	152.62±18.10 <sup>ab</sup>	217.10±36.35 <sup>ab</sup>	203.99±47.20 <sup>a</sup>	134.81 ±19.10 <sup>ab</sup>	91.51 ± 13.93 <sup>a</sup>	65.51 ± 7.07 <sup>a</sup>	102.22 ± 32.77 <sup>a</sup>
AMFNKPL1(20+40) g	186.48±13.92 <sup>ab</sup>	251.47±39.32 <sup>ab</sup>	182.11±20.21 <sup>a</sup>	169.85±13.38 <sup>ab</sup>	71.92 ± 2.46 <sup>a</sup>	61.71 ± 18.25 <sup>a</sup>	82.02 ± 2.31 <sup>a</sup>
AMFNKPL2 (30+60) g	227.50±30.98 <sup>a</sup>	405.18±34.67 <sup>a</sup>	349.06±39.69 <sup>a</sup>	341.50 ± 1.45 <sup>a</sup>	61.8 ± 39.0 <sup>a</sup>	59.83 ± 28.26 <sup>a</sup>	61.38 ± 16.74 <sup>a</sup>
AMFNKPL3 (40+80) g	161.87±12.57 <sup>ab</sup>	284.93±38.60 <sup>ab</sup>	215.56±33.08 <sup>a</sup>	184.55 ± 9.70 <sup>ab</sup>	83.04 ± 6.23 <sup>a</sup>	61.36 ± 15.47 <sup>a</sup>	71.59 ± 11.56 <sup>a</sup>
AMFNKPL4 (50+100) g	159.06±1.98 <sup>ab</sup>	228.64±53.12 <sup>ab</sup>	233.76±39.67 <sup>a</sup>	212.17±16.16 <sup>ab</sup>	113.57 ± 6.70 <sup>a</sup>	49.92 ± 14.32 <sup>a</sup>	63.39 ± 19.99 <sup>a</sup>
AMFCDL1 (20+200) g	131.67±21.99 <sup>ab</sup>	212.86±59.60 <sup>ab</sup>	175.14±64.11 <sup>a</sup>	188.33±66.55 <sup>ab</sup>	104.90 ± 7.60 <sup>a</sup>	89.51 ± 14.12 <sup>a</sup>	62.28 ± 12.71 <sup>a</sup>
AMFCDL2 (30+300) g	175.92±20.74 <sup>ab</sup>	227.97±51.84 <sup>ab</sup>	205.86±52.05 <sup>a</sup>	137.08±28.09 <sup>ab</sup>	76.49 ± 8.02 <sup>a</sup>	58.97 ± 6.78 <sup>a</sup>	33.44 ± 8.43 <sup>a</sup>
AMFCDL3 (40+400) g	172.03±14.94 <sup>ab</sup>	282.84±57.02 <sup>ab</sup>	237.23±69.31 <sup>a</sup>	217.81±46.00 <sup>ab</sup>	81.87 ± 21.65 <sup>a</sup>	95.27 ± 9.39 <sup>a</sup>	73.75 ± 13.04 <sup>a</sup>
AMFCDL4 (50+500)g	138.90±20.62 <sup>ab</sup>	208.48±37.82 <sup>ab</sup>	190.26±33.90 <sup>a</sup>	132.88±12.73 <sup>ab</sup>	79.42 ± 5.86 <sup>a</sup>	64.16 ± 10.15 <sup>a</sup>	60.45 ± 4.87 <sup>a</sup>
CDNPKL1 (200+40) g	158.05 ± 6.91 <sup>ab</sup>	308.21±19.76 <sup>ab</sup>	242.42±49.45 <sup>a</sup>	205.83±62.75 <sup>ab</sup>	83.41 ± 14.06 <sup>a</sup>	70.43 ± 13.35 <sup>a</sup>	49.98 ± 16.07 <sup>a</sup>
CDNPKL2 (300+60) g	154.62±19.24 <sup>ab</sup>	322.37±60.63 <sup>ab</sup>	272.03±59.67 <sup>a</sup>	150.32±10.35 <sup>ab</sup>	96.22 ± 19.71 <sup>a</sup>	111.28±24.52 <sup>a</sup>	101.01 ± 18.26 <sup>a</sup>
CDNPKL3 (400+80) g	151.10±52.28 <sup>ab</sup>	280.98±56.74 <sup>ab</sup>	287.94±72.75 <sup>a</sup>	224.69±31.31 <sup>ab</sup>	104.38 ± 3.22 <sup>a</sup>	110.80±15.63 <sup>a</sup>	90.44 ± 4.04 <sup>a</sup>
CDNPKL4 (500+100) g	212.93±25.73 <sup>ab</sup>	323.77±64.52 <sup>ab</sup>	353.08±44.78 <sup>a</sup>	247.58±20.48 <sup>ab</sup>	85.15 ± 2.38 <sup>a</sup>	84.71 ± 16.00 <sup>a</sup>	82.76 ± 12.94 <sup>a</sup>

Means with the same letter along the column are not significantly different.

AMF: Arbuscular mycorrhiza fungi;

CD: Cow dung;

WAE: Weeks After Establishment.

**Table 4:** Effects of treatments on cassava stems girth (mm).

Treatment	10WAE	12WAE	14WAE	16WAE	18WAE	20WAE
Control	14.83 ± 1.89 <sup>ab</sup>	17.13 ± 0.62 <sup>ab</sup>	17.08 ± 0.44 <sup>ab</sup>	17.48 ± 0.60 <sup>ab</sup>	17.48 ± 0.64 <sup>ab</sup>	17.10 ± 0.54 <sup>ab</sup>
AMF 20 g	14.97 ± 1.53 <sup>ab</sup>	15.40 ± 1.75 <sup>ab</sup>	16.24 ± 1.75 <sup>ab</sup>	15.87 ± 1.61 <sup>ab</sup>	15.67 ± 2.17 <sup>ab</sup>	16.16 ± 1.29 <sup>ab</sup>
AMF 30 g	12.55 ± 0.88 <sup>b</sup>	12.80 ± 0.69 <sup>ab</sup>	12.46 ± 0.57 <sup>ab</sup>	13.10 ± 0.60 <sup>ab</sup>	12.78 ± 0.80 <sup>ab</sup>	12.76 ± 0.63 <sup>ab</sup>
AMF 40 g	15.40 ± 1.06 <sup>ab</sup>	15.06 ± 0.97 <sup>ab</sup>	15.36 ± 0.57 <sup>ab</sup>	15.20 ± 1.05 <sup>ab</sup>	15.37 ± 0.38 <sup>ab</sup>	15.39 ± 0.60 <sup>ab</sup>
AMF 50 g	13.12 ± 1.92 <sup>ab</sup>	13.81 ± 1.90 <sup>ab</sup>	14.85 ± 1.20 <sup>ab</sup>	13.94 ± 1.08 <sup>ab</sup>	14.68 ± 1.28 <sup>ab</sup>	13.66 ± 1.04 <sup>ab</sup>
NPK 40 g	17.15 ± 1.22 <sup>ab</sup>	17.90 ± 0.98 <sup>ab</sup>	17.99 ± 2.15 <sup>ab</sup>	18.45 ± 1.23 <sup>ab</sup>	18.66 ± 1.16 <sup>ab</sup>	18.09 ± 1.50 <sup>ab</sup>
NPK 60 g	17.01 ± 0.86 <sup>ab</sup>	17.04 ± 0.43 <sup>ab</sup>	17.92 ± 0.70 <sup>ab</sup>	18.13 ± 0.65 <sup>ab</sup>	18.07 ± 0.67 <sup>ab</sup>	17.70 ± 0.14 <sup>ab</sup>
NPK 80 g	15.53 ± 1.51 <sup>ab</sup>	14.78 ± 1.45 <sup>ab</sup>	15.09 ± 1.27 <sup>ab</sup>	13.42 ± 0.75 <sup>ab</sup>	14.69 ± 1.37 <sup>ab</sup>	14.94 ± 1.25 <sup>ab</sup>
NPK 100 g	17.33 ± 1.98 <sup>ab</sup>	17.25 ± 2.05 <sup>ab</sup>	17.53 ± 2.25 <sup>ab</sup>	18.14 ± 2.05 <sup>ab</sup>	17.39 ± 2.09 <sup>ab</sup>	17.51 ± 2.21 <sup>ab</sup>
CD 200 g	11.83 ± 1.00 <sup>b</sup>	12.32 ± 1.44 <sup>b</sup>	10.72 ± 2.48 <sup>b</sup>	10.92 ± 2.26 <sup>b</sup>	10.70 ± 2.18 <sup>b</sup>	10.28 ± 2.19 <sup>b</sup>
CD 300 g	13.14 ± 1.21 <sup>ab</sup>	13.95 ± 1.02 <sup>ab</sup>	13.85 ± 1.26 <sup>ab</sup>	14.46 ± 0.68 <sup>ab</sup>	14.46 ± 0.57 <sup>ab</sup>	14.39 ± 0.68 <sup>ab</sup>
CD 400 g	13.02 ± 1.29 <sup>ab</sup>	12.47 ± 1.28 <sup>b</sup>	13.32 ± 1.66 <sup>ab</sup>	13.01 ± 1.43 <sup>ab</sup>	12.45 ± 1.65 <sup>b</sup>	12.82 ± 1.53 <sup>ab</sup>
CD 500 g	14.99 ± 0.63 <sup>ab</sup>	15.01 ± 0.83 <sup>ab</sup>	15.84 ± 1.17 <sup>ab</sup>	16.16 ± 1.28 <sup>ab</sup>	15.93 ± 1.19 <sup>ab</sup>	15.14 ± 1.64 <sup>ab</sup>
AMFNKPL1(20+40) g	16.57 ± 2.38 <sup>ab</sup>	16.31 ± 2.05 <sup>ab</sup>	16.51 ± 2.13 <sup>ab</sup>	17.40 ± 1.63 <sup>ab</sup>	16.65 ± 1.70 <sup>ab</sup>	16.42 ± 1.99 <sup>ab</sup>
AMFNKPL2 (30+60) g	20.72 ± 1.61 <sup>a</sup>	20.14 ± 1.43 <sup>a</sup>	20.76 ± 1.80 <sup>a</sup>	21.08 ± 1.89 <sup>a</sup>	21.19 ± 1.56 <sup>a</sup>	20.87 ± 1.93 <sup>a</sup>
AMFNKPL3 (40+80) g	16.27 ± 1.70 <sup>ab</sup>	17.61 ± 1.00 <sup>ab</sup>	17.69 ± 0.87 <sup>ab</sup>	17.86 ± 0.88 <sup>ab</sup>	17.71 ± 0.69 <sup>ab</sup>	17.84 ± 0.69 <sup>ab</sup>
AMFNKPL4 (50+100) g	15.77 ± 1.06 <sup>ab</sup>	16.54 ± 1.18 <sup>ab</sup>	16.53 ± 1.06 <sup>ab</sup>	16.71 ± 0.99 <sup>ab</sup>	17.06 ± 1.17 <sup>ab</sup>	16.16 ± 1.24 <sup>ab</sup>
AMFCDL1 (20 + 200) g	14.09 ± 1.68 <sup>ab</sup>	13.90 ± 1.77 <sup>ab</sup>	15.34 ± 1.55 <sup>ab</sup>	14.73 ± 2.12 <sup>ab</sup>	14.32 ± 1.63 <sup>ab</sup>	14.50 ± 1.61 <sup>ab</sup>
AMFCDL2 (30 + 300) g	15.10 ± 1.15 <sup>ab</sup>	15.27 ± 0.80 <sup>ab</sup>	14.85 ± 1.08 <sup>ab</sup>	14.63 ± 0.79 <sup>ab</sup>	15.75 ± 0.72 <sup>ab</sup>	14.57 ± 1.22 <sup>ab</sup>
AMFCDL3 (40 + 400) g	16.09 ± 1.08 <sup>ab</sup>	15.62 ± 1.94 <sup>ab</sup>	15.80 ± 1.82 <sup>ab</sup>	16.54 ± 1.86 <sup>ab</sup>	16.38 ± 1.73 <sup>ab</sup>	15.78 ± 2.31 <sup>ab</sup>
AMFCDL4 (50+500)g	13.67 ± 1.64 <sup>ab</sup>	14.11 ± 1.20 <sup>ab</sup>	13.64 ± 0.69 <sup>ab</sup>	13.79 ± 0.68 <sup>ab</sup>	13.82 ± 0.78 <sup>ab</sup>	13.86 ± 0.94 <sup>ab</sup>
CDNPKL1 (200+40)g	16.54 ± 0.90 <sup>ab</sup>	16.91 ± 1.46 <sup>ab</sup>	17.17 ± 1.74 <sup>ab</sup>	17.41 ± 1.86 <sup>ab</sup>	17.52 ± 1.97 <sup>ab</sup>	17.58 ± 2.15 <sup>ab</sup>
CDNPKL2 (300+60) g	16.65 ± 1.22 <sup>ab</sup>	17.40 ± 1.23 <sup>ab</sup>	16.69 ± 0.81 <sup>ab</sup>	16.74 ± 0.85 <sup>ab</sup>	17.75 ± 2.46 <sup>ab</sup>	16.71 ± 0.97 <sup>ab</sup>
CDNPKL3 (400+80) g	16.27 ± 2.60 <sup>ab</sup>	16.68 ± 2.44 <sup>ab</sup>	15.88 ± 2.43 <sup>ab</sup>	17.38 ± 2.15 <sup>ab</sup>	16.97 ± 2.09 <sup>ab</sup>	17.34 ± 2.15 <sup>ab</sup>
CDNPKL4 (500+100) g	18.32 ± 1.21 <sup>ab</sup>	18.47 ± 0.52 <sup>ab</sup>	19.29 ± 0.65 <sup>a</sup>	14.09 ± 2.11 <sup>ab</sup>	16.76 ± 2.94 <sup>ab</sup>	15.99 ± 2.83 <sup>ab</sup>

Means and standard error of the treatments separated using Tukey's HSD (p < .05). Means with the same letter along the column are not significantly different.

AMF: Arbuscular mycorrhiza fungi;

CD: Cow dung;

WAE: Weeks after Establishment.


## Discussion

Several researchers have demonstrated the beneficial effects of mycorrhizal inoculation on growth and yield of cassava Douds et al. [46]; Carretero et al. [16]; Séry et al. [47]. However, the combination of AMF and organic/inorganic fertilizers has not been adequately investigated. The results of this study suggest that *G. deserticola* inoculation in combination with inorganic fertilizer application at the rate of (30g and 60g respectively) has beneficial effects on all the growth parameters studied. In this study, leaf chlorophyll content was a good indicator of growth and this was reflected in the yield of the cassava plants studied. Ekanayake et al. [48], reported that soil inoculated with 10g of *Glomus clarum* and *G. mosseae* enhanced chlorophyll production

in young cassava plants, with the former species supporting more chlorophyll synthesis than the later. Howeler et al. [36], reported an increase in cassava growth under different combinations of P input with AMF as opposed to the sole application of AMF. Most of the soils supporting cassava cultivation in the south-western region of the Nigeria are P deficient thereby underscoring the need for strategies for sustainable soil fertility techniques Salami and Sangoyomi [49]. The physico-chemical properties of the experimental soil do not differ considerably from those reported by these authors.

Arbuscular mycorrhiza fungi are crucial component of the soil ecosystem that enhances nutrient uptake and absorption Bolan, [14]. The increase in growth attributes recorded in this study

could be as a result of these processes. Similarly, Sieverding and Toro [30], opined that mycorrhiza inoculation enhanced nutrient uptake in the shoot of cassava compared to non-inoculated plants. In this study, *G. deserticola* inoculation enhanced the growth and yield of cassava therefore supports reduced inorganic fertilizers inputs. This result compares favourably with previous studies on the effects of mycorrhizal inoculations on cassava growth. Ceballos et al. [37], showed that *Rhizophagus irregularis* inoculation produced 20% increase in cassava yield and 50% reduction in phosphate fertilizer. Similarly Sridevi and Ramakrishnan [50], studying the response of cassava to *Glomus fasciculatum* inoculation at increased NPK levels reported that yield attributes, like number of tubers, tuber yield were optimal under increased NPK and AMF application. In the present study, inoculation of *G. deserticola* enhanced the performance and yield the cassava plants studied.




**Figure 1:** Effects of treatment on the cassava leaf chlorophyll content (ml/g).

**AMF:** Arbuscular Mycorrhiza Fungi

**CD:** Cow Dung

**NPK:** Inorganic Nitrogen: Phosphorus: Potassium fertilizer (15:15:15).



**Figure 2:** Effects of treatments on cassava yield.

**AMF:** Arbuscular Mycorrhiza Fungi

**CD:** Cow Dung

**NPK:** Inorganic Nitrogen: Phosphorus: Potassium fertilizer (15:15:15).

## Conclusion

The cassava variety TME 419 used in this study responded positively to *G. deserticola* inoculation in combination with inorganic fertilization. This finding indicates the potentials of arbuscular mycorrhiza fungi as a biological agent for sustainable agriculture.

## References

1. Montagnac JA, Davis CR, Tanumihardjo SA (2009) Nutritional Value of Cassava for use as a Staple Food and Recent Advances for Improvement Comprehensive Reviews. In Food Science And Food Safety 8(3): 181-194.
2. FAOSTAT (2016) Statistical Database of the Food and Agriculture Organization of the United Nations. USA.
3. El-Sharkawy MA (2006) International research on cassava photosynthesis, productivity, eco-physiology, and responses to environmental stresses in the tropics Photosynthetica 44 (4): 481-512.
4. Edmeades DC (2003) The long-term effects of manures and fertilisers on soil productivity and quality: a review Nutrient Cycling in Agroecosystems 66(2): 165-180.
5. Diacono M, Montemurro F (2010) Long-term effects of organic amendments on soil fertility A review Agron Sustain Dev 30(2): 401-422.
6. Šimon, T Czakó A (2014) Influence of long-term application of organic and inorganic fertilizers on soil properties Plant Soil Environ 60 (7): 314-319.
7. Kotschi J (2013) A Soiled Reputation Adverse impacts of mineral fertilizers in tropical agriculture AGRECOL-Association for Agriculture and Ecology, Heinrich Böll Stiftung, WWF, Germany, p.1-57.
8. Mulvaney RL, Khan SA, Ellsworth TR (2009) Synthetic Nitrogen Fertilizers Deplete Soil Nitrogen: A Global Dilemma for Sustainable Cereal Production. J Environ Qual 38(6): 2295-2314.
9. Ayoola OT, Adeniyi ON (2006) Influence of poultry manure and NPK fertilizer on yield and yield components of crops under different cropping systems in south west Nigeria. African Journal of Biotechnology 5 (15): 1386-1392.
10. Ojeniyi SO, Adejoro SA, Ikotun O, Amusan O (2012) Soil and plant nutrient composition, growth and yield of cassava as influenced by integrated application of NPK fertilizer and poultry manure New York Science Journal 5(9): 62-68.
11. Islami T, Guritno B, Utomo WH (2011) Performance of cassava (*Manihot esculenta* Crantz) based cropping systems and associated soil quality changes in the degraded tropical uplands of East Java, Indonesia. Journal of Tropical Agriculture 49(1-2): 31-39.
12. Smith SE, Facelli E, Pope S, Smith FA (2010) Plant performance in stressful environments. Interpreting new and established knowledge of the roles of arbuscular mycorrhizas. Plant and Soil 326(1-2): 3-20.
13. Böhm J, Hock B (1997) Mycorrhizae: Endomycorrhizae Progress in Botany 58: 555-594.

14. Bolan NS (1991) A critical review on the role of mycorrhizal fungi in the uptake of Phosphorus by plants Plant and Soil 134: (2): 189-207.
15. Smith SE, Read DJ (2008) The mycorrhizal symbiosis. San Diego, Academic Press, USA.
16. Carretero CL, Cantos M, Garcia JL, Azcon R, Troncoso A (2009) Growth response of micro propagated cassava clones as affected by Glomus intraradices colonization. Journal of Plant Nutrition 32(2): 261-273.
17. Qiangsheng W, Renxue X, Zhengjia H (2006) Effect of Arbuscular Mycorrhiza on the Drought Tolerance of Poncirus trifoliata Seedlings. Frontiers of Forestry in China 1(1): 100-104.
18. Ruiz-Lozano JM, Azcon R, Gomez M (1995) Effects of Arbuscular-Mycorrhizal Glomus Species on Drought Tolerance: Physiological and Nutritional Plant Responses. Appl Environ Microbiol 61(2): 456-460.
19. Li H, Ye ZH, Chan WF, Chen XW, Wu FY, et al. (2011) Can arbuscular mycorrhizal fungi improve grain yield, as uptake and tolerance of rice grown under aerobic conditions? Environ Pollut 159(10): 2537-2545.
20. Cely MVT, de Oliveira AG, de Freitas VF, de Luca MB, Barazetti AR, et al. (2016) Inoculant of Arbuscular Mycorrhizal Fungi (Rhizophagus clarus) Increase Yield of Soybean and Cotton under Field Conditions. Front Microbiol 7: 1-720.
21. Tchabi A, Hountondji FCC, Ogunsola B, Lawouin L, Coyne D, et al. (2016) The Influence of Arbuscular Mycorrhizal Fungi Inoculation on Micro-Propagated Hybrid Yam (Dioscorea spp ) Growth and Root Knot Nematode (Meloidogyne spp ) Suppression. Int J Curr Microbiol App Sci 5(10): 267-281.
22. Howeler RH (1991) Long-term effect of cassava cultivation on soil productivity. Field Crops Research 26: 1-18.
23. Habte M, Byappanahalli MN (1994) Dependency of cassava (Manihot esculenta Crantz) on vesicular-arbuscular mycorrhizal fungi Mycorrhiza 4: 241-245.
24. Tawaraya K (2003) Arbuscular Mycorrhizal Dependency of Different Plant Species and Cultivars. Soil Sci Plant Nutl 49(5): 655-668.
25. Burns AE, Gleadow RM, Zacarias AM, Cuambe CE, Miller RE, Cavagnaro TR, et al. (2012) Variations in the chemical composition of cassava (Manihot esculenta Crantz) leaves and roots as affected by genotypic and environmental variation. J. Agric. Food Chem. 60(19): 4946-4956.
26. Straker CJ, Hilditch AJ, Rey MEC (2010) Arbuscular mycorrhizal fungi associated with cassava (Manihot esculenta Crantz) in South Africa. South African Journal of Botany 76(1): 102-111.
27. Bi Voko DRR, Nandjui J, Sery JMD, FotsoB, Amoa JA, et al (2013) Abundance and diversity of Arbuscular mycorrhizal fungal (AMF) communities associated with cassava (Manihot esculenta Crantz) rhizosphere in Abengourou, East Côte d'Ivoire Journal of Ecology and the Natural Environment 5(11): 360-370.
28. dos Santos Heberle E, de Armas RD, Heberle DA, Stürmer SL, Peruch LAM, et al. (2015) Occurrence and Structure of Arbuscular Mycorrhizal Fungal Communities in Cassava after Cultivation of Cover Crops as Observed by the "PCR-DGGE" Technique R Bras Ci Solo 39(5): 1292-1301.
29. Begoude DAB, Sarr PS, Mpon TLY, Owona D A, Kapeua MN, et al (2016) Composition of arbuscular mycorrhizal fungi associated with cassava (Manihot esculenta Crantz) cultivars as influenced by chemical fertilization and tillage in Cameroon J Appl Biosci 98: 9270-9283.
30. Sieverding E, Toro ST (1989) Effect of mixing VAM inoculum with fertilizers on cassava nutrition and VAM fungal association. Agriculture Ecosystems and Environment Netherlands 29(1-4): 397-401.
31. Ekanayake IJ, Oyetunji OJ, Osonubi O, Lyasse O (2004) The effects of arbuscular mycorrhizal fungi and water stress on leaf chlorophyll production of cassava (Manihot esculenta Crantz) Food, Agriculture & Environment 2 (2): 190-196.
32. Okon IE, Solomon MG, Osonubi O (2010) The Effects of Arbuscular Mycorrhizal Fungal Inoculation and Mulch of Contrasting Chemical Composition on the Yield of Cassava under Humid Tropical Conditions. Scientific WorldJournal. 10: 505-511.
33. Carretero CL, Cantos M, García JL, Azcón R, Troncoso A (2008) Arbuscular-Mycorrhizal Contributes to Alleviation of Salt Damage in Cassava Clones Journal of Plant Nutrition 31(5): 959-971.
34. Salami AO, Odebo AC, Osonubi O (2005) The use of Arbuscular Mycorrhiza (AM) as a source of yield increase in sustainable alley cropping system. Archives of Agronomy and Soil Science August 51(4): 385-390.
35. Howeler RH, Edwards DG, Asher CJ (1979) The effect of soil sterilization and mycorrhizal inoculation on the growth, nutrient uptake and critical P concentration of cassava. Intern Symp Trop Root and Tuber Crops, Manila: 519-537.
36. Ceballos I, Ruiz M, Fernández C, Peña R, Rodríguez A, et al. (2013) The in vitro mass-produced model mycorrhizal fungus, Rhizophagus irregularis, significantly increases yields of the globally important food security crop cassava. PLoS ONE 8(8): e70633.
37. Azcón-Aguilar C, Cantos M, Troncoso A, Barea JM (1997) Beneficial effect of arbuscular mycorrhizas on acclimatization of micropropagated cassava plantlets. Scientia Horticulturae 72: 63-71.
38. Gee GW, Bauders A (1986) Particle size analysis: In: Khite A (Eds.), Methods of soil analysis, Part 1 (2<sup>nd</sup> edn), Agronomy Monograph 9, USA.
39. Nelson DW, Sommers LE (1996) Total carbon, organic carbon and organic matter In: Sparks DL (Eds.), Methods of soil analysis, Part 3 Chemical Methods, SSSA Book Series No 5 ASA & SSSA, USA, pp.961-1010.
40. Bremner JM (1996) N-total In: Sparks DL (Eds.), Methods of soil analysis, Part 3, SSSA and ASA, Soil Science Society of America, USA, pp. 1085-1121.
41. Bray RH, Kurtz LT (1945) Determination of total, organic and available form of phosphorus in soils. Soil Science Society of American Journal 59(1): 39-46.
42. Jackson ML (1962) Soil chemical analysis. Constable and Company Limited, London, England.
43. Mathias L, Kabambe VH (2015) Potential to increase cassava yields through cattle manure and fertilizer application: Results from Bunda College, Central Malawi African Journal of Plant Science 9(5): 228-234.



44. Okon IE, Solomon MG, Osonubi O (2010) The Effects of Arbuscular Mycorrhizal Fungal Inoculation and Mulch of Contrasting Chemical Composition on the Yield of Cassava under Humid Tropical Conditions. *Scientific WorldJournal*. 10: 505-511.
45. Adepetu JA (1996) Interpretation of soil test data: Simple Soil, water and plant testing techniques for soil resource management. In: Depetu JA & Nabhan H (Eds. ), *Proceedings of a training course, Nigeria*, pp 1-158.
46. Douds DD, Nagahashi G, Pfeffer PE, Kayser WM, Reider C (2005) On-farm production and utilization of arbuscular mycorrhizal fungus inoculum. *Canadian Journal of Plant Sciences* 85: 15-21.
47. Séry DJM, Kouadjo ZGC, Voko BRR Zézé A (2016) Selecting Native Arbuscular Mycorrhizal Fungi to Promote Cassava Growth and Increase Yield under Field Conditions. *Frontiers in Microbiology* 7: 2063.
48. Ekanayake IJ, Oyetunji OJ, Osonubi O, Lyasse O (2004) The effects of arbuscular mycorrhizal fungi and water stress on leaf chlorophyll production of cassava (*Manihot esculenta* Crantz) *Food, Agriculture & Environment* 2 (2): 190-196.
49. Salami BT, Sangoyomi TE (2013) Soil fertility status of cassava fields in South Western Nigeria. *American Journal of Experimental Agriculture* 3(1): 152-164.
50. Sridevi S, Ramakrishnan K (2013) Effect of inoculation with NPK fertilizer and arbuscular mycorrhizal fungi on growth and yield of cassava. *International Journal of Development Research* 3(9): 46-50.