

Obesity Correlates with Neutrophilia

Abstract

Background: Obesity is a global health problem imposing risk for many serious diseases. Lately, obesity has been described as a state of low-grade inflammation with leukocytes playing a major role. Mild leukocytosis was noticed in obese patients with no co morbidities. This study aimed at defining the association between obesity and leukocytosis.

Methods: In this retrospective study, we reviewed files and medical history of 281 patients. Body Mass Index (BMI) was calculated and differential leukocyte was recorded. Individuals with obesity related conditions like hypertension, diabetes, hyperlipidemia or malignancies were excluded. Data analysis was performed to assess the relationship between variables.

Results: Among the 281 participants included in the study the BMI was used as a determinant of obesity with a cut-off point of 30 (127 were non-obese and 154 were obese). There was a statistically significant difference in the white blood cells (WBC) count among the two groups (p-value <0.001). In addition, 88.9% of those with leukocytosis were obese, while 11.1% non-obese. There is a positive correlation between the WBC count and the BMI (r=0.391, p-value <0.001). Neutrophilia is the rationale behind leukocytosis with the mean of $8.3 \times 10^9 \pm 3$.

Conclusion: The positive correlation between obesity and leukocytosis proves the relationship but does not establish causation. Although most obese patients in this study had a normal WBC count, their counts were higher than the non-obese group due to prominent neutrophilia. The association highlighted will have an impact on patients during the pre-operative work up for bariatric surgery, as neutrophilia is a risk factor for metabolic and cardiovascular complications requiring detection and treatment to minimize this risk. The incidental detection of neutrophilia in obese patients is not yet rationalized and requires further research

Keywords: Obesity; Neutrophilia; Leukocytosis; Bariatric surgery

Research Article

Volume 3 Issue 2 - 2016

Salma Sait*, Alqassas M, Othman S, Shihon SB, Alqalayta L, Alqusair S and Mohamad Qari

Department of Hematology, King Abdulaziz University, Saudi Arabia

***Corresponding author:** Salma Sait, Hematology dept, College of Medicine, King Abdulaziz University, P O Box 80215, Jeddah 21589, Saudi Arabia, Tel: +966542226281; Email: salmahsait@yahoo.com

Received: November 10, 2016 | **Published:** December 19, 2016

Abbreviations: BMI: Body Mass Index; WBC: White Blood Cells; CRP: C-Reactive Protein; DM: Diabetes Mellitus; HTN: Hypertension

Introduction

Obesity is a global issue that threatens millions of people around the world in both developing and developed countries. Prevalence of obesity has duplicated two fold since 1980 and it is predicted that by 2030 it will affect more than one billion people [1,2]. Obesity is a major risk factor for type 2 diabetes, coronary heart disease, hypertension, atherosclerosis, arterial thrombosis, heart failure and sudden cardiac death [3]. According to the WHO, obesity is defined as body mass index (BMI) of 30, while a value greater than 40 is considered extreme or morbid obesity. Recently, obesity has been described as a state of low-grade inflammation [4-6], with leukocytes playing an important role in this state. Leukocytosis is an increase in the total number of White blood cell (WBC) count with a value greater than $11,000/\text{mm}^3$ ($11 \times 10^9/\text{L}$) due to any cause such as infection, inflammation, allergic reaction, malignancy, hereditary disorders, or other miscellaneous causes [7].

A Study conducted in Turkey reported that WBC, neutrophils and lymphocyte counts are significantly affected by BMI status [8]. In China, it was found that levels of WBCs and C-reactive protein (CRP) in morbidly obese patients who underwent surgical weight loss significantly decreased after the surgery [9].

Furthermore, a relation between leukocytosis and obesity-related co morbidities has been established. A study conducted by Ryder et al, showed that a high WBC count in obese patients is associated with insulin resistance [10]. Another cross-sectional study also concluded that an elevated WBC count is linked to the prevalence and future development of metabolic syndrome in the young population [11]. However, to what extent a high WBC count in obese patients can be considered alarming is still unknown. Most of the studies in this area focused on leukocytosis in obese patients with obesity-related comorbidities; little work has been done regarding leukocytosis in asymptomatic obese individuals. We therefore aim in this paper, to study the level of correlation between leukocytosis and obesity in a cohort of healthy and obese subjects.

Material and Methods

This retrospective study was conducted during the month of June 2016 in King Abdulaziz University Hospital, Jeddah, Saudi Arabia. It was approved by the Research Ethics committee of KAU. Medical records of 326 adult patients were reviewed; 131 of whom were obese subjects going for bariatric surgery while 195 were volunteer donors. The sample size was calculated using the following equation, with confidence interval of 95% and margin of error=6%.

$$Sample\ Size = \frac{z^2 \times p(1-p)}{1 + \frac{z^2 \times p(1-p)}{e^2 N}}$$

Information obtained from the medical records included age, gender, height, weight and differential leukocyte count. Body mass index was calculated by dividing (weight in kg) over (height in m²); patients with BMI≥30 were considered obese while those with BMI<30 were non-obese according to WHO’s classification of body mass index (BMI). Leukocytosis was determined the total WBC count ≥11.5mmol/L as stated by the KAUH hematological laboratory. Patients of both sexes ages 18 and above, were included in the study. Exclusion criteria included the following: Diabetes Mellitus (DM), hypertension (HTN), hyperlipidemia, metabolic syndrome, cardiovascular diseases, malignancies and recent infection (body temperature >38°C). In addition to diseases or drugs known to influence the immune system and secondary obesity (hypothyroidism, Cushing syndrome), SPSS version 21 was used for statistical analysis. The relation between the two variables was studied using chi-square and the one-way ANOVA test. A p-value <0.05 was considered as significant.

Results

The records of 326 patients from KAUH were reviewed and analyzed. 45 patients were excluded because they had associated co morbidities including hypertension, diabetes, hyperlipidemia, thyroid disease and malignancies. Of the 281 patients included in the study, 127 were non-obese with BMI<30 and 154 were obese with BMI>30. The characteristics of the participants studied are shown in Table 1. As shown in Figure 1&2, there is a positive correlation between BMI and Total WBC count-Pearson’s correlation coefficient= 0.391, p-value <0.001. Leukocytosis was found in 18 of the participants studied; 88.9 % of whom were obese and only 11.1% were non obese. To assess the effect of severity of obesity with leukocytosis, we further grouped the participants according to their BMI as shown in Table 2. When comparing the means of WBC count between the groups using the one-way ANOVA test, a significant difference was found as shown in Figure 3. We also found that BMI of 35 or higher is significantly associated with leukocytosis (p-value=0.01). While measuring odds ratio between leukocyte count and obesity; it was found to be 7.2%. Among the various WBC types, neutrophilia is the rationale behind leukocytosis with the mean of 8.3 X 10⁹/l. (4.5-11-11.5x10⁹/l.). Furthermore, we deduced that there was no gender difference when relating obesity with leukocytosis using chi-square test, p-value=0.373.

Table 1: The characteristics of the participants studied.

	Non-obese group (n=127)	Obese group (n=154)
Male/Female	101/26	65/89
Age, Years	30.9 (range= 19-54 years)	33.1 (range= 18-60 years)
BMI (kg/m ²)	25.7(±2.5)	42.9 (±8.6)
Mean WBC count	6.4(±1.7)	8.4 (±2.3)

Table 2: The association between leukocyte count and BMI class.

	BMI class	Number of subjects	
		Normal	Leukocytosis
BMI	Normal (18.5-24.9)	43 (16.3%)	1 (5.6%)
	Overweight (25-29.9)	82 (31.2%)	1 (5.6%)
	obese class1 (30-34.9)	24(9.1%)	1(5.6%)
	obese class2 (35-39.9)	38(14.4%)	7 (38.9%)
	morbid obese >40	76(28.9%)	8 (44.4%)
	Total	263 (100%)	18(100%)

*Frequencies were found using chi-squared.




Figure 1: Correlation between WBC count and BMI.




Figure 2: Distribution of Leukocytosis among BMI classes.




Figure 3: One-way ANOVA test to compare the mean of each group.

Discussion

Our study aims to determine whether there is an association between obesity and leukocytosis. We found that the more obese an individual is more likely he or she is to have a high WBC count. An explanation for this is that obesity is known to be a chronic inflammatory condition with white blood cells infiltrating into adipose tissue and pro-inflammatory cytokines (tumor necrosis factor- α and interleukin-6) are produced by adipocytes and WBCs [12-14]. It has been postulated that one of the major determinants of the significant association of obesity and a high leukocyte count is the associated presence of metabolic syndrome [15]. However, our study shows that even in the absence of metabolic syndrome, obese individuals are found to have leukocytosis. We found that there is a correlation between BMI and total leukocyte count ($r = 0.360$, $P < 0.001$). The mean WBC count of the obese group is 8.5 ± 2.5 compared to 6.67 ± 1.8 for the non-obese group. Nonetheless, the absolute leukocyte number of obese individuals is still within the normal range ($4-11.5 \text{ mmol/L}$) [16]. This supports findings of previous studies which demonstrated a positive association between WBC count and BMI [8,9,17]. Furthermore, while looking at the differential leukocyte counts, Neutrophils were found to be elevated in morbidly obese patients mean $8.3 \times 10^9 \pm 3$. (Normal range: $2-7.5 \text{ K/mmol}$). This may be because chemokines are produced by adipose tissue which play a role in bone marrow granulopoiesis and also demarginate intravascular neutrophils [18-20].

The odds ratio for leukocytosis in obese patients with $\text{BMI} > 30$ is 7.2, which implies that obese individuals who have $\text{BMI} > 30$ are 7.2 times more likely to develop leukocytosis than those who are non-obese. Similarly, a cross-sectional study conducted in the Israel found that for every one kg/m^2 increase in BMI, the risk of having leukocytosis was increased by 11% [21]. The study also showed no significant difference between leukocytosis in obese females when compared to age-matched obese males, supporting our findings of no difference between genders. This supports the hypothesis that high body weight affects inflammatory markers, regardless of variation in body fat distribution peculiar to each gender. Children aged 6-12 years, has a significant neutrophilia and higher CRP concentration correlating with obesity. Thus it may be used as an obesity-related inflammatory marker in young

obese patients [22]. Given our retrospective study design, the limitations we encountered included omitting the assessment for the smoking status, CRP level and the waist circumference.

Moreover, within clinical settings leukocytosis should not be ignored, as an elevated leukocyte count even if within the normal range has been shown to be a risk factor for metabolic syndrome and cardiovascular disease [11,23-25]. A retrospective 5-year cohort study found that patients who developed Coronary Artery Disease initially had a high baseline leukocyte count when compared to normal patients [26]. However, the patho-physiology of leukocytosis and in turn, cardiovascular disease, caused by an increase in BMI is not clearly understood. It could be explained by the co-existence of metabolic syndrome [27]. We therefore recommend further prospective cohort studies to determine the clinical course of obese patients with leukocytosis.

Conclusion

The positive correlation between obesity and leukocytosis proves the relationship but does not establish causation. Although most obese patients in this study had a WBC count within the normal range, their counts were higher than the non-obese group due to prominent neutrophilia. Therefore, work-up of patients to rule out the common causes and subsequently treating the cause is essential. The association highlighted will have an impact on patients during the pre-operative work up for bariatric surgery, as neutrophilia is a risk factor for metabolic and cardiovascular complications that needs detection and treatment to minimize this risk. The incidental detection of leukocytosis in any obese patient is not yet rationalized and further research looking into the inflammatory and metabolic biomarkers are warranted to explain the etiology of this association.

References

1. Kelly T, Yang W, Chen CS, Reynolds K, He J (2008) Global burden of obesity in 2005 and projections to 2030. *Int J Obes (Lond)* 32(9): 1431-1437.
2. World Health Organization (WHO) (2016) Obesity and Overweight. World Health Organization, USA.
3. Klein S, Burke LE, Bray GA, Blair S, Allison DB, et al. (2004) Clinical implications of obesity with specific focus on cardiovascular disease: a statement for professionals from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism: endorsed by the American College of Cardiology Foundation. *Circulation* 110(18): 2952-2967.
4. Hotamisligil GS (2005) Role of endoplasmic reticulum stress and c-Jun NH_2 -terminal kinase pathways in inflammation and origin of obesity and diabetes. *Diabetes* 54(Suppl 2): S73-S78.
5. Hotamisligil GS, Shargill NS, Spiegelman BM (1993) Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance. *Science* 259(5091): 87-91.
6. Shoelson SE, Lee J, Goldfine AB (2006) Inflammation and insulin resistance. *J Clin Invest* 116(7): 1793-1801.
7. Cotran RS, Kumar V, Abbas AK, Collins T, Robbins SL, et al. (1999) Robbins Pathologic Basis of Disease. (6th edn), Walter Burns Saunders, Pennsylvania, USA, pp. 644-696.

8. Furuncuoğlu Y, Tulgar S, Dogan AN, Cakar S, Tulgar YK, et al. (2016) How obesity affects the neutrophil/lymphocyte and platelet/lymphocyte ratio, systemic immune-inflammatory index and platelet indices: a retrospective study. *Eur Rev Med Pharmacol Sci* 20(7): 1300-1306.
9. Chen SB, Lee YC, Ser KH, Chen JC, Chen SC, et al. (2009) Serum C-reactive protein and white blood cell count in morbidly obese surgical patients. *Obes Surg* 19(4): 461-466.
10. Ryder E, Ewald MD, Mosquera J, Fernández E, Pedrañez A, et al. (2014) Association of obesity with leukocyte count in obese individuals without metabolic syndrome. *Diabetes Metab Syndr* 8(4): 197-204.
11. Fadini GP, Marcuzzo G, Maescotti MC, de Kreutzenberg SV, Avogaro A (2012) Elevated white blood cell count is associated with prevalence and development of the metabolic syndrome and its components in the general population. *Acta Diabetol* 49(6): 445-451.
12. Das UN (2010) Obesity: genes, brain, gut, and environment. *Nutrition* 26(5): 459-473.
13. Sell H, Eckel J (2010) Adipose tissue inflammation: novel insight into the role of macrophages and lymphocytes. *Curr Opin Clin Nutr Metab Care* 13(4): 366-370.
14. Gregor MF, Hotamisligil GS (2011) Inflammatory mechanisms in obesity. *Annu Rev Immunol* 29: 415-445.
15. Yoshimura A, Ohnishi S, Orito C, Kawahara Y, Takasaki H, et al. (2015) Association of peripheral total and differential leukocyte counts with obesity-related complications in young adults. *Obes Facts* 8(1): 1-16.
16. Schwartz J, Weiss ST (1991) Host and environmental factors influencing the peripheral blood leukocyte count. *Am J Epidemiol* 134(12): 1402-1409.
17. Solá E, Jover A, López-Ruiz A, Jarabo M, Vayá A, et al. (2009) Parameters of inflammation in morbid obesity: lack of effect of moderate weight loss. *Obes Surg* 19(5): 571-576.
18. Suwa T, Hogg JC, English D, Van Eeden SF (2000) Interleukin-6 induces demargination of intravascular neutrophils and shortens their transit in marrow. *Am J Physiol Heart Circ Physiol* 279(6): H2954-H2960.
19. Rusten LS, Jacobsen FW, Lesslauer W, Loetscher H, Smeland EB, et al. (1994) Bifunctional effects of tumor necrosis factor alpha (TNF alpha) on the growth of mature and primitive human hematopoietic progenitor cells: involvement of p55 and p75 TNF receptors. *Blood* 83(11): 3152-3159.
20. Ulich TR, del Castillo J, Guo KZ (1989) In vivo hematologic effects of recombinant interleukin-6 on hematopoiesis and circulating numbers of RBCs and WBCs. *Blood* 73(1): 108-110.
21. Herishanu Y, Rogowski O, Polliack A, Marilus R (2006) Leukocytosis in obese individuals: possible link in patients with unexplained persistent neutrophilia. *Eur J Haematol* 76(6): 516-520.
22. Nascimento H, Rocha S, Rego C, Mansilha HF, Quintanilha A, et al. (2010) Leukocyte Count versus C-Reactive Protein Levels in Obese Portuguese Patients Aged 6-12 Years Old. *Open Biochem J* 4: 72-76.
23. Jung CH, Lee WY, Kim BY, Park SE, Rhee EJ, et al. (2013) The risk of metabolic syndrome according to the white blood cell count in apparently healthy Korean adults. *Yonsei Med J* 54(3): 615-620.
24. Madjid M, Awan I, Willerson JT, Casscells SW (2004) Leukocyte count and coronary heart disease: implications for risk assessment. *J Am Coll Cardiol* 44(10): 1945-1956.
25. Prentice RL, Szatrowski TP, Fujikura T, Kato H, Mason MW, et al. (1982) Leukocyte counts and coronary heart disease in a Japanese cohort. *Am J Epidemiol* 116(3): 496-509.
26. Takeda Y, Suzuki S, Fukutomi T, Kondo H, Sugiura M, et al. (2003) Elevated white blood cell count as a risk factor of coronary artery disease: inconsistency between forms of the disease. *Jpn Heart J* 44(2): 201-211.
27. Kip KE, Marroquin OC, Kelley DE, Johnson BD, Kelsey SF, et al. (2004) Clinical importance of obesity versus the metabolic syndrome in cardiovascular risk in women: a report from the Women's Ischemia Syndrome Evaluation (WISE) study. *Circulation* 109(6): 706-713.