

Isolation and characterization of psittacine beak and feather disease virus in Saudi Arabia using molecular technique

Abstract

Psittacine Beak and Feather Disease (PBFD) is a contagious, fatal viral disease that affects the beak, feathers, and immune system of wild and captive old and New World Psittacine. PBFD is characterized by feather abnormalities, beak and claw deformities and eventually dying as a result of immunosuppression. The disease is caused by member of the genus *circovirus* in the family *Circoviridae*. The disease diagnosed by clinical signs and laboratory detections. The serological diagnosis by haemagglutination (HA) and haemagglutination inhibition (HI) is subjective but molecular technique using Polymerase chain reaction (PCR) is the most reliable means for confirming the presence of the PBFVDV ambience DNA genome in whole blood or tissue samples. Many birds were presented to the veterinary clinic with clinical signs of PBFD. No reports document the presence of PBFVDV in KSA to date. Therefore, the objective of this work was to investigate the presence of PBFVDV in KSA using PCR. Total 175 samples (blood and feathers) from clinically-suspect, and apparently-health birds from 10 different psittacine species were collected, DNA was extracted and conventional PCR was performed. The viral DNA was identified in six samples (6/175) four Grey parrots (*Psittacus erithacus*), and two ring necks (*Psittacula eupatria eupatria*). PBFVDV was identified for the first time in the Kingdom of Saudi Arabia (KSA). During this investigation, we developed a technique for faster and simpler processing of multiple feather samples. The isolated PBFVDV could be characterize and use as positive control for further research purpose. We recommend that the Rep gene feather-based PCR technique be established as a routine diagnostic tool in quarantine facilities across the country.

Keywords: psittacine beak and feather disease, polymerase chain reaction, Saudi Arabia, grey parrot, *psittacus erithacus*

Volume 2 Issue 1 - 2017

A Hakami,¹ A Al-Ankari,² M Zaki,² A Yousif²

¹Sabya Municipality, King Faisal University, Saudi Arabia

²Avian Diseases, King Faisal University, Saudi Arabia

Correspondence: Alaudeen Hakami, Sabya Municipality, Main road, Sabya, Gizan, Saudi Arabia, Tel 00966557669995, Email hakamialaa@hotmail.com

Received: May 11, 2017 | **Published:** June 06, 2017

Abbreviations: PBFD, psittacine beak and feather disease; HI, haemagglutination inhibition; HA, haemagglutination; PCR, polymerase chain reaction; KSA, kingdom of Saudi Arabia; RT, room temperature

Introduction

Psittacine beak and feather disease (PBFD) was first described in Australian cockatoos in early 1970s. The disease was characterized by varying degrees of symmetric feather dystrophy and loss. PBFD outbreaks of wild populations are still common in Australia while outbreaks of captive birds occur worldwide.¹ Antibody detection tests, commonly used in identification of PBFVDV are HA and HI, but their results are subjective.² However, to overcome this problem, PCR have been developed for identification and classification of several pathogens including PBFVDV. These molecular techniques offer higher sensitivity and specificity compared to serological test.³ In addition, PCR based diagnosis can be used for identification of both clinically suspect, and apparently health carrier cases of PBFD.⁴ A previously described Rep gene-based PCR assay⁵ was used in this investigation. This technique utilizes a highly conserved region of the PBFVDV ORF 1 (the rep gene) that can reliably use for sequence comparisons and phylogenetic tree analysis. The clinical manifestations of PBFD have been observed in the KSA. However, no scientific reports document the presence of PBFVDV in KSA psittacine populations to date.⁶ Therefore, the objective of this work was to investigate the presence of PBFVDV in KSA psittacine populations and to determine if it is

feasible to monitor the presence of the virus routinely. The most serious problem facing this investigation was the absence of a proper positive control. The PCR experiment had to be optimized without the importation of possibly infectious material from other laboratories. This decision was made in order to eliminate possible risks associated with handling and importation of exotic avian disease agents into the country. The knowledge of the presence of PBFVDV in the KSA and availability of the technology for its diagnosis is extremely important for the development of proper disease control programs for the preservation of the psittacine bird populations in the country.

Materials and methods

Summary description of the experimental design

This investigation was designed to detect the presence of PBFVDV in psittacine populations in the KSA. The detection method selected was an ORF1 PCR assay that was proven highly sensitive in the detection of the virus from various clinical specimens. The bird samples targeted were feathers and blood (when available). These sample types were reported to be the best for PCR-based detection of the virus. The samples were collected from the different bird populations and species where access was permitted. It was recommended that a total of 250 birds be sampled for this investigation because of the reported high prevalence of the virus in wild and domestic psittacine populations worldwide; ultimately, only 175 samples were available for PCR analysis. DNA was extracted from the different clinical samples using commercial kits that ensured high yield and

purity. DNA was then tested for the presence of the PBFVDV ORF 1 sequences using the reference primers. Avian mitochondrial DNA from each extracted sample was targeted for PCR amplification in separate reactions to ensure that sufficient amounts of sample DNA were extracted, and introduced in the diagnostic PBFVDV PCR master mix (Quality control).

Sample collection and preservation

Samples from 19 clinically-suspected (Figure 1), and 156 apparently-health birds were collected. Samples included chest feathers and whole blood. Blood was collected either by puncturing the wing vein using sterile disposable needle, or by clipping one of the nails off the claw. Blood was collected directly into 1.5 ml sterile centrifuge tubes. Feather samples were plucked from the chest area or, collected from the bird's environment and were placed in sterile test tubes. Both sample types were labeled and stored in separate freezer boxes at -80°C in the Avian Diseases Laboratory, College of Veterinary Medicine and Animal Resources, King Faisal University, Al-Ahsa, KSA. The bird species sampled during this investigation belonged to the following genera: Sulphur-crested cockatoos (*Cacatua galerita*), Indian ringneck (*Psittacula eupatria eupatria*), Local ringneck parrot (*Psittacula eupatria*), Lovebird (*Agapornis roseicollis*), Cockatiel (*Nymphicus hollandicus*), Senegal parrot (*Poicephalus senegalus*), African grey (*Psittacus erithacus*), Fisher (*Agapornis fisheri*), Macaw (*Ara macaw*), and Amazon (*Amazona autumnalis*).



Figure 1 Clinically suspected birds showed symmetric feather dystrophy and loss.

Samples from different locations in Eastern and Southern provinces of KSA were collected (Figure 2).

A = Eastern provinces, B=Southern provinces. Bird samples were collected over a period of three years starting from 2008 and ending early 2010. The reason for this was to identify a possible introduction time for the virus. Only 8 samples were collected at the end of 2008, 48 samples during the first months of 2010 and the rest of the 175 samples during 2009.



Figure 2 Samples from different locations in Eastern and Southern provinces of KSA were collected.

DNA extraction from blood

DNA was extracted from blood samples using DNeasy Blood and Tissue kit (Qiagen, Cat. 69504, USA). Extraction was performed according to the manufacturer's instructions. Briefly, blood tubes were allowed to thaw at room temperature (RT). Twenty microliters of proteinase K were pipetted into the bottom of the microcentrifuge tubes containing the blood sample. The volume was then adjusted to 220µl with PBS. Two hundred microliters of buffer AL were further added and mixed by pulse-vortexing for 15seconds, or until clotted blood was completely submerged in the lysis buffer. Samples were incubated for 10min at 56°C to allow for digestions, and then spun down briefly to remove drops from the inside of the lid. Two hundred microliters of 100% ethanol were added, mixed for 15 seconds pulse-vortexing, and briefly centrifuged again. The mixture was then transferred to a DNeasy mini spin column inserted in a 2ml collection tube, and centrifuged at 8,000rpm for 1min, the flow-through and collection tube were discarded and the spin column transferred into a new collection tube. Five hundred microliters of buffer AW1 were added to the tube, and centrifuged as before. DNeasy mini spin column was then transferred to a clean collection tube, 500µl buffer AW2 were added, and centrifuged at 14,000rpm for 3min. DNeasy mini spin column was transferred to a new 1.5ml microcentrifuge tube, 200µl buffer AE were added and the column was incubated at RT for 1min to allow for DNA elution. Eluted DNA was collected by centrifugation at 14,000rpm for 1min. DNA extracts were preserved at -80°C until testing.

DNA extraction from feather

Viral DNA was extracted from feather samples using JETQUICK Tissue DNA Spin Kit, (Cat. No. 450 050, GENOMED GmbH, Germany). Briefly, feathers were allowed to thaw at RT before processing. Distal, differentiated portions were cut off and discarded. The growing tips were retained for DNA extraction. Tips were cut into small pieces using a sharp scalpel on a fresh piece of disposable aluminum foil for each. Cut tips were then crushed using the broad end of the scalpel blade. Crushed tips were transferred into fresh microfuge tube, and lysis buffer was added according to the manufacturer's instructions. For confirmation of direct extraction from crushed tips, duplicates of some of the samples were homogenized in a sterile mortar and pestle using sterile sand and PBS prior to addition to lysis buffer. Twenty microliters of proteinase K were pipetted into the bottom of the 1.5ml microcentrifuge tube containing the crushed, or homogenized, tips. Two hundred microliters of buffer T1 were added, and mixed thoroughly by inverting the reaction tube several times. Tubes were then incubated for 1-2h at 56°C to allow for digestion. Tubes were then spun down briefly to remove drops from the inside of the lid. Two hundred microliters of Buffer T2 were added, mixed thoroughly by vortex, and incubated for 10min at 70°C. The mixture was allowed to cool down for approximately 1min. Two hundred microliters of absolute ethanol were added to each extraction tube. The mixture was then transferred to the reservoir of the micro-spin column and centrifuged at 10600rpm for 1min. The flow-through was discarded and 500µl of reconstitution buffer TX were pipette into the reservoir. The spin columns were centrifuged for 1min at 10600rpm. The flow-through was discarded and five hundred microliters of buffer T3 were added to the reservoirs. Tubes were centrifuged as before. The flow-through was discarded, columns were replaced in the collection tubes, and centrifuged for 1min at 13,000rpm to remove residual buffer T3. The micro-spin columns were transferred to new

1.5ml microcentrifuge tubes, 200µl of 10mM Tris-HCL buffer [pH 9.0] pre-warmed to 70°C were added, incubated at room temperature for 5min and finally centrifuged subsequently for 2min at 13,000rpm.

Determination of extracted dna quality and quantity

DNA concentration and purity was determined spectrophotometrically using SmartSpec™ Plus spectrophotometer (BioRad, Hercules, CA, USA). Purity was determined by 260/280nm ratios using distilled water for blanking. DNA integrity was determined by electrophoresis in 1% Ethidium bromide-stained agarose gels using TAE at 5V/cm for 20min at RT. DNA was visualized in a UV transilluminator (Gel Doc XR gel documentation system, BIO-RAD laboratories, Milan, Italy). PCR A 603-nt fragment (Nucleotides 178-780) of the PBFVDV genome was amplified using PCR with the primers 5'-TTAACAACCCTACAGACGGCGA-3' and 5'-GGCGGAGCATCTCGCAATAAG-3'.⁵ Primers amplify a region of the viral genome within the coding region of the Rep protein. A 530-nt fragment of the avian mitochondrial DNA 12S rRNA gene was also amplified using the primers L-12SA 5'-AAACTGGGATTAGATACCCCACTAT-3' and H-12SB2 5'-TTCCGGTACACTTACCTTGTTACGAC-3'.⁷ The mitochondrial DNA was amplified as a control for the presence of avian DNA in extracts and the absence of reaction inhibitors. Amplification reactions were carried out using the HotStar Taq® Plus Master Mix PCR kit (QIAGEN, Hilden, Germany). Each reaction tube contained 1 unit of HotStar Taq® Plus DNA polymerase in 1×PCR buffer containing (KCl and (NH₄)₂SO₄), 200uM of each dNTP, 0.5µM of each of the forward and reverse primers, 1×CoralLoad dye mix, and 3µl of the sample DNA and controls. The reaction tubes were centrifuged at 2000×g briefly, and placed in a BioRad MyCycler™ thermal cycler (BioRad, Hercules, CA, USA). Thermal cycling conditions for amplification of viral and avian DNA targets genes were: an initial denaturation step (95°C for 4 min), 38 amplification cycles (95°C for 30sec, 54.5°C for 30sec, and 72°C for 50sec), and a final extension step (72°C for 5min). After amplification, 15µl of each sample were analyzed by electrophoresis on a 1.2% agarose gel stained with Ethidium bromide (0.5µg/ml). DNA bands were visualized by UV irradiation in a Gel Doc XR gel documentation system (BIO-RAD laboratories, Milan, Italy).

Results

Recovery of DNA from avian samples

Blood and feather samples subjected for DNA extraction yielded concentrations from 1-11 Kochev µg/ml (0.2-2.2µg/sample). The DNA extracted was highly pure (OD₂₆₀/OD₂₈₀ values ranging from 1.7-2.5).

Amplification of mitochondrial DNA from avian DNA extracts

Avian mitochondrial DNA amplification was used prior to PBFVDV PCR assay to ensure that DNA concentration and quality were not factors in PBFVDV PCR results. Successful amplification of a 530bp PCR product was achieved from tested avian DNA extracts at the lower and higher DNA concentration values (Figure 3). There were no indications of PCR inhibition in DNA extracts from either blood or feathers. Ten microliters of avian blood were extracted using the DNeasy Blood & Tissue Kit (QIAGEN Sciences, Maryland, and USA). PCR was conducted using the 12S rRNA gene primers.⁸ Lane M, standard size marker, L₁ negative control, L_{2&3} test samples.




Figure 3 Amplification of mitochondrial DNA from Avian DNA extracts.

NDetection of PBFVDV DNA and development of PCR positive control

DNA from 10µl of avian blood collected from a suspected bird was extracted using the DNeasy Blood & Tissue Kit (QIAGEN Sciences, Maryland, and USA). PCR was conducted using the PBFVDV Rep gene PCR primers as described above. A 603bp PCR product was amplified from the avian sample. This was the first report of PBFVDV in Saudi Arabia (Figure 4, lane 2). The PCR product obtained from that first PBFVDV positive sample was used as a positive control after sequence confirmation (as described below). PBFVDV PCR-products were used as templates after 1:1000 dilutions. PCR reactions produced a clear positive PCR band of the correct size (603bp). Ten microliters of avian blood collected from a suspect bird (showing clinical presentation of PBFVDV) were extracted using the DNeasy Blood & Tissue Kit (QIAGEN Sciences, Maryland, USA). PCR was conducted using the replicase-associated PBFVDV gene PCR primers. Lane 1 Negative control, Lane 2 DNA extract from PBFVDV positive sample, lane 3 DNA extract from PBFVDV sero-negative sample, Lane 4 PBFVDV PCR positive product. Lane M standard size marker.




Figure 4 PBFVDV Positive controls.

PCR screening of avian extracts for PBFVDV DNA

A total of 6 psittacine bird samples were positive using PBFVDV PCR (Figure 5) (Table 1). Two positive samples were collected during 2008, 3 during 2009, and only 1 during 2010. All positive samples were collected from clinically suspected birds. Four of the positive samples were collected from Grey parrots (*Psittacus erithacus*) and 2 from ring neck birds (*Psittacula eupatria*). It was interesting to notice that the two birds from which both blood and feathers were collected, only feathers were positive to PBFVDV. Only two blood samples were positive to PBFVDV. All positive samples were collected from the Eastern province. PBFVDV PCR screening samples Blood and feather samples were tested using the replicase-associated PBFVDV gene PCR primers. Lane 1-20 (upper) screened blood samples, Lane 1-20 (lower) screened feather samples. Lane 4 (upper) and lane 1 (lower) PBFVDV control positive. Lane 1 (upper) and lane 5 (lower) control negative, Lane 9 (upper) and Lane 8 (lower) standard size markers. Lanes 1-3, 5-8, 10-20 (upper), Lane 2-7, 9-20 (lower) test samples.




Figure 5 PBFDV PCR screening samples.

Table I Summary of PBFDV PCR screening from psittacine birds collected in the KSA

Host species	Host common name	Sampling location	Date	Clinical status	Blood	Feather
<i>Psittacus erithacus</i>	Grey parrot	Avian clinic	2008	S*	+ve	**
<i>Psittacus erithacus</i>	Grey parrot	Avian clinic	2008	S	+ve	**
<i>Psittacus erithacus</i>	Grey parrot	Alqatif bird marketing	2009	S	-ve	+ve
<i>Psittacus erithacus</i>	Grey parrot	Al-ahsa bird market	2009	S	**	+ve
<i>Psittacula eupatria</i>	Local ringneck	Alqatif private farm	2009	S	**	+ve
<i>Psittacula eupatria</i>	Indian ringneck	Al-ahsa bird market	2010	S	-ve	+ve

Discussion

PBFD is the most common viral disease of wild and captive psittacine birds. It has the potential to become a major threat 42 species of wild Parrots throughout the continents.⁹ To date, PBFVDV was not reported in the KSA. Five years ago, our clinical observations claimed that PBFVDV could present in the country. In order to prove or rule out this assumption, 175 psittacine blood and feather samples were collected during the years 2008-2010 and processed.¹⁰ A survey in South Africa investigated only 161 bird samples and yielded 75 positive samples.¹¹ Therefore, considering the expected total number of psittacine birds available, 175 samples could be deemed to achieve the primary goal of detection of the virus in the Eastern and Southern provinces of Saudi Arabia. We developed a simple feather-tip processing technique to reduce the need for using multiple sterile mortars. The technique relied on cutting, then crushing feather tips on sterile pieces of aluminum foil. This reduced the cost and time required for processing feather samples drastically. The technique used for processing blood and feathers was validated by assessing the DNA concentration and quality spectrophotometrically, and with the use of mitochondrial DNA PCR analysis. Our primary focus was on suspect samples showing the typical clinical presentation of PBFD. These samples were taken mainly from the avian clinic, the Veterinary Teaching Hospital, College of Veterinary Medicine and Animal Resources. It was decided that standard positive samples will not be allowed into our laboratory to avoid the possibility of contamination of the tests, and/or the environment with a novel virus. One blood sample collected in late 2008 produced the expected size PCR product in gel analysis. We intended to sequence the PBFD PCR positive products, and subsequently, use as control positive for field investigations. Positive samples identified were at the lower limit of the test detection capacity (Figure 4). Serial dilution of positive samples did not result in PCR amplification. This forced us to develop a PCR positive control for the PBFVDV PCR assay using diluted PCR products. Up to 103 dilution of the positive control gave a strong PCR product in Rep

gene PCR assays (Figure 5). The optimized control positive DNA will be used for research purposes. Only six samples tested positive for PBFVDV (3.42%). All positive samples were suspected cases (31.6% of suspect birds). The majority of suspected cases (68.42%) were PBFVDV-negative. Surprisingly, PBFVDV was not detected in any of the apparently normal birds. Latently infected birds can cause a severe environmental contamination.^{7,12} Latent infections have been reported to be the majority of infections in captive psittacine populations.¹³ The small number of positive samples detected in this relatively large sample size and the absence of detectable latent infections may be attributed to:

- I. Intensive breeding of psittacines in Saudi Arabia is a relatively a new practice; which would reduce the chance of horizontal and vertical transmission of the virus reported by others.¹³
- II. Psittacines imported into Saudi Arabia are from the new world; which suffers a lower prevalence of the virus.¹⁴

Another interesting finding is that positive samples were only found in 2 out of 13 bird species (Table 1) investigated. The ring-neck is bred easily in Saudi Arabia. The African grey parrot is expensive and owners are willing to spend money to try to treat infected birds. Therefore, it is possible that the virus was detected in both but not in other species where breeding is less likely, or the cost is low enough to discourage taking care of sick birds (which would lead to discarding the bird when clinical signs appear). Moreover, two of the positive samples were only positive upon testing feathers and not blood. This pointed to the importance of feather sample testing for screening of birds for PBFVDV.

Conclusion

It is important not to consider the 4% to be a real indication of the prevalence of the disease in KSA. It should be taken only as an indicator for the need to further investigate the prevalence of the disease in the different avian farms and at ports. In support of this conclusion is the

fact that the six positive samples were from suspected cases. PCR is the only tool to be recommended to detect cases with Pbfd, as some other diseases and manifestation show similar clinical signs.¹⁵ This technique is an excellent tool to be used for easy an epidemiology studies for the Pbfd in Saudi Arabia. This study recommends all imported birds are subject to be tested for Pbfd by using PCR in order to prevent entry of the virus.

Acknowledgements

This project was funded by the Deanship of Scientific Research, King Faisal University. The technical assistance of Dr. Mohammad Al-kahtani, King Faisal Special Hospital and Research Centers, Riyadh, was appreciated.

Conflict of interest

We declare that there is no financial interest or any conflict of interest exists.

References

1. Pass DA, Perry RA. Psittacine Beak and Feather Disease: An update. *Aust Vet J.* 1984;61(3):69–74.
2. Johne R, Raue R, Grund C, et al. Recombinant expression of a truncated capsid protein of beak and feather disease virus and its application in serological test. *Avian Pathol.* 2004;33(3):328–336.
3. Bonne N, Shearer P, Sharp M, et al. Assessment of recombinant beak and feather disease virus (BFDV) capsid protein as a vaccine for psittacine beak and feather disease (Pbfd). *J Gen Virol.* 2009;90(3):640–647.
4. Hess M, Scope A, Heinz U. Comparative sensitivity of polymerase chain reaction diagnosis of psittacine beak and feather disease on feather samples, cloacal swabs and blood from budgerigars (*Melopsittacus undulatus*, Shaw 18005). *Avian Pathol.* 2004;33(5):477–481.
5. Ritchie PA, Anderson IL, Lambert DM. Evidence for specificity of psittacine beak and feather disease viruses among avian hosts. *Virology.* 2003;306(1):109–115.
6. Hakami A, Al-Ankary A, Zaki M, et al. *Beak and Feather Disease Virus in the Kingdom of Saudi Arabia.* 2010.
7. Ritchie B. Function and Control. In: *Avian Viruses.* USA: Wingers Publishing; 1995. p. 223–252.
8. Ritchie B, Niagro F, Latimer K, et al. A review of psittacine beak and feather disease. *J Assoc Avian Vet.* 1989;3:143–149.
9. Saif Y, Barnes H, Glisson G, et al. *Disease of poultry.* 11th ed. USA: Iowa State Press; 2003.
10. Shearer PL, Bonne N, Clark P, et al. Beak and feather disease virus infection in cockatiels (*Nymphicus hollandicus*). *Avian Pathol.* 2008;37:75–81.
11. Kondiah K, Albertyn J, Bragg R. Genetic diversity of the Rep gene of beak and feather disease virus in South Africa. *Arch Virol.* 2006;151:2539–2545.
12. Dahlhausen S, Radabaugh M. Update on psittacine beak and feather disease and avian polyomavirus – epidemiology and diagnostics. *Aust Vet Proced.* 1997:51–57.
13. Rahaus, M, Desloges N, Probst S, et al. Detection of beak and feather disease virus DNA in embryonated eggs of psittacine birds. *Veterinari Medicina.* 2008;53(1):53–58.
14. Bert E, Tomassone L, Peccati C, et al. Detection of beak and feather disease virus (BFDV) and avian polyomavirus (APV) DNA in psittacine birds in Italy. *Vet Med B Infect Dis Vet Public Health.* 2005;52:64–68.
15. Schat K. Circovirus infections. *Diseases of Poultry.* In: Saif Y, Fadly A, Glisson J, McDougald L, Nolan K, Swayne D, editors. 11th ed. USA: Iowa State Press; 2003. p. 181–201.