

Toxic trace elements in selected edible rhizomes of medicinal plants using INAA and ICP-MS techniques

Abstract

Consumption the crop is one of the main sources of dietary exposure to toxic trace elements. In order to assess the level of toxic trace elements from selected herbs and to make an assumption on their consumption level of concern, fourteen elements in *Zingiber zerumbet* (lempoyang), *Boesenbergia rotunda* (temu kunci), *Zingiber officinale* var. rubrum (halia bara) and *Zingiber officinale roscoe* (halia) have been determined via ICP-MS (Cd, Be, Ti, Pb) and INAA (As, Al, Ba, Cr, Co, Sb, Sr, Th, U and V) methods. It was revealed that the concentrations of these elements in most selected rhizomes did not exceed the standard dangerous toxic level. However, 47% cadmium accumulation was detected in *Zingiber officinale* var. rubrum grown in poly-bags.

Keywords: toxic element, herbs, rhizomes, inaa, icp-ms

Volume 6 Issue 4 - 2017

AbuBakar Mohd Daran,¹ Kamaludin Rashid,² Halijah Ibrahim,³ Mahanom Jalil,² Yusmin Mohd Yusof,² Shamrulazhar Shamzir Kamal,³ Shahril Efzueni,² Reza Farzinebrahimi²

¹Department of Physics, University of Malaya, 50603 Kuala Lumpur, Malaysia

²Department of Biology, University of Malaya, Malaysia

³Institute of Biological Sciences, University of Malaya, Malaysia

Correspondence: Reza Farzinebrahimi, Institute of Biological Sciences, Faculty of Science Building, University of Malaya, 50603 Kuala Lumpur, Malaysia, Tel 0060173030586, Email rfebrahimi@siswa.um.edu.my

Received: October 29, 2016 | **Published:** April 21, 2017

Abbreviations: Cd, cadmium; Be, beryllium; Ti, thallium; Pb, lead; As, arsenic; Al, aluminum; Ba, barium; Cr, chromium; Co, cobalt; Sb, antimony; Sr, strontium; Th, thorium; U, uranium; V, vanadium; MRL, minimal risk level; INAA, instrumental neutron activation analysis; ICP-MS, inductively coupled plasma/mass spectrometry

Introduction

The interaction between chemical extracted compounds from plants and the human body go through the same identical process to those well understood for the chemical compounds in conventional drugs. Similarly, the herbal medicine may cause the same potential to cause a side effect.¹ *Zingiber zerumbet* (lempoyang), *Boesenbergia rotunda* (temu kunci), *Zingiber officinale* var. rubrum (halia bara) and *Zingiber officinale roscoe* are belonging to Zingiberaceae or Ginger family consisting of aromatic perennial herbs with creeping horizontal or tuberous rhizomes planted locally. The same species can be found throughout tropical Africa, Asia, and the America. These rhizomes are a common edible ingredient in many countries, in South East Asia and serve as culinary herbs due to their aromatic flavor to promote appetite.

Their popularities as folk medicine have drawn further interest amongst researchers to investigate on various perspectives about their medicinal properties. Eng-Chong et al.,² Sivasothy et al.,³ & Sontakke et al.,⁴ have been reported on medicinal crop areas. A list of possible treatments for illnesses using the described rhizomes is shown in Table 1.

Trace elements may take by medicinal plants from the surrounding mineral or contaminated soil environment via the root system and translocation within at various part of the plant such as in the rhizome, stems, and leaves. Some plants can absorb trace elements from the soil in high quantity and do not present toxicity symptoms or have their growth affected. These plants are suitable for remediation of polluted soils known as Phytoremediation.⁵ Accumulation of toxic trace element above minimal risk level (MRL) value or provisional tolerable index level, ingested by humans for a long term, will give

an adverse effect to human health depending on the type of trace elements present.

The present study was undertaken to determine the level of toxic trace elements found in the rhizome samples and to make an assumption on their consumption level of concern if ingested by an adult based on the MRL tabulated by the Agency for Toxic Substances and Disease Registry.⁶

Materials and method

The rhizomes of four species, namely *Z. zerumbet* (A), *B. rotunda* (B, C), *Z. officinale* var. rubrum (D, E) and *Z. officinale Roscoe* (F) were collected from the surrounding area in the University of Malaya. The *B. rotunda* was separated into a center piece or the top part and the tubular shape and *Z. officinale* var. rubrum were taken from the wild and planted in polybag condition.

The samples were pre-rinsed with deionized water and dried at 39°C in the oven for five hours. The dried samples were powdered using mortar and pestle to make a dried rhizome powder. The powders were sieved using a standard set of sieves. About 15- 100g of grounded sample was weighed and sealed in the polyethylene vials and tagged as A, B, C, D, E and F according to the species and their origin. A duplicate was used up for each analyte sample and blank for error correction.

Standard coal ash SRM-1632a based on (NIST)⁷ was employed as a multi element comparison standard. The Analyte and standard samples were stacked up in a rotary rack at different positions ready for irradiation.

Irradiation and counting

The irradiation facility was carried out by TRIGA MK II (Malaysian Institute of Nuclear Technology, MINT) for 6hours at a thermal flux (2×10^{12} ncm⁻²Sec⁻¹). After irradiation, the vial is returned from the reactor by atmospheric pressure and allowed to decay to the level of activity is within acceptable limits for handling. Cooling time was arranged according to the nuclear properties of

radioisotopes. Medium-lived species were counted after four days of cooling while a decay time of nearly a month was set aside for a long-lived isotope. Each sample was placed in the appropriate position on the detector for counting.

For the gamma spectra measurement, a solid state horizontal hyper pure germanium (HPGe) detector was used. A nuclear Data 66 multichannel analyzer was used for pulse height analysis. The gamma spectra analyses were carried out by using the software Genie 2000 and gamma vision supplied by ORTEC and Canberra, respectively.

Heavy metals and non-measurable by Instrumental Neutron Activation Analysis (INAA), such as Pb, Be, Tl, and Cd were analyzed using a Perkin-Elmer SCIEX Elan 6000 Inductively Coupled Plasma/Mass Spectrometry (ICP-MS).

The dry duplicate sample was weighed (0.5g) and transferred into a beaker (10mL) followed by concentrated HNO₃ (5mL). The sample was pre-digested for one hour, and the beaker was heated up on a hot plate at 60-70°C.

Table 1 Medicinal plants and its uses

Species	Parts used	Treatment of
Boesenbergia rotunda	Swollen roots and leaves	Rheumatism, pain killer, gout, gastrointestinal disorders, stomach ache, dyspepsia and peptic ulcer ²
	Rhizome	Anti cancer ¹² wound healing ¹²
Zingiber zerumbet	Rhizome	Inflammation and pain-mediated diseases, worm infestation and diarrhea ³ Anti cancer ¹³
Zingiber officinale var. rubrum	Rhizome	Anti cancer, nausea, antibacterial activity ¹⁴
Zingiber officinale Roscoe	Rhizome	Nausea (seasickness), morning sickness and ^{4,15}

Table 2 Result for samples analyzed using ICP-MS and INAA techniques

	Elements	Concentration (ppm)					
		A	B	C	D	E	F
ICP-MS method	Cd	0.02±0.01	0.04 ±0.02	0.04±0.01	0.44±0.01	0.10±0.01	0.22±0.01
	Be	<0.01	<0.01	<0.01	<0.01	0.02±0.01	<0.01
	Tl	<0.01	0.02±0.01	<0.01	0.07±0.01	0.02±0.01	<0.01
	Pb	1.04±0.10	1.45±0.07	0.66±0.07	0.62±0.20	2.29±0.11	1.07±0.12
	As	0.09±0.01	0.07±0.01	0.13±0.01	0.40±0.02	<0.05	0.05±0.01
	Al	254±40	242±35	213±54	<1.0	<1.0	<1.0
INAA method	Ba	<1.0	23.7±0.7	8.74±2.01	14.7±1.6	10.1±1.9	4.05±0.81
	Cr	1.52±0.94	1.55±0.21	3.44±0.28	3.84±0.43	0.54±0.09	0.38±0.08
	Co	0.61±0.05	0.23±0.01	0.20±0.01	0.27±0.01	0.36±0.01	<0.1
	Sb	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
	Sr	<1.0	<1.0	<1.0	<1.0	10.1±1.0	9.3±1.0
	Th	0.26±0.01	0.99±0.19	1.21±0.07	0.55±0.02	0.62±0.08	1.03±0.03
	U	<0.05	0.09±0.02	0.11±0.01	0.05±0.02	<0.05	<0.05
	V	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1

Table 3 Estimation of tolerable dietary intake of rhizomes

Elements	ATSDR (MRL) ^a	@ 60 Kg BW	WHO (1999) ^b	Consumption Level of Concern
	ppm BW/day	mg/day	ppm	g/day (sample)
Cd	0.0005 (Int.)	0.03	0.3	68 (D)
Be	0.002 (Chr.)	0.12		6000 (E)
Tl	0.002	0.12		1714 (D)
Pb	0.007c	0.42	10	183 (E)
As	0.005 (Acute)	0.3	1	750 (C)
Al	1 (Int.)	60		236 (A)
Ba	0.2 (Int.)	12		506 (B)
Cr	0.005 (Int.)	0.3		78 (D)
Co	0.01 Int.)	0.6		983 (A)
Sb	0.0004 (0.024		480 (I)
Sr	2 (Int.)	120		1200 (E)
Th	15 pCi/Ld	0.13		109 (C)
U	0.002 (Acute)	0.12		1090 (C)
V	0.01 (Int.)	0.6		6000 (I)

A 2mL of HNO₃ and 3mL of HClO₄ were dropped slowly until the solution becomes clear and dissolved. The solution was filtered out using filter paper then was added up with deionized water (30mL). The blank was prepared using the same reagents undergoing the similar digestion procedure. If necessary, the sample and blank were diluted accordingly prior analysis. Moreover, enough samples based

on IAEA-SOIL-7 were digested with some digestion groups to assure the accuracy of the analytical procedures.

The elemental compositions for the four selected rhizome species, determined by ICP-MS and INAA are shown in Table 2. The samples are labeled as A, B, C, D, E and F with all concentrations are expressed

in parts per million (ppm). A total of 14 elements was detected. The Cd, Be, Tl and Pb by ICP-MS, and As, Al, Ba, Cr, Co, Sb, Sr, Th, U and V were revealed by INAA technique.

A, *Z. zerumbet* (lempoyang); B, *B. rotunda* (temu kunci)- centre piece; C, *Boesenbergia rotunda* (temu kunci)- tubular shape; D, *Z. officinale* var *rubrum* (halia bara) - polybag environment; E, *Z. officinale* var *rubrum* (halia bara)- wild environment; F, *Z. officinale* Roscoe (halia)

Result and discussion

The high concentration of aluminum (>200ppm) was observed in *Z. zerumbet* and *B. rotunda* in center piece and tubular shape. However, the trace amount was recorded in *Z. officinale* var *rubrum* (polybag and wild environment) and Al ions in *Z. officinale* Roscoe translucent very slowly to the upper parts of plants.⁸ The distribution of toxic trace elements in the studied rhizomes showed that the Al and Ba are found to be present at the major level, Cr, Pb, and Tl at the minor level and Cd, Co, Sb, Sr, Th, U and V at the trace level. Most plants contain no more than 0.2mg Al g⁻¹ dry mass, and this caused that the rhizomes of sample A, B and C is not an Al accumulator.

B. rotunda sp. (samples B and C) was accumulated with a higher mean of Ba concentration compared to *Z. officinale* var *rubrum* sp. (samples D and E). A little amount was detected from *Z. zerumbet* (sample A). Sample D was recorded rich of Cd, Tl, As, Ba and Cr compare to other specimens. Even compared to the sibling sample E, the ratio of these toxic minerals can go from 1.5-8 folds higher. In a contrary, sample E absorbed more Pb and Sr. Since sample D and E grew in different cultures thus the sources of toxic elements up taken by the plants may come from the soil compost, the fertilizer or the polybag. Heavy metals are often used in pigments or as stabilizers in polymer materials. Thus, polymers inevitably contain hazardous metals or toxic elements such as As, Cd, Cr and Zn to a certain degree.⁹

Similarly, sample B showed higher concentrations of Pb and Ba over sample C that in contrary marks higher concentration in Cr. It led to Pb and Ba translocated at the centerpiece of the rhizome while Cr resides more in the tuberous shape. Other elements observed a balanced distribution throughout the rhizome of *B. rotunda* sp. The Sr was only detected in sample E and F. Concentration of Be, Sb and V were below the detection level (undetected) for all samples. Comparatively, sample A, B and C contain a high amount of toxic trace mineral observed, whereas sample D, E and F are due to the negligible contribution from Al.

The estimation of tolerable dietary intake of the selected rhizomes was calculated based on the published data from.^{6,10} According to the guideline for the medicinal plant,¹⁰ the concentrations of As, Cd and Pb do not exceed the limits of 1, 0.3 and 10ppm respectively. Based on Table 3, the concentration of As, Cd and Pb were measured below the recommended limit except in sample D where it registered 47% higher for Cd. For Duration, Acute = 1 to 14days, Intermediate = 15 to 364days, and Chronic = 1year or longer. a6, b10, c16, d17.

The minimal risk level (MRL) tabulated by ATSDR was used to estimate the number of rhizomes may be consumed per day. For example, the MRL of Cd taken orally continuously for 15days to a year without giving adverse health effect is 0.0005ppm BW/day. Assuming an adult of 60kg BW may require 0.03mg of the Cd/day for biosystem sustainability function. The consumption level of concern for Cd was calculated from the ratio of adult MRL to the maximum

value in Cd column detected from the samples. As shown in Table 2, sample D was recorded the highest concentration of Cd, and the tolerable dietary intake of rhizome D was only 68g. Similarly, the MRL for Cr was attained when someone ingested a dry weight of 78g of the same rhizome.

An herbal infusion of these rhizomes may dilute the concentration of toxic trace element to some extent due to low extraction efficiency and in some cases the inorganic toxic trace element such as were not detected in water infusion.¹¹

Conclusion

The obtained amount of toxic trace elements in medicinal rhizomes is being less than the permissible levels allowed by the MRL standard of ATSDR. Cadmium, Tellurium, Arsenic, and Chromium showed high concentration in *Z. officinale* var *rubrum* (growth in polybag environment). However, the level of Cadmium detected higher than the recommended level. In *Z. officinale* var *rubrum* (growth in the wild environment) the same pattern did not copy. It could be a coincidence the sample that grown in Agra-polybag may contain specific toxic trace element used for strengthening and stabilizing poly bags.

Ingestion of these rhizomes as a medicinal herb is considered safe as long as the period of ingestion is kept short and with the quantified amount. Water infusion further reduced the risk of metal toxic poisoning possibly due to low extraction efficiency.

Acknowledgments

The authors gratefully acknowledge the financial support provided by the University of Malaya. The authors would like to express their deepest gratitude to Dr. Mohd Shuhaimi Hamzah and Mr. Md Shuhaimi Elias from Malaysian Institute of Nuclear Technology (MINT) for rendering the INAA and ICP-MS services and also for running the samples.

Conflicts of interest

Author declares there are no conflicts of interest.

Funding

None.

References

1. Tapsell LC, Hemphill I, Cobiac L, et al. Health benefits of herbs and spices: the past, the present, the future. *Med J Australia*. 2016;185(4 Suppl):S4-S24.
2. Eng-Chong, T, Yean-Kee L, Chin-Fei C, et al. *Boesenbergia rotunda*: From Ethnomedicine to Drug Discovery. *Evid Based Complement Alternat Med*. 2012;25.
3. Sivasothy Y, Chong WK, Hamid A, et al. Essential oils of *Zingiber officinale* var. *rubrum* Theilade and their antibacterial activities. *Food Chem*. 2011;124(2011):514-517.
4. Sontakke S, Thawani V, Naik MS. Ginger as an antiemetic in nausea and vomiting induced by chemotherapy: a randomized, cross-over, double blind study. *Indian J Pharmacol*. 2003;35:2-36.
5. Khalili H, Soltani R, Negahban S, et al. Reliability of Disk Diffusion Test Results for the Antimicrobial Susceptibility Testing of Nosocomial Gram-positive Microorganisms: *Is E-test Method Better?* *Iran J Pharm Res*. 2012;11(2):559-563.
6. Ismail M, Bagalkotkar G, Iqbal S, et al. Anticancer Properties and Phenolic Contents of Sequentially Prepared Extracts from Different Parts of Selected Medicinal Plants Indigenous to Malaysia. *Molecules*. 2012;17(5):5745-5756.

7. NIST. Trace Elements in Coal, (Bituminous). USA: National Institute of Standards and Technology (NIST); 1978.
8. Hombach M, Zbinden R, Böttger EC. Standardisation of disk diffusion results for antibiotic susceptibility testing using the sirscan automated zone reader. *BMC Microbiology*. 2013;13:225.
9. Manoharan A, Pai R, Shankar V, et al. Comparison of disc diffusion & E test methods with agar dilution for antimicrobial susceptibility testing of *Haemophilus influenzae*. *Indian J Med Res*. 2003;117:81–87.
10. Bagalkotkar G. Isolation and Characterisation of Compounds from ‘Naga Buana’ (*Phyllanthus Pulcher*) and ‘Simit Matinggi’ (*Casaria Capitellata*) and their Cytotoxic Effects on Cancer Cell Lines, Master Thesis, Faculty of Medicine and Health Science. Malaysia: *Universiti Putra Malaysia, serdang*; 2007. pp 240.
11. Lalitha MK. Manual on antimicrobial susceptibility testing. India: Department of Microbiology, Christian Medical College; 2005.