

# Water quality assessment and apportionment of northern Pakistan using multivariate statistical techniques – a case study

## Abstract

The present study examines the physico-chemical characteristics in Gilgit and Hunza rivers, northern Pakistan. Twenty-nine water samples from different locations of Gilgit and Hunza rivers were collected during July 2012. The analytical results of the samples point out high concentration of bicarbonates. Large variations in the level of various parameters including chloride, total alkalinity, total hardness, sulphates and nitrates were observed. Multivariate statistical techniques were employed to interpret the data and results of physico-chemical properties showed that values of all parameters were in accordance with the permissible limits proposed by World Health Organization guidelines. The high values of total alkalinity showed that surface water of the study area is of bicarbonate type.

**Keywords:** physico-chemical characteristics, gilgit and hunza river, multivariate statistical analysis

Volume 2 Issue 1 - 2018


**Muhammad Usama Zafar, Waqar Ahmad**  
*Department of Environmental Science, Federal Urdu University, Pakistan*

**Correspondence:** Waqar Ahmad, Department of Environmental Science, Federal Urdu University, Pakistan, Email waqarevs@gmail.com

**Received:** November 09, 2017 | **Published:** January 04, 2018

## Introduction

The Indus is one of the largest river of Pakistan with drainage basin area (9,70,000 km<sup>2</sup>). It is nourished by glaciers and snows of the Himalayas, the Karakorum and the HinduKush that originate in the Indian State of Jammu and Kashmir and the Northern Areas of Pakistan. The parts of the HinduKush and the Karakorum ranges in the northern territory of Pakistan are drained by Gilgit River (current study area) which is bordered with Afghanistan and China in the North. The Hunza River basin is also a part of the present study area, actually the sub basin of the Gilgit River but owing to its substantial size and significance, it is considered as a separate basin. Like Gilgit River, Hunza River also drains the Karakorum Mountains consisting of a large glaciated area situated in the North. As far as water analysis is concerned, little work has been carried out in these areas. Mercury in Pan Amalgamation was found in high concentration in 37 river samples collected from 24 different sites of Gilgit and Hunza.<sup>1</sup> Physico-chemical properties along with some heavy metals (arsenic, chromium, copper, mercury and lead) of Handrap Lake and Nullah in Ghizer district were tested and no trace metals were found (World Wildlife Fund). As Northern areas of Pakistan are mountainous rural regions with a population of 900,000 living in villages typically encompassing 50-200 households.<sup>2</sup> In this region, the main source of water supply is from melting snows. It comes through channels (river lets) and small streams to the mouth of villages and is considered as the only supply of water for the villages as there are no wells and hand pumps which can substitute for this water. The water from melting snows run down the hills collecting various materials on its way and converted to turbulent mountainous streams.<sup>3</sup> The utilization of such water for domestic purpose may cause harmful diseases. So here we propose a study, where water samples were selected from 29 locations of Gilgit and Hunza Rivers (Figure 1) to check the quality of water from study areas. The following objectives were covered.



**Figure 1** Map representing twenty nine sampling sites from Gilgit and Hunza Rivers.

## Research objectives:

- To obtain water quality data from different location of Gilgit and Hunza Rivers.
- To examine physical and chemical contamination.
- To identify whether the concentration of dissolved solids are within the permissible limits of World Health Organization guidelines.
- To classify possible factors which are responsible for the variation in water quality of Gilgit and Hunza Rivers.

E. To use multivariate statistical techniques to study homogeneity and heterogeneity among sampling stations and to differentiate water quality variables according to temporal variation of Gilgit and Hunza Rivers.

## Materials and methods

### Sampling and on site evaluation

Twenty nine locations were chosen from Gilgit and Hunza Rivers for the estimation of physico-chemical parameters. The first 12 locations were from Gilgit River and the last 17 locations were

selected from Hunza River (Table 1). Samples were collected in the month of July 2012. The collection was carried out in such a way that samples did not get contaminated with other substances. At each site, surface water was collected and kept in polythene plastic bottles formerly washed in 10% nitric acid for 24 hours and rinsed with distilled water. 500 ml water was collected in each bottle and six parameters were noted at the site with the assistance of Sension 156 HACH potable multi-parameter, USA. The parameters were temperature (°C), pH, electrical conductivity (µS/cm), total dissolved solids (mg/L), salinity (%) and dissolved oxygen (mg/L). Two-three drops of nitric acid were introduced in the bottles so that chemical characteristics of the water could not be deteriorated.

**Table 1** Nearest town, elevation and map location of water collection of Gilgit and Hunza valleys

S. no.	Locations	Elevation in meters	Co-ordinates
1	Napura Baseen	1683	35o50N, 74o15E
2	Napura Baseen (spring)	1700	35o50N, 74o15E
3	Kargah	1674	35o50N, 74o15E
4	Gilgit city	1574	35o54N, 74o21E
5	Gilgit tap water	1574	35o54N, 74o21E
6	Jutial	1748	35o53N, 74o20E
7	Nomal	2507	36o08N, 74o12E
8	Nalter (spring)	2968	36o07N, 74o10E
9	Nalter (Lake)	2968	36o07N, 74o10E
10	Danyore	1580	36o08N, 74o51E
11	JuglotGah Nala	1610	36o09N, 74o51E
12	HaramoshNala	1600	35o07N, 74o08E
13	Aliabad Nala	1700	36o09N, 74o52E
14	Aliabad tapwater	1700	36o09N, 74o52E
15	Atabad	2400	36o20N, 74o52E
16	Gulmit	2412	36o20N, 74o52E
17	Hussaini	2433	36o20N, 74o52E
18	Ghalapur Nala	2500	36o36N, 74o51E
19	Khyber Nala	2678	36o34N, 74o48E
20	Passu	2700	36o46N, 74o90E
21	Gulkin Nala	2403	36o24N, 74o52E
22	Batura Glacier	2540	36o30N, 74o52E
23	Batura Lake	2540	36o30N, 74o52E
24	Shimshal River	2850	36o20N, 75o01E
25	Shimshal	2850	36o20N, 75o01E
26	Morkhun	2780	36o40N, 74o52E
27	Boiber tributary	3075	36o40N, 74o52E
28	BoiberNala	3075	36o40N, 74o52E
29	Sost River	3075	36o41N, 74o52E

### Laboratory and statistical analyses

In the laboratory, chemical analyses were carried out to find chloride, total hardness, total alkalinity, nitrate and sulphate. Chloride, total hardness, and total alkalinity were detected by titration method while nitrate and sulphate were evaluated by spectrophotometer; detail

of the methods is given in Table 2. For statistical analysis, “Minitab” version 11.12 was used. A Pearson Correlation matrix among all parameters was created to check whether the concentration of one parameter affect the concentration of another. Multivariate techniques (cluster analysis and principal component analysis) were applied to the datasets by normalizing the variables.

**Table 2** Analysis parameters and their analytical procedures

Serial no.	Variables	Abbreviation	Analytical method	Units
	physical	parameters		
1	Temperature	Temp	Sension HACH,156	oC
2	pH	pH	Sension HACH,156	No
3	Electrical Conductivity	EC	Sension HACH,156	µS cm <sup>-1</sup>
4	Total Dissolved Solids	TDS	Sension HACH,156	mg L <sup>-1</sup>
5	Salinity	Sal	Sension HACH,156	%
	Chemical	Parameters		
6	Dissolved Oxygen	DO	Sension HACH,156	mg L <sup>-1</sup>
7	Chloride	Cl-I	Titration (Silver Nitrate)	mg L <sup>-1</sup>
8	Total Hardness	Ca+Mg	Titration (EDTA)	mg L <sup>-1</sup>
9	Total Alkalinity	CO <sub>3</sub> +HCO <sub>3</sub>	Titration (H <sub>2</sub> SO <sub>4</sub> )	mg L <sup>-1</sup>
10	Sulphate	SO <sub>4</sub>	Spectrophotometer	mg L <sup>-1</sup>
11	Nitrate	NO <sub>3</sub>	Spectrophotometer	mg L <sup>-1</sup>

The procedures followed were those described by APHA.<sup>18</sup>

### Results

The values of twenty nine water samples with eleven parameters are presented in Table 3. World Health Organization limits for safe drinking water are shown at the bottom of Table 3. We took averages of all parameters from two rivers and then compared these averages. Temperature, pH, DO, salinity, chloride and nitrate had similar values. The other parameters TDS, conductivity, total alkalinity, total hardness and sulphate showed dissimilarities in results. The higher

values were obtained from Hunza River explaining Hunza River in part because of the Boiber tributary which exceeded the WHO (1993) approved thresholds of pH, conductivity, total alkalinity and total hardness as was the case with Boiber Nala tributary. The values of all parameters satisfied the corresponding thresholds (except in a few cases), whereas total alkalinity was found to be high. This means that all samples have high values of alkalinity. Total Dissolved Solids, D.O, salinity, chloride, sulphate and nitrate met the acceptable limits designed by WHO (1993).

**Table 3** Values of all sampling sites with eleven parameters from Gilgit and Hunza Rivers

S.no	Locations	Temperature	ph	D.o	T.d.s	Conductivity	Salinity	Chloride	Total Alkalinity	Total Hardness	Sulphate	Nitrate
1	Baseenpur	13.4	8.54	0.98	23	40.2	0	14	420	40	10	11.1
2	Baseenpur (Spring)	12.31	7.57	0.91	22.3	35.7	0.1	10	460	32	10	27.4
3	Kargah	13	7.6	1.8	24.3	38.2	0.1	10	700	32	11	10.9
4	Gilgit City	27.6	8.1	1.7	31	69.5	0	10	480	48	18	37.4
5	Gilgit tap water	25.2	7	1.65	35	83.6	0.1	9.93	1200	40	11	32.9
6	Jutial	12.1	7.48	1.89	27.6	43.6	0.1	12	560	28	19	12.4
7	Nomal	12.9	8.65	1.29	110	68.6	0.1	8	1020	120	22	5
8	Nalter (Spring)	7.4	8.2	2.2	24.8	38.5	0.1	10	360	132	11	36.7
9	Nalter lake	9.4	7	0.49	61.4	145.7	0	7.94	960	80	21	9.5

Table Continued....

S.no	Locations	Temperature	ph	D.o	T.d.s	Conductivity	Salinity	Chloride	Total Alkalinity	Total Hardness	Sulphate	Nitrate
10	Danyore	16.5	7.5	1.73	86.7	204	0	9.93	1020	100	37	21
11	JuglotGahNala	12.4	7.77	1.5	211	505	0.1	12	860	80	23	9.7
12	HaramoshNala	14.2	7.4	1.74	102.2	240	0.1	11.91	900	140	43	26.3
13	Aliabad Nala	12.3	7.1	1.63	36.5	86.1	0	9.93	840	40	18	22.5
14	Aliabad Tawafer	13.2	7.42	1.48	47.7	115.4	0	12	920	220	119	18.5
15	Atabad	10.2	7.4	2.01	69.7	164	0	7.94	700	100	21	29.4
16	Gulmit	12.1	7.86	1.47	85.7	20.3	0	12	980	72	32	18.5
17	Hussaini	10.2	7.75	1.41	71.21	168.9	0	6	820	100	24	16.4
18	GhalapurNala	11.8	8.49	1.36	166.9	70.4	0	12	700	116	9	13.7
19	Khyber Nala	16.7	8.54	1.85	171.8	316	0.1	12	740	172	78	10.8
20	Passu	11.3	7.2	1.7	52.6	124.4	0	11.9	880	80	34	33.2
21	GulkinNala	10.8	8.37	2.07	19.3	36.3	0	12	540	36	9	17.6
22	Batura Glacier	11.1	8.55	2.02	52.3	96.2	0	12	600	48	19	33.8
23	Batura Lake	11.1	8.3	1.98	119.9	219	0.1	10	1000	168	46	31.5
24	Shimshal River	10.9	7	2.01	78.3	184.8	0	14.89	800	120	48	36.5
25	Shimshal	11	7.2	1.92	181	423	0.1	9.93	860	240	91	39.9
26	Morkhun	13.3	8.75	1.26	182	426	0.1	8	840	136	79	36
27	BoiberTributery	14.5	8.55	1.46	277	513	0.3	10	820	176	86	12.1
28	BoiberNala	13.2	8.73	1.7	155	290	0.1	8	560	196	60	42.7
29	Sost River	13.5	8.26	1.7	109.5	187.5	0.1	14	900	140	50	12.4
	WHO (1993)	No guideline	6.5-8.5	No guideline	500 mg/L	400 µS/cm	No guideline	250 mg/L	250 mg/L	150 mg/L	500 mg/L	50 mg/L

**The Pearson correlation matrix of all parameters**

Correlations of all parameters from 29 samples of Gilgit and Hunza Rivers are shown in Table 4. Eleven parameters from 29 locations from two rivers were analyzed for correlation. The significance level at degrees of freedom 27 is checked using r Table in the following order.

1. Greater than 0.349 but less than 0.449 (p<0.05);
2. Greater than 0.449 but less than 0.554 (p<0.01);
3. Greater than 0.554 (p<0.001)

Temperature did not illustrate any significant correlation with all parameters. It was negatively correlated with pH, D.O, TDS, conductivity and total hardness and positively correlated with salinity, chloride, total alkalinity, sulphate and nitrate. pH was found to be significantly correlated with total dissolved solids and total alkalinity at (p<0.05) and positively correlated with all parameters except chloride and nitrate. No significant correlation was found between dissolved oxygen and all other parameters. It was positively correlated with three parameters and negatively correlated with the other parameters. TDS showed significant correlations with four parameters i.e. conductivity, salinity, total hardness and sulphate at (p<0.001).

Conductivity also expressed strong significant positive correlations with salinity, hardness and sulphate and weak positive correlation with total alkalinity. Positive significant correlations were observed among salinity, total hardness and sulphate at (p<0.05) level and chloride did not show any significant correlation with any other parameter. Total alkalinity and total hardness indicated significant positive correlation with sulphate at (p<0.05) and (p<0.001) respectively. There was no significant correlation between sulphate and nitrate. This showed that total hardness and sulphate are the two parameters which are highly correlated with most of the other parameters. The characteristics of three groups derived from agglomerative cluster analysis are presented (Table 5 & Figure 2). Temperature was similar in all clusters. The pH of the water was slightly more alkaline for cluster I and III. Dissolved oxygen was almost similar in all groups whereas total dissolved solids were remarkably higher in group III. Conductivity of the samples showed greater values for groups II and III. Salinity was found to be higher in cluster III while chloride was little bit higher in group I. Total alkalinity was found to be low in cluster I as compared to the other two clusters. Total hardness of group III was higher than the other two groups. Sulphate and nitrate showed greater values for group III. Principal Component analysis (PCA) was applied on normalized data sets (11 variables) separately for 29 locations (Figure 3) to find similarities or dissimilarities among variables. PCA of the

data sets produced first five PCs with eigen values > 1 explaining 80.8 percent of total variance with respect to water quality data. Eigen value measures the significance of the factor and values greater than one are considered as significant contribution to the analysis.<sup>4,5</sup> Six parameters (temperature, pH, D.O, salinity, chloride and nitrate) formed a close association with sulphate. Hardness and TDS centroids occurred close together but were distinct from the previous 7 parameters. The conductivity and alkalinity centroids were found to be located quite apart from the other parameters in ordination space, indicating they have the least correlation with the other parameters. The scree plot (Figure 4) was used to explain the number of PCs to be retained in order to understand the fundamental data structure.<sup>6</sup> The scree plot of the present study showed that first five PCs have eigen


values greater than one (Figure 4), together the first four components explains 70.4% of the total variance inherent in the data set. Among three PCs, PC1 explaining 33% of total variance has weak positive loadings on TDS, conductivity, salinity, total hardness and sulphate. PC2 (14.5% of total variance) has moderate positive loadings on D.O and nitrate which may indicate that increased concentration of nitrate will presumably increase the consumption of dissolved oxygen<sup>7</sup> PC3 with 13.4% of total variance caused moderate positive loading over pH and alkalinity. Overall analysis of PCA shows that variations of water quality are related to dissolved solids and salts (e.g., sulphate, nitrate, total alkalinity and total hardness) which may change the pH and dissolved oxygen concentration of water of the study area.

**Table 4** Correlation matrix among all parameters


	Temperature	ph	D.o	Tds	Conductivity	Salinity	Chloride	Total Alkalinity	Total Hardness	Sulphate
ph	-0.006									
D.o	-0.067	0.2								
Tds	-0.039	0.379	0.216							
Conductivity	-0.01	0.149	-0.134	0.877						
Salinity	0.088	0.288	-0.165	0.592	0.561					
Chloride	0.014	-0.023	0.131	-0.093	-0.137	-0.125				
Total Alkalinity	0.155	-0.358	-0.071	0.289	0.273	0.067	-0.133			
Total Hardness	-0.166	0.189	0.039	0.638	0.603	0.354	-0.099	0.253		
Sulphate	0.001	0.083	-0.18	0.565	0.631	0.349	0.032	0.301	0.843	
Nitrate	0.126	-0.122	-0.152	-0.089	0.084	-0.09	-0.131	-0.156	0.205	0.142

**Table 5** Characteristics of three groups derived from Ward's clustering of the water quality variables of the samples collected from 29 locations


Water quality variables	Cluster I (1,2,4,8,3,18,6,21,22)	Cluster II (5,7,16,9,10,23,12,29,14,13,20,15,17,24)	Cluster III (11,27,25,26,19,28)
Temperature	13.28	13.07	13.51
ph	8.1	7.56	8.25
D.o	1.66	1.59	1.61
T.d.s	43.5	76.17	196.3
Conductivity	52.07	143.73	412.16
Salinity	0.04	0.035	0.13
Chloride	11.33	10.45	9.98
T.Alkalinity	535.56	924.2	780
T.Hardness	56.89	108.5	166.66
Sulphate	12.89	37.5	69.5
Nitrate	22.33	22.4	25.2



**Figure 2** Dendrogram resulting from Ward's clustering of 29 samples collected from Gilgit and Hunza Rivers.



**Figure 3** Principal Component analysis (PCA) based on eleven parameters of water samples collected from Gilgit and Hunza valleys.



**Figure 4** Scree plot of 29 water samples with eleven parameters.

## Discussion

The present study is the first comprehensive study in which we investigated water resources (river, stream, lakes and nullahs) of Gilgit and Hunza valleys. Mean conductivity was satisfactory among all samples except that water from Juglot Gah Nala exhibited high conductivity. This sample also had a high amount of TDS, perhaps this might be due to the presence of minerals in the water originating from solid rocks and glaciers. Of all parameters, only total alkalinity crossed the tolerable threshold of drinking water. The rocks of these valleys contain high amounts of carbonate and bicarbonate which mix with the water passing through it, causing the high water alkalinity. Correlation analysis results explained that there are specific relationship pattern among pH, TDS, conductivity, total hardness and sulphate and these results are also confirmed by principal component

analysis. Correlation analysis results showed the direct significant positive relationship ( $P < 0.001$ ) between electrical conductivity with total hardness and sulphate. It means that total hardness and sulphate drive the conductivity of surface water of Rivers as reported in ground water from urban areas of Karachi.<sup>8</sup> The cluster analysis of the overall data set showed three major groups while the discriminating variable for the groups were six variables (temperature, pH, D.O, salinity, chloride and nitrate) which showed homogeneity within groups but heterogeneity between the three groups. TDS and hardness also exhibited considerable differences between the groups. Likewise, vast differences in the mean values were observed between the clusters for conductivity and total alkalinity. Cluster analysis highlights that at present, most of the samples collected from Hunza River have high values of TDS, conductivity, total alkalinity and hardness which may indicate the high concentration of salts in Hunza River as compared to Gilgit River. In the present study, we found the lowest temperature (7.4°C and 9.4°C) from Nalter site (lake and spring respectively). The low temperature of the water is due to the fact that the water which we collected from Nalter, was coming from glaciers. Similar temperature (8°C) was also observed by Islamuddin (2010) who worked over Nalter Lake. Water temperature from Nomal valley, located at lower height (2507 m) than Nalter valley (2968 m) was 13°C in current study whereas<sup>9</sup> described the temperature of the same valley about 25°C due to the difference in collection seasons. All the physico-chemical properties from the present study and the Islamuddin<sup>10</sup> study are concurrent with each other and are in accordance with the limits of WHO (1993) except total alkalinity (Carbonate and bicarbonate) were relatively higher in both studies. Our physico-chemical characteristics also match the findings of Jhelum River, District Muzaffarabad Azad Kashmir study<sup>11</sup> and with that of some studies of lower Indus Basin<sup>12-14</sup> but total alkalinity was lower in their studies as compared to current studies. The concentrations of all chemical parameters were found within permissible limits in samples with the exception of total alkalinity in the samples. The alkaline water has a bitter taste and slippery feel. High alkaline water like in the current study can cause drying of skin. Alkalinity is important for fish and aquatic life because it acts as a buffer against rapid pH changes.<sup>15,16</sup> High alkalinity is also important in agricultural activity. Use of high alkaline water affects plant growth because excessive salts raise osmotic pressure in soil solution and reduce water availability. This high alkalinity results in lower leaf-area index.<sup>17</sup> Finally it is concluded that physico-chemical properties of surface water of study areas are within drinking permissible limits of WHO (1993) but the water is considered as bicarbonate type. It is still important to explore the bacterial contamination of the study area but it is not a current study objective. However Nano-filtration techniques should be installed at the mouth of water supply to reduce the total alkalinity of Gilgit and Hunza Rivers.<sup>18</sup>

## Acknowledgments

None.

## Conflict of interest

Authors declare there is no conflict of interest in publishing the article.

## References

1. Biber K. Mercury fate and transport in Hunza and Gilgit rivers, Northern Areas, Pakistan-A numerical modeling approach. *GSA annual meeting in Minneapolis*. 2011;43(5):241–328.

2. Nanan D, White FI, Azam H Afsar, et al. Evaluation of water, sanitation and hygiene education intervention on diarrhea in northern Pakistan. *Bull World Health Organ.* 2003;81(3):160–165.
3. McCarrison R, Captain MB. Observation on endemic goiter in Chitral and Gilgit Valleys. *Agency Surgeon Gilgit Kashmir.* 1906;1–38.
4. Alvin CR. *Methods of Multivariate Analysis.* John Wiley & Sons INC, USA; 2002. p. 1–727.
5. Otto M. Multivariate methods. In: Kellner R, Mermet JM, Otto M, Widmer HM (Eds.), *Analytical Chemistry, Wiley-VCH, Germany;* 1998. p. 1–916.
6. Vega MR, Pardo E Barrado, Deban L. Assessment of seasonal and polluting effects on the quality of river water by exploratory data analysis. *Water Research.* 1998;32(12):2581–3592.
7. Singh KP, Malik A, Mohan D, et al. Mutivariate statistical techniques for the evaluation of spatial and temporal variations in water quality of Gomti River (India): A case study. *Water Res.* 2004;38(18): 3980–3992.
8. Farooq MA. The study of ground water pollution resulting from storm water infiltration in Urban Areas of Karachi. PhD Dissertation. Federal Urdu University of Arts, *Science and Technology Karachi.* 2008.
9. Ahmed K, Shah S. Physical and microbial assessment of drinking water of Nomal Valley, Northern Areas, Pakistan. *Pak J of Zool.* 2007;39(6):367–373.
10. Islamuddin. Hydro-chemical investigations of high altitude alpine lakes of Gilgit and Ghizer districts, Northern Areas of Pakistan. University of Peshawar, Pakistan; 2010. p. 1–246.
11. Sarwar S, Ahmed F, Khan J. Assessment of the Quality of Jhelum River water for irrigation and drinking at Distt. Muzaffarabad Azad Kashmir. *Sarhad J Agric.* 2007;23(4):1041–1046.
12. Farooq MA. Surface water quality assessment and apportionment of pollution sources of lower Indus Basin from Kashmore to Keti-Bunder using multivariate statistical technique- A case study. Research Project, Report submitted to Higher Education Commission Pakistan; 2012.
13. Pirzada TM, Talpur MA, Rajper AD, et al. Quality Assessment of Potable Water of Khairpur City. *J Chem Soc Pak.* 2011;33(4):604–611.
14. Lashari KH, Korai AL, Sahoto GA, et al. Limnological studies of Keenjhar Lake, Distt. Thatta, Sindh, Pakistan. *Pak J Ana Env Chem.* 2009;10(1):39–47.
15. Benjamin M. *Water Chemistry.* McGraw-Hill, USA; 2002.
16. Hemond H, Frechner-Levy E. *Chemical Fate and Transport in the Environment.* 2nd ed. Elsevier Science, USA; 2000.
17. Tyagi NK. Managing saline and alkaline water for high productivity. In: Kijne JW, Barker R, Molden D, editors. *Water productivity in agriculture: Limits and opportunities for improvement.* Central Soil Salinity Research Institute, India; 2003. p. 1–19.
18. APHA, AWWA, WEF. *Standard Methods for the Examination of Water and Waste Water,* 21st edn. American Public Health Association, USA; 2005.