

Vulvar Procidence of Intestinal Anses: A Rare Complication of Uterine Rupture

Keywords: Vulvar procidence; Intestinal anses; Uterine rupture; Placental retention; Hemorrhagic shock

Clinical Images

It is a 20 year old paucipar, received in our structure after delivery in a health post, for placental retention, vulvar externalization of intestinal loops. Childbirth was performed by low voice one hour before with birth of a macrosome (4000g), fresh stillborn. There was dystocia of the shoulders to expulsion, managed by uterine expressions. There was no history of Caesarean section. On arrival, there was a state of hemorrhagic shock. The gynecological examination found vulvar bleeding, a uterus at the level of the umbilicus, a procidence of maternal intestinal loops through the vulva. The biological assessment found a hemoglobin level at 7.5g/dl, GB 9.103/ul, platelets at 42.103/ul, serum creatinine at 16.78 mg/l.

Emergency laparotomy revealed a left uterine rupture extending up to the left appendix, where the slender loops entered the vagina with an externalized necrotic portion (Figure 1). The placenta sat in the abdomen. Gestures: hysterectomy of interannexal haemostasis and intestinal resection of 30 cm with ileo-ileal anastomosis.



Figure 1: Vulvar Procidence of Intestinal Anses.

Clinical Images

Volume 4 Issue 2 - 2018

Thiam M^{1,2*}, Gueye I^{1,2}, Dieme Mef^{3,4}, Niang MM^{3,4}, Diouf AA^{3,4}, Ba PA^{1,5}, Gassama O^{3,5} and Cisse ML^{1,2}

¹Department of Health Sciences, University of Thiès, Senegal

²Department of Obstetrics and Gynecology, Thies Regional Hospital, Senegal

³Gynecological and Obstetrical Clinic, The Dantec University Teaching Hospital of Dakar, Senegal

⁴Cheikh Anta Diop University of Dakar, Senegal

⁵Department of Surgery and Specialties, Thies Regional Hospital, Senegal

***Corresponding author:** Mariétou Thiam, Assistant Professor at faculty of Health sciences, University of THIES, Obstetrician and Gynecologist at the THIES Regional Hospital, Senegal, Email: maricoulibaly10@gmail.com

Received: February 07, 2018 | **Published:** February 12, 2018

The patient had been resuscitated post-operatively for 8 days.

She had received 7 iso-rhesus iso-group blood bags. She was out at J13 postoperatively.

Acknowledgement

The corresponding author is the guarantor of submission.

Conflict of Interest

Authors declare no conflict of interest.