

# Increased expression of *Toxoplasma gondii* gra1 suppresses host cell apoptosis

## Abstract

Suppression of host cell apoptosis enable *Toxoplasma gondii* to proliferate, but the mechanism is not well understood. We explore the relationship between the expression of *T. gondii* dense granule protein 1 (GRA1) and host HeLa cell apoptosis. We inserted the *T. gondii* gra1 coding gene into a pET32a vector and produced recombinant GRA1 (rGRA1) in *E. coli* Rosetta strain. We then immunized rabbits with rGRA1 to produce anti-GRA1 serum and infected HeLa cells with *T. gondii* tachyzoites. We determined the expression of *T. gondii* GRA1 by Western blotting with anti-GRA1 serum and determined apoptosis of HeLa cells via the Annexin V-FITC/PI method. All the experiments were repeated for three times in the same condition. The expression of GRA1 decreased gradually after *T. gondii* infection. The rate of HeLa cell apoptosis increased more rapidly in the infected cells than in the uninfected cells. Our results suggest that the suppression of host cell apoptosis is related to the expression of *T. gondii* GRA1 expression.

**Keywords:** *toxoplasma gondii*, GRA1, apoptosis, ROPs, GRAs, STAT

Volume 4 Issue 5 - 2017

Liang Wu,<sup>1</sup> Jin Shen,<sup>1</sup> Chenyu Tang,<sup>2</sup> Xugan Jiang,<sup>1</sup> Jiajian Wang,<sup>2</sup> Xiaoling Jin,<sup>2</sup> Jinbo Qiu,<sup>2</sup> Shengxia Chen,<sup>1</sup> Jianping Cao<sup>3</sup>

<sup>1</sup>School of Medicine, Jiangsu University, China

<sup>2</sup>Jingjiang College of Jiangsu University, China

<sup>3</sup>National Institute of Parasitic Diseases, China

<sup>#</sup>Liang Wu and Jin Shen contribute equally to this work

**Correspondence:** Shengxia Chen, School of Medicine, Jiangsu University, Zhenjiang 212013, China, Tel +86 13952815350, Email chensxia@ujs.edu.cn

**Received:** April 25, 2017 | **Published:** May 16, 2017

## Introduction

*Toxoplasma gondii* is an obligate intracellular protozoan parasite. In immunocompromised hosts, such as AIDS patients, it becomes an opportunistic pathogen causing severe toxoplasmosis.<sup>1</sup> Apoptosis is a common biological phenomenon that removes damaged or unwanted cells during development, ensuring tissue homeostasis in multi-cellular organisms.<sup>2</sup> Cell apoptosis also plays a major role in innate and adaptive immune defense by controlling and eradicating invasive pathogens.<sup>3</sup> *T. gondii*, however, can actively invade any nucleated host cell and, by down-regulating apoptosis in the cell, can evade elimination by the host's defense system.<sup>4</sup> Although the mechanism of this down-regulation is not yet understood, it is known that *T. gondii* secreted proteins including rhoptries (ROPs) and dense granule proteins (GRAs) play an important role in this regulation.<sup>5</sup> Many studies have examined the function of ROPs, but few have studied the function of GRAs.<sup>6</sup> In this study, we investigate variation in the expression of GRA1 and in the rate of cell apoptosis during *T. gondii* invasion into HeLa cells.

## Material and methods

### Ethics statement

All animal experiment in the manuscript were reviewed in advance by the Laboratory Animal Management Committee of Jiangsu University and also met the guidelines of the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

### Cell culture and *T. gondii*

The HeLa cell line was stored in liquid nitrogen in our laboratory and cultured in DMEM medium (HyClone, Argentina) and incubated with 10% fetal bovine serum at 37°C and at 5% CO<sub>2</sub> (Tianhang Biological Technology Co., Ltd, China). The cells were stripped from the cell culture flask with a 0.25% trypsin (GBICO, USA) and 0.02% EDTA-Na<sub>2</sub> solution. The passage operation was repeated every 2-3 days. *T. gondii* RH strain tachyzoites were maintained in HeLa cells. Before

infection with *T. gondii* tachyzoites for 12h, the HeLa cell medium was changed to a DMEM medium with 2% fetal bovine serum. The *T. gondii* RH strain tachyzoites were added into the culture medium and this was changed to new fresh DMEM medium (with 2% fetal bovine serum) 4hours after infection. After HeLa cell lysis, the tachyzoites were collected by centrifugation and purified using a 3-mm filter membrane.<sup>7</sup>

### Cloning of the GRA1 gene and anti-GRA1 serum production

The purified tachyzoites were used for total RNA extraction performed using the RNeasy Total RNA Isolation Kit according to the manufacturer's instructions (Generay Biotech Co., Ltd. China). The GRA1 gene used for His-GRA1 recombinant protein production was amplified from cDNA by PCR assay. The GRA1 primers 5'-CGGATCCCCGAAGGCGCGACAACCAG-3' (sense) and 5'-GAATTCTACTCTCTCTCTCTGTTAGGAACCCAATGTC-3' (anti-sense) yielded a 495-nucleotide product specific to the GRA1 coding gene (GenBank No. HM067753.1). The PCR product was digested with BamHI and EcoRI, ligated into a pET32a vector using T4 DNA ligase (TAKARA, Dalian, China), and transformed into an *E. coli* Rosetta strain. The right directional clone was confirmed by sequencing. The His-GRA1 recombinant protein was expressed in *E. coli* Rosetta strain that had been induced at 30°C for 10h with 0.4mM isopropyl-β-D-thiogalactopyranoside (IPTG). The bacterial pellets were re-suspended in lysis buffer (Tris-HCl 10mM [pH 8.0], NaH<sub>2</sub>PO<sub>4</sub> 100mM, Urea 8M). Cellular debris was removed by centrifugation at 16,000 g for 30min. The cell-free extract was used for rHis-ACP purification by Ni-NTA Agarose (QIAGEN, Hilden, Germany) according to the manufacturer's instructions. This His-ACP recombinant protein was identified by SDS-polyacrylamide gel electrophoresis staining with Coomassie blue.

The specific anti-serum of GRA1 was obtained from New Zealand rabbits (male, 3kg, Laboratory Animal center of Jiangsu University) by three subcutaneous immunizations. The first injection comprised

1mg of recombinant GRA1 in Freund's complete adjuvant, with two further injections also with Freund's incomplete adjuvant administered at three week intervals. The rabbit was then bled and the specific anti-serum collected for the Western blotting assay.

### Western blotting assay

Freshly released tachyzoites were boiled in SDS-PAGE sample buffer and separated on 12% polyacrylamide gels utilizing the Laemmli discontinuous buffer system.<sup>8</sup> Proteins were transferred to nitrocellulose membranes (BOSTER, China), and the nitrocellulose strips were saturated for 1h in 5% non-fat milk in 15mM Tris-HCl (pH 8.0), 150mM NaCl, and 0.05% Tween 20 (TNT). They were then incubated with anti-GRA1 serum diluted 1:2,000 in TNT for 1h. After washing, the strips were incubated with goat anti-rabbit IgG conjugated to HRP (BOSTER, China) and detected using the ECL (BOSTER, China) plus Western blotting system.

### *T. gondii* GRA1 expression

The parasites ( $5 \times 10^6$ /well) were added to confluent HeLa cell monolayers in a 6-well plate (ratio of cells to parasites equals one). After infection for 1h, all the infected HeLa cells were completely washed with PBS to eliminate parasites that had not invaded cells. We used anti-GRA1 serum and anti-*T. gondii* actin mAb (in mouse ascitic fluids, obtained from Professor Dominique of the University of Geneva, Switzerland) to determine the expression of GRA1 during infection.

### Analysis of apoptosis

The parasites ( $5 \times 10^6$  per well) were added to confluent HeLa cell monolayers in a 6-well plate (at a cell to parasite ratio equal to one). One hour after infection, all infected HeLa cells were completely washed with PBS to eliminate parasites that had not invaded cells. After cell detachment, HeLa cells were stained with Annexin V using the Annexin V-FITC/PI kit (Vazyme, Nanjing, China). Apoptotic cells were identified and quantified by flow cytometry (FACSCalibur, BD Company, USA). Briefly, HeLa cells ( $1 \times 10^6$  per well) were washed in PBS and incubated with 1× binding buffer, propidium iodide (PI) and Annexin V-FITC for 30min at room temperature. The apoptotic cells were analyzed on a fluorescence-activated cell sorter.


### Statistical analyses

Statistical analyses were performed in SPSS. Student's t tests were performed under the assumption of equal variance and using a two-tailed test, where  $P \leq 0.05$  was considered significant.

## Results


### Expression of rGRA1 and anti-GRA1 serum production

Recombinant plasmids containing GRA1 gene were confirmed by BamHI/EcoRI restriction enzyme analysis. The band corresponding to the 495bp product was visualized on a 1.5% agarose gel (Figure 1). As shown in the SDS-PAGE gel, the rGRA1 expression was induced by IPTG and its molecular weight was about 45kDa. This is consistent with the expected molecular mass given that the expression vector produced a recombinant protein fused to a 20kDa His-thioredoxin tag (Figure 2). In the Western blotting assay, the GRA1 (25kDa) in tachyzoite lysis could be recognized by the rabbit anti-GRA1 serum, while the rabbit anti-GRA1 serum did not react with the HeLa cell lysis (Figure 3).




**Figure 1** The GRA1 gene product amplified by RT-PCR from *T. gondii* (RH strain).

M, molecular weight marker; Lane 1, GRA1 gene; Lane 2, negative control.



**Figure 2** SDS-PAGE analysis of rGRA1 expressed in *E. coli*.

The *E. coli* lysates were electrophoresed on 12% SDS-PAGE and stained with Coomassie brilliant blue R-250. M: molecular weight marker; Lane 1: IPTG-induced *E. coli* with pET-32a; Lane 2: Purified rGRA1. The rGRA1 fused with His-thioredoxin tag (45kDa) is shown with an arrow.



**Figure 3** Western blotting analysis of expressed rGRA1.

Lane 1: HeLa cells; Lane 2: *T. gondii*.


### GRA1 expression in *T. gondii* infection

The expression of GRA1 decreased gradually following the infection of HeLa cells by *T. gondii*. One hour post-infection the expression of GRA1 peaked and then decreased rapidly to its lowest level at 36 post-infection (Figure 4).

### HeLa cell apoptosis


The rate of HeLa cell apoptosis increased with time for both infected and uninfected cells, but did so more rapidly in the uninfected

group. At eight hours post-infection, there was no significant difference in apoptosis rates between infected and uninfected individuals. However, by 12 hours post-infection, the rate of apoptosis of uninfected cells had increased dramatically. At 36 hours post-infection, we observed the greatest difference in the rate of apoptosis in these two groups (Figure 5).



**Figure 4** Western blotting analysis of *T. gondii* GRA1 expression over 36h.

Actin: *T. gondii* actin expression; GRA1: *T. gondii* GRA1 expression.



**Figure 5** Rates of apoptosis of HeLa cells infected with *T. gondii* and uninfected cells over a 36-hour period.

## Discussion

The obligate intracellular protozoan *T. gondii* has evolved an intimate relationship with its host that extends to the cellular and molecular levels.<sup>9</sup> This pathogen requires an appropriate host cell environment for its replication and is able to modify its host cell functions.<sup>10</sup> Intuitively, the active modification of host cell growth must be a complex process, as host cells are not inherently programmed to provide an environment conducive to pathogens.<sup>11,12</sup> Host cells have evolved primary lines of defense as countermeasures to pathogen invasion, establishment, and replication.<sup>13,14</sup> Defenses to limit pathogen growth include apoptosis, reactive oxygen and nitrogen intermediates.<sup>15</sup> Previously, we reported that *T. gondii* infection suppressed apoptosis of HeLa cells for 36 hours post-infection in uninfected cells and in cells treated with actinomycin D.<sup>16</sup> *T. gondii* uses a mixture of specialized parasite secreted proteins, including ROPs, GRAs and MICs to invade cells.<sup>17,18</sup> Many of these parasite secreted proteins are anchored on the parasitophorous vacuole membrane (PVM) and can interact with the host cells directly. The kinase activity ROPs, such as ROP2,<sup>19</sup> ROP16<sup>20</sup> and ROP18,<sup>21</sup> have been exhaustively studied and shown to down-regulate host cell apoptosis.<sup>22</sup> Moreover, kinase activity ROPs can affect the activation of signal transducer and activator of transcription (STAT) signaling pathways, leading to a downstream modulation of the secretion of IL-12 in host cells.<sup>23,24</sup>

GRA1 belongs to a dense granule protein family consisting of 12 distinct polypeptides that are stored within dense granules of *T. gondii*.<sup>25</sup> During cell invasion, GRA1 was exocytosed into the parasitophorous vacuole (PV) and associated with PVM. Unlike *T. gondii* ROPs, the role of GRAs is not yet clearly understood. Indeed, GRA1 could activate TGF- $\beta$  transcription through phosphorylation and activate apoptosis in monocytes.<sup>26</sup> The role of GRA1 in human cell apoptosis has rarely been studied. In this study, we showed that the expression of GRA1 decreased gradually 36 hours post-infection, while HeLa cell apoptosis increased more rapidly in infected cells compared to uninfected cells. These findings suggest that GRA1 suppresses HeLa cell apoptosis, in a manner similar to ROPs.

## Conclusion

We suggest that GRA1 is anchored in the PVM and along with other secreted proteins, such as ROP2, could down-regulate human cell apoptosis. This would provide a longer growth period for *T. gondii* in host cells and also benefit the parasite by avoiding elimination of infected cells.

## Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (No. 81301453), the Laboratory of Parasite and Vector Biology of China, MOPH (No. WSBKTKT201302), the China Postdoctoral Science Special Foundation (No. 2015T80518), the China Postdoctoral Science Foundation (No. 2014M561598), Jiangsu Postdoctoral Science Foundation (No. 1402171C), and the Senior Talent Studying Initial Funding of Jiangsu University (No. 13JDG023, 13JDG127).

## Conflict of interest

The author declares no conflict of interest.

## References

- Pereira-Chioccola VL, Vidal JE, Su C. *Toxoplasma gondii* infection and cerebral toxoplasmosis in HIV-infected patients. *Future Microbiol.* 2009;4(10):1363–1379.
- Feustel SM, Meissner M, Liesenfeld O. *Toxoplasma gondii* and the blood-brain barrier. *Virulence.* 2012;3(2):182–192.
- Van de Veerdonk FL, Kullberg BJ, Netea MG. Pathogenesis of invasive candidiasis. *Curr Opin Crit Care.* 2010;16(5):453–459.
- Luder CG, Campos-Salinas J, Gonzalez-Rey E, et al. Impact of protozoan cell death on parasite-host interactions and pathogenesis. *Parasit Vectors.* 2010;3:116.
- Treeck M, Sanders JL, Elias JE, et al. The phosphoproteomes of *Plasmodium falciparum* and *Toxoplasma gondii* reveal unusual adaptations within and beyond the parasites' boundaries. *Cell Host Microbe.* 2011;10(4):410–419.
- Bradley PJ, Sibley LD. Rhoptries: an arsenal of secreted virulence factors. *Curr Opin Microbiol.* 2007;10(6):582–587.
- Wu L, Chen SX, Jiang XG, et al. Separation and purification of *Toxoplasma gondii* tachyzoites from *in vitro* and *in vivo* culture systems. *Exp Parasitol.* 2012;130(1):91–94.
- Cleveland DW, Fischer SG, Kirschner MW, et al. Peptide mapping by limited proteolysis in sodium dodecyl sulfate and analysis by gel electrophoresis. *J Biol Chem.* 1977;252(3):1102–1106.
- Laliberte J, Carruthers VB. Host cell manipulation by the human pathogen *Toxoplasma gondii*. *Cell Mol Life Sci.* 2008;65(12):1900–1915.

10. Sinai AP, Roepe PD. Autophagy in Apicomplexa: a life sustaining death mechanism? *Trends Parasitol.* 2012;28(9):358–364.
11. Lim DC, Cooke BM, Doerig C, et al. *Toxoplasma* and *Plasmodium* protein kinases: roles in invasion and host cell remodelling, *Int J Parasitol.* 2012;42(1):21–32.
12. Nelson MM, Jones AR, Carmen JC, et al. Modulation of the host cell proteome by the intracellular apicomplexan parasite *Toxoplasma gondii*. *Infect Immun.* 2008;76(2):828–844.
13. Randow F, Munz C. Autophagy in the regulation of pathogen replication and adaptive immunity. *Trends Immunol.* 2012;33(10):475–487.
14. Hood MI, Skaar EP. Nutritional immunity: transition metals at the pathogen–host interface. *Nat Rev Microbiol.* 2012;10(8):525–537.
15. Hohl TM, Feldmesser M. *Aspergillus fumigatus*: principles of pathogenesis and host defense. *Eukaryot Cell.* 2007;6(11):1953–1963.
16. Shen J, Chen YT, Wu L, et al. Effect of *Toxoplasma gondii* tachyzoite infection on HeLa cell apoptosis. *J Jiangsu Univ.* 2014;24(5):399–402.
17. Paredes–Santos TC, de Souza W, Attias M. Dynamics and 3D organization of secretory organelles of *Toxoplasma gondii*. *J Struct Biol.* 2012;177(2):420–430.
18. Nadipuram SM, Kim EW, Vashisht AA, et al. *In vivo* biotinylation of the *Toxoplasma* parasitophorous vacuole reveals novel dense granule proteins important for parasite growth and pathogenesis. *MBio.* 2016;7(4):e00808–e00816.
19. Sinai AP, Joiner KA. The *Toxoplasma gondii* protein ROP2 mediates host organelle association with the parasitophorous vacuole membrane. *J Cell Biol.* 2001;154(1):95–108.
20. Jensen KD, Hu K, Whitmarsh RJ, et al. *Toxoplasma gondii* rhoptyr 16 kinase promotes host resistance to oral infection and intestinal inflammation only in the context of the dense granule protein GRA15. *Infect Immun.* 2013;81(6):2156–2167.
21. Cheng L, Chen Y, Chen L, et al. Interactions between the ROP18 kinase and host cell proteins that aid in the parasitism of *Toxoplasma gondii*. *Acta Trop.* 2012;122(3):255–260.
22. Kemp LE, Yamamoto M, Soldati–Favre D Subversion of host cellular functions by the apicomplexan parasites. *FEMS Microbiol Rev.* 2013;37(4):607–631.
23. Schneider AG, Abi Abdallah DS, Butcher BA, et al. *Toxoplasma gondii* triggers phosphorylation and nuclear translocation of dendritic cell STAT1 while simultaneously blocking IFN $\gamma$ –induced STAT1 transcriptional activity. *PLoS One.* 2013;8(3):e60215.
24. Asai T, Tena G, Plotnikova J, et al. MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature.* 2002;415(6875):977–983.
25. Nam HW. GRA proteins of *Toxoplasma gondii*: maintenance of host–parasite interactions across the parasitophorous vacuolar membrane. *Korean J Parasitol.* 2009;47:S29–S37.
26. D’Angelillo A, De Luna E, Romano S, et al. *Toxoplasma gondii* Dense Granule Antigen 1 stimulates apoptosis of monocytes through autocrine TGF– $\beta$  signaling. *Apoptosis.* 2011;16(6):551–562.