

# Effect of Zinc Oxide and Titanium Dioxide Nanoparticles on Supported Lipid Bilayers

## Research Article

## Abstract

Nanoparticles (NP) are broadly exploited in biomedical sciences in order to develop various methods of targeted drug delivery, novel biosensors and new therapeutic pathways. However, relatively little is known in the negotiation of NPs with complex biological environments. NP interaction with cell membranes can damage the cell membrane and cause toxicity; therefore, examining interactions between NPs and cell membranes is crucial to understanding NP toxicity mechanisms and the development of safe and non-toxic NP-based commercial and therapeutic applications. To gain a physical understanding of NP-membrane interactions, we used a simplified system built of well-defined synthetic lipid bilayers and NP. The interaction of nanoparticles (NPs) with supported lipid bilayers (SLB) can lead to structural modification of the SLB and affect the structure and dynamic of lipids. In this work  $\text{TiO}_2$  and ZnO nanoparticles were chosen because of wide applications and usage in industry of papers, inks, medicines, food products, cosmetics, toothpastes and skin care products and among others. Therefore, a better understanding of the interactions between NP and lipid membrane may help to better clarify the potential risk of NPs. The interaction between ZnO and  $\text{TiO}_2$  NPs and lipid membranes was studied by using scanning supercritical angle fluorescence microscopy. The biological response to elucidated changes in lipid membrane structure/characteristics under ZnO or  $\text{TiO}_2$  NP influence was tested by fluorescence correlation spectroscopy. It was found the significant reduction of lipids diffusion mobility, which can be explained as a result of lipid-ZnO aggregates binding, depending on ZnO concentration. The  $\text{TiO}_2$  has a little effect on lipids diffusion mobility.

**Keywords:** ZnO Nanoparticles;  $\text{TiO}_2$  Nanoparticles; Lipid membrane; Supercritical angle fluorescence

Volume 2 Issue 3 - 2015

**Nora Grinceviciute<sup>1</sup>, Dorinel Verdes<sup>2</sup> and Valentinas Snitka<sup>1\*</sup>**<sup>1</sup>Department of Mathematics and Life Sciences, Kaunas University of Technology, Lithuania<sup>2</sup>University of Zurich, Switzerland

**\*Corresponding author:** Valentinas Snitka, Department of Mathematics and Life Sciences, Research Centre for Microsystems and Nanotechnology, Kaunas University of Technology, Studentu 65, LT-51369, Kaunas, Lithuania, Tel: +370 37 451588; Fax: +370 37 451588; Email: vsnitka@ktu.lt

Received: April 30, 2015 | Published: May 25, 2015

**Abbreviations:** NP: Nanoparticles; SLB: Supported Lipid Bilayers; BLMs: Bilayer Lipid Membranes; SAF: Supercritical Angle Fluorescence; FCS: Fluorescence Correlation Spectroscopy; PBS: Phosphate Buffered Saline; DLS: Dynamic Light-Scattering

## Introduction

Lipid membranes are essential components of any living cell and therefore, bilayer lipid membranes (BLMs) are simplified planar models of cell membranes, commonly employed for both fundamental and applied studies. BLMs as a model system are of great importance for improving our understanding of cell interaction with environment, various disease mechanism and membrane properties for drug screening purposes. Important area of nanomedicine research is the interaction of nanoparticles with cell and a possible toxicity [1-5]. Apart from their mechanical function as a cell wall, lipid membrane performs a function of a transfer mechanism for molecules diffusing through the membrane. As this process could distort the membrane structure, characterization tools that enable us to study the structure (ordering) of the membrane also provide information about the influence of NPs on the membranes. ZnO and  $\text{TiO}_2$  nanoparticles and their interaction with biological objects were investigated by several authors using different methods [6-10], however the mechanism of NP interaction with membrane is still an open question. Supercritical Angle Fluorescence Microscopy (SAF) is a technique that allows detection and characterisation of fluorescent species (proteins, biomolecules, pharmaceuticals,

etc.) and their behaviour very close or even adsorbed or linked at surfaces.

The method is able to observe molecules at a distance of up to 100 nanometers from the surface even in the presence of high concentrations of fluorescent species around [11-12]. In this work, we address two issues related to the interpretation of the fluorescence of phospholipid systems. While useful information can be retrieved from SAF microscopy, including the detection of very subtle effects, also the interpretation of the fluorescence correlation spectroscopy feature has not yet been settled, at least not in person NP's influence for lipid bilayer. Here, we specifically focus on the effect of different types and concentrations of NP's on lipid diffusion in supported lipid bilayers (SLBs). We used two different NPs: ZnO and  $\text{TiO}_2$  with 1 mg/ml and 2.5 mg/ml concentrations. The NP's are attached to the supported lipid bilayer by ionic interactions between positively charged NP's suspension and negatively charged lipid. The diffusion properties are determined by fluorescence correlation spectroscopy (FCS). FCS used to determine diffusion coefficient of SLB's alone and with NP's, to demonstrate the NP influence on lipid bilayer structure and molecular dynamics.

## Materials and Methods

All reagents and solvents used in this study were analytical grade. ZnO nano-powder ZincOx10 was obtained from Nanogate AG, Titanium dioxide nano powder from Na Bond, phosphate

buffered saline (PBS), Tris buffer, EDTA, NaCl, CaCl<sub>2</sub>, NaN<sub>3</sub> and 2-propanol from Sigma Aldrich. Lipids 1,2-dioleoyl-sn-glycero-3-phospho-L-serine (DOPS) and 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC) were obtained from Sigma Aldrich and 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE) from ATTO-TECH.

### Preparation of ZnO and TiO<sub>2</sub> suspension

Weighted 5 mg of ZnO/TiO<sub>2</sub> powder was sonicated for 15 min at 20°C in an ultrasonic bath along with 50 µl of 2-propanol. Next, 5 ml of PBS was added and the suspension (1mg/ml) was sonicated for 1 h at 20°C. For the higher concentration suspension preparation (2,5mg/ml) 12, 5 mg of ZnO/TiO<sub>2</sub> powder was sonicated for 15 min at 20°C in an ultrasonic bath along with 125 µl of 2-propanol and 5 ml of PBS was added and the suspension was sonicated for 1 h at 20°C.

### Substrate preparation

As support for the lipid bilayers, glass cover slips were used. The pieces (ca. 2.5 × 4 cm) were cleaned in ethanol and sonicated for 15 min at 20°C in an ultrasonic bath. Before use, these substrates were stored in ethanol. Before use, substrates were rinsed several times with Milli-Q water, dried with nitrogen gas and affected with low pressure plasma (Electronic Diener) for 10 minutes. For the bilayer preparation the substrates were glued onto an aluminium sample cell.

### Supported lipid bilayers

Supported lipid bilayers were prepared following a modified protocol of Richter et al. [13]. In short, (DOPS) and DOPC) and (DOPE) in chloroform were used as received and mixed at a ratio of 35:65:0, 000016. The lipid was dried overnight under vacuum to remove the solvent and then resuspended in Tris buffer (149 mMNaCl, 5 mM CaCl<sub>2</sub>, 10 mMTris) to a final concentration of 0.1 g/L. The resulting solution was extruded 27 times through a membrane (pore size 0.1 µm) to produce unilamellar vesicles. The vesicle solution was put directly in the measurement cell containing the glass substrate. There the solution was left for more than 1 h and after that it was replaced by the pure Tris buffer. Subsequently, this solution was exchanged again by a Ca<sup>2+</sup> free Tris buffer (10 mMTris, 133 mMNaCl) containing 5 mM EDTA in order to remove the Ca<sup>2+</sup> ions and finally by PBS. All buffers contained 0.05% NaN<sub>3</sub> and were adjusted to pH 7.4.

### SAF and FCS measurements

Images of fluorescently labelled lipid membranes were recorded with a custom-made scanning microscope allowing detection of fluorescence and the supercritical angle fluorescence (SAF) images of an emitter simultaneously. The SAF channel detects only fluorophores in close proximity to the surface (~100-200 nm) and efficiently rejects the fluorescence from the bulk solution using parabolic lenses as the decisive optical element. A detailed description of the optical setup is provided in the work [14]. All measurements were conducted by passing buffered solutions of NP's over the lipid membrane through the flow cell at a constant pump rate of 250 µl/min (0.42 mm/s). This flow rate is clearly slow enough to have no effect on the supported lipid bilayer integrity. Raw scan images are presented as measured using

appropriate linear scaling for the signal intensities. Note that the background subtraction in all images is based on a fixed intensity threshold. The shown images are selected representative images from time-lapse experiments which were conducted at varying time intervals, because the applied setup (non-commercial) does not contain the option of a software controlled time-lapse measurement.

Fluorescence Correlation Spectroscopy (FCS) is a method that was introduced and established as a very suitable approach to characterize membranes [14,15]. A single-molecule method with the dramatically improved sensitivity provides sufficient statistical significance to also use it for general characterization of membranes, mainly through the diffusion properties of their constituents (diffusion coefficients, concentrations, molecular brightness, intra molecular dynamics and molecular interactions) [16,17]. FCS was employed by several authors to measure the lateral lipid mobility and to assess the impact of NP's on lipid dynamics in supported lipid membranes [18-22]. A hardware correlator translates the photon arrival pulses into intensity fluctuations and calculates the correlation in real time. The correlation curves were acquired and fitted to analytical expressions. For an averaged correlation measurement, a separate correlation measurements of 100 seconds duration were taken and corresponding standard deviations for every point of the experimental curves were calculated from multiple experiments. The diffusion coefficient of the lipids in the supported lipid bilayer was determined from the experimentally obtained correlation time and the known waist of the confocal observation volume.

### Zeta potential measurements

All measurements were made at a temperature of 25°C on a DelsaNanoC (Beckman Coulter) fitted with a Flow Cell coupled to Auto Titrator. For all measurements, a voltage of 60 V was applied across an electrode spacing of 16 mm, the angle detection being of 15°. The solutions used for pH titration were 0.1M HCl and 0.1 M NaOH. The measurement for electrophoretic mobility was converted into zeta potential using Smoluchowski's formula [23-25].

### Atomic force microscopy measurements

The Atomic Force Microscopy (AFM) investigations for the ZnO and TiO<sub>2</sub> nanoparticles were done in the tapping mode (NT-MDT Inc.) using commercial silicon cantilevers NSG11 with a force constant of 5 Nm<sup>-1</sup>. The ZnO or TiO<sub>2</sub> water solution (1mg/1ml) was filtered by 200 nm polyethersulfone pore membrane (Chromafil PES-20/25, Macherey-Nagel) and placed onto the glass substrate (Carl Roth, 50x24mm, #1) and then dried and measured.

### Dynamic light scattering measurements

Size characterization of the samples was made by dynamic light-scattering (DLS) measurements using the same DelsaNanoC fitted with a Size Cell, which uses a 4 mW He-Ne laser operating at a wavelength of 658 nm, 30mW and a detection angle of 15°. Dispersions (1% Solution of TiO<sub>2</sub> or ZnO in water) were made at 3.24 mg/ml in Nano-pure water with 16 minutes of sonication on ice bath according with NRCWE protocol used for similar analysis. Size characterization of the samples was made by DLS

measurements using the same DelsaNanoC fitted with a Size glass cell, which uses a detection angle of  $160^\circ$ . Approximately 2 ml of the suspension was introduced to a glass size cell using a plastic pipette tip.


### Transmission electron microscopy measurements

The TEM analyses were performed using a JEOL JEM 2010F electron microscope operating at an accelerating voltage of 200 kV, with a point to point resolution of 0.19 nm (Joint Research Center, Ispra). The microscope was equipped with a Gatan electron energy loss imaging filter with an energy resolution equal to 0.7 eV and a multiscan CCD 1k x 1k CCD camera for image acquisition.

## Results and Discussion

### Characterization of nanoparticles

AFM and TEM were used to determine the size of NPs. The grain analysis function was used and then the distribution of NPs size was built. It gives the size of ZnO of 20.2 nm (standard deviation (SD) - 9.47 nm) and  $\text{TiO}_2$  gave the size value of 35.3 nm (SD-9.20 nm) (Figure 1). DLS was used by the instrument to determine the size distribution of particles by measuring dynamic fluctuations of light scattering intensity caused by the Brownian motion of the particle. This technique yields a hydrodynamic diameter that is calculated via the Stokes-Einstein equation from the aforementioned measurements. The measurement gives as result the average hydrodynamic diameter of the particles, the peak values in the hydrodynamic diameter ( $d_h$ ) distribution and the polydispersity index (PDI) that describes the width of the particle size distribution (Figure 2a-2f). Also, zeta potential measurements were performed. The magnitude of the zeta potential indicates the degree of electrostatic repulsion between adjacent, similarly charged particles in dispersion. ZnO zeta potential variation with pH 7-8.5, dispersions are stable and positively charged (with pH 7.4 ~ +3 mV). It is known, that low zeta potentials tend ZnO suspension to coagulate or flocculate. In  $\text{TiO}_2$  zeta potential variation with pH 5-9, dispersions are stable and surface is positive (with pH 7.4 ~ +33 mV). In this case, a high zeta potential will confer  $\text{TiO}_2$  stability, i.e., the suspension resists aggregation.




**Figure 1:** The experimental set-up for SAF imaging and FCS measurements.

### ZnO and $\text{TiO}_2$ NP's interaction with SLB

The SLB contained one part of negatively charged phospholipids (DOPS) and four parts of neutral phospholipids (DOPC) because the chosen ratio is similar to the composition of the inner

leaflet of neuronal membranes, which is typically enriched with negatively charged phospholipids in the range of 15-30 mol %. Generally, the interaction of nanoparticles with hydrophilic interfaces requires negative surface charges implying a strong contribution of electrostatic forces during the adsorption event. As shown in Figure 3, the fluorescence intensities of SLB after 16 h of exposure to a suspension of ZnO or  $\text{TiO}_2$  were recorded. The reduced fluorescence intensity of the membrane with ZnO (2.5 mg/ml) reflects on the possible ZnO aggregates deposition on SLB. The measured zeta potential of ZnO suspension shows the high probability of aggregates formation, which could quench the fluorescence intensity. The possible explanation why we obtained difference with different NP's is because the various (from 3 to 33 mV) zeta potentials lead to stability of suspension ( $\text{TiO}_2$  case) or big aggregates attraction to lipid bilayer (ZnO case).



**Figure 2:** AFM measurements and characterization, TEM images and selected area electron diffraction pattern and DLS Differential number vs. Diameter for 1% solution in water of ZnO (a, c, e) and  $\text{TiO}_2$  (b, d, f) nanoparticles, respectively.


To confirm NP's aggregates attachment to SLB surface, FCS measurements were used (Figure 4) indicating the impact of NP's on the lipids diffusion in the supported lipid bilayer at different concentrations. To investigate the ZnO and  $\text{TiO}_2$  NP's interaction and aggregation with supported lipid bilayers at different concentrations we employed SAF microscopy combined with the fluorescent correlation spectroscopy (FCS) as a sensitive and efficient method to study the interaction of nanoparticles

with supported lipid bilayers under continuous and constant flow conditions. For analysing the FCS data (Figure 4) we used a diffusion model expanded by a second diffusion coefficient in a linear combination ( $D, D_a, D_b$ )

$$G(t) = a * G_0 * (1 + 4D_a / \omega_0^2 t)^{-1} + b * G_0 * (1 + 4D_b / \omega_0^2 t)^{-1}$$

$$\text{with } D = a * D_a + b * D_b$$


Where  $G_0$  is the intercept of the FCS curve and inverse proportional to the number of the fluorescent molecules and  $\omega_0$  is the radius of the detection volume. Both NP's show slight variations ( $D$  is in range  $2\text{-}3\text{E-}12 \text{ m}^2 \text{ s}^{-1}$ ) in the diffusion coefficient values, with no defined trend. Although ZnO with 2.5 mg/ml concentration has a different impact on lipid diffusion ( $D\text{-}4.91\text{E-}13 \text{ m}^2 \text{ s}^{-1}$ ). The effect could be due to the unstable suspension caused aggregations connections to SLB surface. The influence of addition of the ZnO or  $\text{TiO}_2$  NP's to the lipids suspension and NP's effect on the supported lipid membranes was studied by analysing the FCS measurements of the labelled DOPE (diffusion coefficient- $D$ ) lipids contained inside membranes before and after nanoparticles addition. Figure 4 illustrates three independent measurements of the diffusion coefficient ( $D$ ) without addition of nanoparticles and after addition 1 mg/ml and 2.5 mg/ml ZnO or  $\text{TiO}_2$  concentration, respectively. FCS measurements reveal that the significant reduction of  $D$  occurs only after addition of 2.5 mg/ml ZnO; other samples remained quite stable for a longer period of time.



**Figure 3:** ZnO and  $\text{TiO}_2$  interaction with lipid membrane. SAF images of the scanned areas acquired after 16h of exposure to a suspension of ZnO (top row) or  $\text{TiO}_2$  (bottom row) with different concentrations on the lipid membrane bilayer.

The significant reduction of  $D$  can be explained as a result of lipid-ZnO aggregates binding. This effect is enlarged with the increasing ZnO concentration. The higher ZnO concentration leads to the enhancement of the  $D$  reduction. This is in line with other studies, which reported that a noticeable influence over lipid membranes is observed at higher concentrations of cholesterol, proteins or other nanoparticles [26-32]. This effect is explained by reduction of free area because of tighter lipid packing. To determine if the decrease of the lipids diffusion coefficient is followed by a destruction of the bilayer homogeneity and

integrity, scanned SAF images were performed at different times (Figure 3). The scanned images reveal that the bilayer integrity is not altered for 16 hours after nanoparticles addition. Thus, the higher concentration of ZnO fluorescence intensity is significantly lower from others presented in Figure 3. This let us conclude that the ZnO addition induces a significant alteration of the mobility of lipids inside the membrane before any aggregation processes of nanoparticles start to occur. The possible explanation is that the ZnO zeta potential (with pH 7.4  $\sim$  +3 mV) could affect the lipids mobility due to the unstable suspension caused aggregations connections to SLB surface. In opposite,  $\text{TiO}_2$  zeta potential variation (with pH 7.4  $\sim$  33 mV) shows stable suspension. Experimental results suggest that the nanoparticle-membrane interaction depends on the NP charge as well as the zeta potential. Cationic nanoparticles are known to penetrate through the cell membrane [1]. It seems evident that the binding of  $\text{TiO}_2$  with the membrane takes place through relatively weak contacts since NP did not immerse deeply in the membrane but floated on top of the bilayer surface, enabling diffusion of the membrane, with no adsorption to the membrane. This is in line with other study, where reported that a noticeable and statistically significant ( $p < 0,001$ ) influence on phospholipid vesicles were observed after incubation of ZnO (but not  $\text{TiO}_2$ ) nanoparticles [33].



**Figure 4:** Fluorescence intensities of lipid membrane without (blue line) and with ZnO (a; red and green lines) or with  $\text{TiO}_2$  (b; red and green lines). Diffusion curves (from FCS measurements) indicating the difference in the presence (blue) and absence (green) of 2.5 mg/ml ZnO. The solid black lines represent the fitting curves for corresponding FCS curves.

This is known that agglomerates larger than  $0,2 \mu\text{m}$  were found attached to the membrane [34], whereas smaller aggregates and single particles could probably cross the membrane through pores or ion channels [35]. In addition, aggregates adhere to surfaces may therefore cause effects without even entering cells. The effects of NPs on SLB are important for understanding the potential effects of NPs on biological membranes. Limited biological relevance of the results of experiments with SLB does not mean that these results have no value for understanding effects on biological membranes. It is indicated that ZnO (but not  $\text{TiO}_2$ ) can cause the changes of the membrane characteristics if it can get close to the phospholipid bilayer.

## Conclusion

In this study, we investigated the impact of ZnO and  $\text{TiO}_2$ , being essential constituents of drugs and cosmetics, on the dynamics of lipid molecules in bilayer membranes. Using electrostatically

attached NP's of positive charge on negatively charged supported model membranes, we demonstrated that the diffusion mobility of lipids within the ZnO membrane slowed down significantly, depending on concentration. The weak interaction was found between a TiO<sub>2</sub> and lipid bilayers. As an explanation we suggest a mechanism of strong interaction between positively charged nanoparticles and the negatively charged lipids, thus forming tight nanoparticles-lipid complexes in the case of ZnO. The finding that diffusion of lipid molecules within the membrane significantly depends on interaction with NP represents a new step in understanding the interactions between NP and cells by demonstrating the non disruptive nature of the NP and lipid bilayer interaction and showing the importance of NP surface properties.

### Acknowledgement

This research is funded by the European Social Fund under the Global Grant measure (Grant No. VP1-3.1-SMM-07-K-03-044).

### References

- Heikkilä E, Martinez-Seara H, Gurtovenko AA, Vattulainen I, Akola J (2014) Atomistic simulations of anionic Au 144 (SR) 60 nanoparticles interacting with asymmetric model lipid membranes. *Biochim Biophys Acta* 1838(11): 2852-2860.
- He XC, Lin M, Li F, Sha BY, Xu F, et al. (2015) Advances in studies of nanoparticle-biomembrane interactions. *Nanomedicine* 10(1): 121-141.
- Karlsson HL, Cronholm P, Hedberg Y, Tornberg M, De Battice L, et al. (2013) Cell membrane damage and protein interaction induced by copper containing nanoparticles-Importance of the metal release process. *Toxicology* 313(1): 59-69.
- Adjei IM, Sharma B, Labhassetwar V (2014) Nanoparticles: cellular uptake and cytotoxicity. In *Nanomaterial* 811: 73-91.
- Lu B, Smith T, Li R, Xia T, Nel A, et al. (2014) Nanoparticle-Membrane Interactions Studied with Lipid Bilayer Arrays. *Biophys J* 106(2): 415a.
- Sanjay SS, Pandey A, Ankit P, Chattopadhyaya M (2013) Fabrication of surfactant sensing membrane with znO nano-composite. *Proceedings of the National Academy of Sciences, India Section A: Physical Sciences* 83(3): 279-285.
- Sundaram Sanjay S, Yadav RS, Pandey AC (2013) Synthesis of lamellar porous photocatalyticnanozno with the help of anionic surfactant. *Adv Mater Lett* 4(5).
- Asl SK, Sadrmezhaad S (2010) The seeding effect on the microstructure and photocatalytic properties of znO nano powders. *Materials Letters* 64(18): 1935-1938.
- Maurya A, Chauhan P, Mishra A and Pandey AK (2012) Surface functionalization of TiO<sub>2</sub> with plant extracts and their combined antimicrobial activities against *E. faecalis* and *E. Coli*. *J Res Updates Polymer Sci* 1(1): 43-51.
- Šimundić M, Drašler B, Šuštar V, Zupanc J, Štukelj R, et al. (2013) Effect of engineered TiO<sub>2</sub> and ZnO nanoparticles on erythrocytes, platelet-rich plasma and giant unilamellar phospholipid vesicles. *BMC vet Res* 9(1): 7.
- Ruckstuhl T, Verdes D (2004) Supercritical angle fluorescence (SAF) Microscopy. *Opt Express* 12(18): 4246-4254.
- Verdes D, Ruckstuhl T, Seeger S (2007) Parallel two-channel near- and far-field fluorescence microscopy. *J Biomed Opt* 12(3): 034012.
- Richter R, Mukhopadhyay A, Brisson A (2003) Pathways of lipid vesicle deposition on solid surfaces: a combined QCM-D and AFM study. *Biophys J* 85(5): 3035-3047.
- Schwille P, Korlach J, Webb WW (1999) Fluorescence correlation spectroscopy with single-molecule sensitivity on cell and model membranes. *Cytometry* 36(3): 176-182.
- Bacia K, Scherfeld D, Kahya N, Schwille P (2004) Fluorescence correlation spectroscopy relates rafts in model and native membranes. *Biophys J* 87(2): 1034-1043.
- Eggeling C, Ringemann C, Medda R, Schwarzmann G, Sandhoff K, et al. (2009) Direct observation of the nanoscale dynamics of membrane lipids in a living cell. *Nature* 457: 1159-1121.
- Sezgin E, Schwille P (2011) Fluorescence techniques to study lipid dynamics, *Cold Spring Harb perspect Biology* 3(11): a009803.
- Haustein E, Schwille P (2007) Fluorescence correlation spectroscopy: novel variations of an established technique. *Annu Rev Biophys Biomol Struct* 36: 151-169.
- Chiantia S, Ries J, Chwastek G, Carrer D, Li Z, et al. (2008) Role of ceramide in membrane protein organization investigated by combined AFM and FCS. *Biochim Biophys Acta*. 1778(5): 1356-1364.
- Lingwood D, Ries J, Schwille P, Simons K (2008) Plasma membranes are poised for activation of raft phase coalescence at physiological temperature. *Proc Natl Acad Sci* 105(29): 10005-10010.
- García-Sáez AJ, Schwille P (2010) Stability of lipid domains. *FEBS Lett* 584(9): 1653-1658.
- García-Sáez AJ, Carrer DC, Schwille P (2010) Fluorescence correlation spectroscopy for the study of membrane dynamics and organization in giant unilamellar vesicles. *Methods Mol Biol* 606: 493-508.
- Smoluchowski M (1921) *Handbuch der Electricität und des Magnetismus (Graetz) (vol. II)*, Barth, Leipzig, Germany, pp. 366.
- Delgado AV, González-Caballero F, Hunter RJ, Koopal LK, Lyklema J (2005) Measurement and interpretation of electrokinetic phenomena. *Pure Appl Chem* 77(10): 1753-1805.
- Kaszuba M, Corbett J, Watson FM, Jones A (2010) High-concentration zeta potential measurements using light-scattering techniques. *Philos Trans A Math Phys Eng Sci* 368(1927): 4439-4451.
- Kahya N, Scherfeld D, Bacia K, Schwille P (2004) Lipid domain formation and dynamics in giant unilamellar vesicles explored by fluorescence correlation spectroscopy. *J Struct Biol* 147(1): 77-89.
- Kahya N, Schwille P (2006) How phospholipid-cholesterol interactions modulate lipid lateral diffusion, as revealed by fluorescence correlation spectroscopy. *J Fluoresc* 16(5): 671-678.
- Sahoo H, Schwille P (2013) Influence of glycosaminoglycans on lipid dynamics in supported phospholipid bilayers. *Soft Matter* 9(14): 3859-3865.
- Bockmann RA, Hac A, Heimburg T, Grubmuller H (2003) Effect of sodium chloride on a lipid bilayer. *Biophys J* 85(3): 1647-1655.
- Sum AK, Faller R, de Pablo JJ (2003) Molecular simulation study of phospholipid bilayers and insights of the interactions with disaccharides. *Biophys J* 85(5): 2830-2844.
- Blondelle SE, Lohner K, Aguilar MI (1999) Lipid-induced

- conformation and lipid-binding properties of cytolytic and antimicrobial peptides: determination and biological specificity. *Biochim Biophys Acta* 1462 (12): 89-108.
32. Macháň R, Hof M (2010) Lipid diffusion in planar membranes investigated by fluorescence correlation spectroscopy. *Biochim Biophys Acta Biomembr* 1798(7): 1377-1391.
  33. Šimundić M, Drašler B, Šuštar V, Zupanc J, Štukelj R, et al. (2013) Effect of engineered TiO<sub>2</sub> and ZnO nanoparticles on erythrocytes, platelet-rich plasma and giant unilamellar phospholipid vesicles. *BMC vet Res* 9: 7.
  34. Rothen-Rutishauser BM, Schurch S, Haenni B, Kapp N, Gehr P (2006) Interaction of fine particles and nanoparticles with red blood cells visualized with advanced microscopic techniques. *Environ Sci Technol*, 40(14): 4353-4359.
  35. Porter AE, Muller K, Skepper J, Midgley P, Welland M (2006) Uptake of C60 by human monocyte macrophages, its localization and implication for toxicity: studied by high resolution electron microscopy and electron tomography. *Acta Biomater* 2(4): 409-419.