

Kangaroo Mother Care for Low Birth Weight Babies: A Randomized Controlled Trial in a Tertiary Care Hospital of Bangladesh

Research Article

Volume 7 Issue 2 - 2017

Maksudur Rahman*, **MAK Azad Chowdhury**,
Md Mahbubul Hoque, **Nishat Jahan** and
Liton Chandra Shaha*Department of Neonatology, Bangladesh Institute of Child health(BICH), Dhaka Shishu (Children) Hospital, Bangladesh*

*Corresponding author: Maksudur Rahman, Associate professor, Department of neonatology, Bangladesh Institute of Child health(BICH), Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh, Email: maksudr@gmail.com

Received: September 22, 2017 | **Published:** October 11, 2017**Abstract**

Background: KMC is a safe, low cost and effective method of caring for low birth-weight (LBW) babies to reduce neonatal morbidity and mortality.

Objectives: To assess the effect of KMC on LBW babies when compared with conventional method of care.

Methods: This randomized controlled trial study was conducted in neonatal ward of Dhaka Shishu (Children) Hospital, Dhaka, from November, 2014 to October 2015. All preterm LBW neonates with birth weight 1250gm to 1800 gm., gestational age > 30 weeks to < 35 weeks, haemodynamically stable were given KMC and compared with conventional method of care. All the babies were monitored and data regarding weight, temperature, apnea, nosocomial infection, feeding, hospital stay and mortality etc. were recorded during the hospital stay and followed up to 40 weeks of gestational age. Statistic analysis was done by χ^2 , student t test etc. with SPSS-17 version.

Results: comparison of 40KMC babies with 40 control showed that rate of weight gain per day in KMC group was 18.11 ± 7.67 gms and in control group, it was 13.02 ± 4.45 gms ($p < 0.001$). Apnea occurred less in KMC group (8% vs. 15%) ($p=0.228$) and episodes of hypothermia was also recorded less in KMC group (10% vs 18%) ($p=0.210$). Culture positive sepsis was 15% in KMC group and 20% in control group ($p=0.556$) The mean time to achieve full enteral feed in KMC group was 9.08 ± 2.35 days and 14.735 ± 4.54 days ($p < 0.001$) in control group Ninety percent of KMC and 60% of control group babies were discharged with exclusive breast feeding ($p=0.002$). Mean hospital stay was 15.61 ± 10.56 days vs 18.15 ± 4.5 days, in KMC and control group respectively that was statistically insignificant, $p=0.15$. The mortality was 2(5%) in KMC group and those was 6(15%) in control group and that was also statistically insignificant ($P=0.136$).

Conclusion: In this study, KMC is effective for caring of LBW babies and has better effect on weight gaining and establishment of exclusive breast feeding. So KMC is recommended for caring of low birth weight infants.

Keywords: Kangaroo mother care; KMC; Dhaka Shishu Hospital

Abbreviations: KMC: Kangaroo Mother Care

Introduction

Bangladesh has one of the top 10 highest burdens of newborn deaths in the world [1]. Nearly 45% of all newborn deaths are directly related to preterm birth and its complications [2]. In conventional methods of care, Preterm and LBW (Low birth weight) babies need incubators which are costly, highly technical and need skilled personnel and proper maintenance [3]. Moreover incubators separate babies from their mother, depriving them of the necessary contact. So good care of LBW babies is difficult, and hypothermia and nosocomial infections are frequent in low resource developing countries. Under such circumstances KMC (Kangaroo Mother Care) is a good alternative and some instance more effective than incubator in caring of preterm and LBW babies. To decrease neonatal morbidity and mortality, KMC is now considered as the most feasible, readily available, and preferred

intervention in developed and developing countries [4]. It is a special way of caring for the preterm or low birth weight infants where baby is kept in skin-to-skin contact with the mother and breastfed exclusively [5]. It is a safe and effective method of caring for low birth weight (LBW) infants, especially in low-resourced settings [4]. It has also been described as a cost-effective, [6,7] high-impact intervention [8,9] for improving newborn survival. Beneficial physiological and behavioral effects of KMC for the infant has been well documented. Physiological effects include better thermoregulation, improved cardiorespiratory stability, lower risk of infection and faster growth. Behavioral effects relate to better sleep cycles, less crying and an analgesic effect during painful procedures. Beneficial effects for the mothers include better breastfeeding (increased milk production, exclusivity, duration, early initiation) and psychosocial effects (reduced anxiety, more maternal satisfaction, improved maternal-infant attachment and bonding) [10,11]. This study was conducted to

assess the effect of KMC on LBW babies when compared with conventional method of care.

Materials and Methods

This randomized controlled trial study was conducted in neonatal ward of Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh from November, 2014 to October 2015. All preterm, neonates with birth weight 1250gm to 1800gm., gestational age > 30 weeks to < 35 weeks (Gestational age determined by New Ballard Score), haemodynamically stable (criteria of stable baby was 1) normal heart rate 100-160 per minutes, 2) respiratory rate 30-59 per minute with breaths comfortably, no sign of respiratory distress pink in room air or with 40% oxygen, no prolonged or frequent apnea) were included in this study. Babies having major life threatening congenital malformation, perinatal asphyxia and required ventilator or ion tropic support and babies with critically ill mother were excluded. Neonates in this study were randomly enrolled by inclusions and exclusions criteria, and were allocated either KMC group or control group by simple randomization with concealed sealed envelope technique. Forty neonates (40) were in KMC group and forty neonates (40) were in control group. Neonates in control group were managed under radiant warmer. A written informed consent was obtained from all mothers. Details of antenatal period and delivery were recorded. During hospital stay, both the groups were monitored for daily weight gain, episodes of hypothermia, apnea, nosocomial infection, physiological parameters (heart rate, respiratory rate, axillary temperature and oxygen saturation). Data regarding feeding was also recorded. After discharge, the neonates were followed weekly till 40 weeks of corrected gestational age. All information was recorded according to follow up sheet. In KMC group mother or caregiver was explained detail about KMC adoption in presence of their family. KMC was initiated as soon as the baby was stable. If the mother was not available, KMC was started with the help of any other family member. In KMC position, baby was kept skin to skin contact with mother bare chest in between the breasts. Baby was remained froglike position, head turned in one side with neck slight extended, wearing front open sleeveless shirt, cap, shocks and a nappy. There was a binder for support the baby around baby and mother. Comfortable bed or chair was provided to the mother or caregiver practicing KMC in ward. KMC was given at least 2 hours at a time and at least 12 hours in a day. When the baby was not in KMC, the baby placed in cot with adequate cloths and coverings. Infant in both the groups were discharged when following criteria were achieved: Baby's general health was good, no evidence of infection and no I/V medications. Baby was feeding well and receiving breast milk directly or by cup and spoon gaining weight (at least 15gm/kg/day) for 3 consecutive days. Maintaining body temperature well without assistance for at least 3 consecutive days and Mother and family members were confident to care the baby. Besides KMC and radiant warmer care, babies of both groups were treated with conventional hospital protocol. Data were analyzed by chi-square (χ^2) test, unpaired student's 't' test with the help of SPSS version-17.

Results

In this study, comparison of 40KMC babies with 40 control showed that rate of weight gain per day in KMC group was 18.11 ±

7.67gms and in control group, it was 13.02 ± 4.45gms (p< 0.001). Apnea occurred less in KMC group (8% vs. 15%) (p=0.228) and episodes of hypothermia was also recorded less in KMC group (10% vs 18%) (p=0.210). But both are statistically insignificant. Culture positive sepsis was 15% in KMC group and 20% in control group (p=0.556). The mean time to achieve full enteral feed in KMC group was 9.08 ± 2.35 days and 14.735 ± 4.54 days (p < 0.001) in control group. Ninety percent of KMC and 60% of control group babies were discharged with exclusive breast feeding (p=0.002). The rest received cup feeding along with the breast feeding. In this study mean hospital stay was 15.61 ± 10.56 days vs 18.15 ± 4.5 days, in KMC and control group respectively that was statistically insignificant, p=0.15. The mortality was 2(5%) in KMC group and those was 6(15%) in control group and that was also statistically insignificant (P=0.136) (Tables 1-4).

Table 1: Distribution of the patients according to particulars of the patients of comparison groups.

Variables	Case(n=40)	Control (n=40)	P value
Age at admission (day) (Mean ± SD)	1.80 ± 1.09	2.10 ± 1.19	0.244
Sex of the baby	No. (%)	No. (%)	0.348
Male	24(60%)	28(70%)	
Female	16(40%)	12(30%)	

Table 2: Feeding outcome between KMC and conventional care group.

Hospital Course	Case (n=40) No. (%)	Control (n=40) No. (%)	p value
Start of first feed on			0.175ns
1st day	13(33.0%)	8(20.0%)	
2-3 days	20(50.0%)	19(48.0%)	
4-5 days	7(17.0%)	13(32.0%)	
Time of achieve full enteral feeding in days	9.08 ± 2.35	14.735 ± 4.54	<0.001*
Exclusive breast feeding			<0.001*
Yes	36(90.0%)	24(60.0%)	
No	4(10.0%)	16(40.0%)	

Statistical analysis was done by Chi-square test and unpaired' test.

P <0.05 was taken as statistically significant,

n =Total number of subjects

Table 3: Showing rate of weight gain between KMC and conventional care group.

Variables	Case (n=40) Mean ± SD	Control (n=40) Mean ± SD	p value
Weight gain started(days)	5.87 ± 1.80	8.57 ± 2.28	<0.001*

Birth weight regained (days)	10.79 ± 2.87	13.16 ± 3.74	<0.001*
Rate of weight gain (gms)	18.11±7.67	13.02 ± 4.45	<0.001*

Statistical analysis was done by unpaired' test.

P <0.05 was taken as statistically significant, n =Total number of subjects

Table 4: Distribution of the patients according to hospital course of the patients.

Hospital course	Case (n=40) No. (%)	Control (n=40) No. (%)	p value
Episode of hypothermia Yes No	4(10%) 36(90%)	7(18%) 33(82%)	0.210
Episode of apnea Yes No	3(7.5%) 37(92.5%)	6(15%) 34(85%)	0.228
Culture proven sepsis Yes No	6(15.0%) 34(85.0%)	8(20.0%) 32(80.0%)	0.556
Exclusive breast feeding Yes No	36(90.0%) 4(10%)	24(60.0%) 16(40.0%)	0.002*
Mortality yes No	2(5%) 38(95%)	6(15%) 34(85%)	0.136
Mean hospital stay (days: mean ± SD)	15.61 ± 10.56	18.15 ± 4.5	0.154

Statistical analysis was done by Chi-square test and unpaired' test.

P <0.05 was taken as statistically significant, n =Total number of subjects

Discussion

This study was conducted at Dhaka shishu (children) hospital from November, 2014 to October, 2015. In this period, 40KMC babies were compared with same number of babies who did not receive KMC. Although KMC is a key intervention package in newborn health initiatives, there is limited systematic information available on the barriers to KMC practice that mothers and other stakeholders face while practicing KMC [12]. A systematic review showed five top most barriers to KMC practice. This were 1) issues with facility environment/resources 2) negative impressions of staff attitudes or interactions 3) fear/anxiety of hurting the infant 4) lack of help with KMC practice and other obligations and 5) low awareness of KMC/infant health. In this study though it was not analyzed statistically few barriers were observed regarding practicing of KMC. The main barrier was absent of appropriate helper of mother that means any person, nurse or family member who helps mother during KMC practice, during her discomfort and bathing etc. Without a continuous presence of a helper KMC cannot be continued. Unwillingness of practicing KMC,

that was overcome with repeated counseling, absence of proper monitoring of KMC practice at home, absence of appropriate privacy of mother, sweating of mother specially during summer season were important barriers of KMC practice in this study.

As with any other intervention KMC has certain costs too. For implementation of KMC in existing neonatology ward, it was needed bed, chair with back support, baby's clothes, cap, socks, nappy, mother's gown, binder which cost about 122.25US\$/baby and incubator cost about 5000US\$/baby plus maintenance cost. The cost of 2 days training program for health personnel about KMC initially, was about 375US\$. The less hospital stay also lessened the cost of KMC. Besides above this, So KMC is a cheap and cost effective intervention.

A global health recommendation is that infants should be exclusively breast fed for the first six months of life as because breast milk mediates unequalled beneficial effects regarding nutritional, immunological, and cognitive outcomes [13]. Managing these preterm babies are clinically more challenging in initiation and sustainability of breastfeeding as their suckling behavior is not mature and requires neonatal care for a sustainable period. In Bangladesh exclusive breastfeeding rate has declined from 62 to 54 and there is no data on preterm breastfeeding rate. As preterm rate is high in Bangladesh it might have influence on declining exclusive breastfeeding rate [14]. A study showed that among Canadian babies weighing less than 2500gm only 58% had been breastfed at birth, compared to 73% in term group. Among LBW infants only 3% were discharged on exclusive breastfeeding [15]. Study done in Malaysia showed, only 40.2% of VLBW babies were being breastfed at the time of discharge from hospital despite breastfeeding incentive programs adopted by the hospital [16]. In this study we have found that early feeding could have been started in LBW and VLBW babies who were managed in KMC and also the time of achieving full enteral feeding was earlier and rate of exclusive breastfeeding (90% vs 60%) was found higher in these babies. A study from Sweden done by Whitelaw et al found that 71 preterm babies weighing less than 1500 g submitted to KMC had a two time higher prevalence of breast feeding than the control group at six weeks of life (55% versus 28%) [17]. Kangaroo Mother Care (KMC), which has an extensive research base, is regarded as a successful way to empower mothers to become familiar with their infants, strengthens their mothering at their own pace, and increases breastfeeding rate and duration [18,19].

As it is known that neonate usually loses weight in his first week of life than starts gaining weight, similar trend was seen in this study. But in this study babies with KMC had early gaining weight and early regaining weight than babies with conventional care (p<001). Moreover weight gain was significantly higher in KMC babies than nonKMC babies (p<001). Similarly in another study, it was found that KMC babies achieved significantly better growth (KMC; 23.99gms vs Conventional Care; 15.58gms, p<0.0001) [20]. Meta analysis of KMC studies confirmed greater wt gain in KMC infants than in infants not managed in KMC [21].

In this study apnea occurred in 8% babies in KMC group and 15% cases in control group which was statistically insignificant (p=0.228). In the study of Chowdhury et al. apnea in KMC group was found less (KMC: 2, 8% vs Control: 8, 32%, p= 0.03) [22].

The episodes of hypothermia was 10% in KMC group and 18% in control group that was also statistically insignificant ($p=0.210$) in this study. Several studies showed that incidence of hypothermia were less in KMC group. Cattaneo et al. showed in their study that hypothermia was more in babies that did not get KMC (31.5 vs 13.5 episode/100 infants) [23].

There were no statistical differences between two groups regarding culture positive sepsis. It was found 15% cases in KMC group and 20% babies in control group ($p=0.556$). But Chowdhury et al. [22] showed culture positive sepsis was more in babies who did not get KMC than who got KMC [22].

In this study mean hospital stay was 15.61 ± 10.56 days vs 18.15 ± 4.5 days in KMC and control group respectively that was statistically insignificant ($p=0.15$). There was no difference in hospital stay between KMC and no-KMC group in the study done by Rao et al [20]. (KMC vs no-KMC: 12.78 days vs 12.86 days, $P=0.929$). But Chowdhury et al. [22] found less hospital stay in KMC group in their study and which was statistically significant.

Mortality was 2(5%) in KMC group and those was 6(15%) in control group and that was statistically insignificant, $P=0.136$. But meta-analysis of three RCTs showed a significant reduction in neonatal mortality [relative risk (RR) 0.49, 95% confidence interval (CI) 0.29-0.82] compared with standard care [23].

Conclusion

In this study, KMC is effective for caring of LBW babies and has better effect on weight gaining and establishment of exclusive breast feeding. So KMC is recommended for caring of low birth weight infants.

Acknowledgements

Professor (Dr.) MAK Azad Chowdhury,

Member of Technical committee of KMC, Bangladesh

Head of KMC Project, Dhaka Shishu (Children) Hospital, Dhaka.

Prof. and head, Department of neonatology & Academic director

Bangladesh Institute of Child Health (BICH), Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh.

Conflict of Interest

None.

References

1. Lawn JE, Cousens S, Zupan J (2005) 4 million neonatal deaths: When? Where? Why? *Lancet* 365(9462): 891-900.
2. Ahmed FA (2012) UNICEF Saving preterm babies in Bangladesh hospitals.
3. Mohammadzadeh A, Farhat A, Jafarzadeh M, Hasanzadeh L, Esmali H (2011) Advantages of kangaroo mother care in less than 2000 grams low birth weight neonates. *M JIRI* 25(1): 11-15.
4. Bergh AM, Manu R, Davy K, Van RE, Quansah AG, et al. (2013) Progress with the implementation of kangaroo mother care in four regions in Ghana. *Ghana Med J* 47(2): 57-63 .
5. World health organization (2003) Kangaroo Mother Care: a practical guide. Department of reproductive Health and research, WHO, Geneva, Switzerland, p. 1-54.
6. Darmstadt GL, Bhutta ZA, Cousens S, Adam T, Walker N, et al. (2005) Evidence-based, cost effective interventions: how many newborn babies can we save? *Lancet* 365(9463): 977-988.
7. Victora CG, Rubens CE, GAPPS Review Group (2010) Global report on preterm birth and stillbirth (4 of 7): delivery of interventions. *BMC Pregnancy Childbirth* 10(S1): S4.
8. Lawn JE, Kerber K, Enweronu LC, Cousens S (2010) 3.6 Million neonatal deaths-what is progressing and what is not? *Semin Perinatol* 34(6): 371-386.
9. Barros FC, Bhutta ZA, Batra M, Hansen TN, Victora CG, et al. (2010) Global report on preterm birth and stillbirth (3 of 7): evidence for effectiveness of interventions. *BMC Pregnancy and Childbirth* 10(S1): S3.
10. Ludington HSM, Morgan K, Abouelfetoh A (2008) A clinical guideline for implementation of kangaroo care with premature infants of 30 or more weeks' postmenstrual age. *Adv Neonatal Care* 8(S3): S3-S23.
11. Ruiz JG, Charpak N (2007) Evidence-based clinical guidelines for an optimal use of the kangaroo mother method in preterm and/or low birth weight infants at birth. Bogotá, Colombia: Fundación Canguro and Department of Clinical Epidemiology and Biostatistics, School of Medicine, Pontificia Universidad Javeriana.
12. Seidman G, Unnikrishnan S, Kenny E, Myslinski S, Cairns-Smith S, et al. (2015) Barriers and Enablers of Kangaroo Mother Care Practice: A Systematic Review. *PLoS ONE* 10(5): e0125643.
13. Kramer MS, Kakuma R (2002) The optimal duration of exclusive breastfeeding. A systemic review. *Adv Exp Med Biol* 554: 63-77.
14. Bangladesh Demographic Health Survey (BDHS) (2014) National Institute of Population Research and Training Ministry of Health and Family Welfare Dhaka, Bangladesh, pp. 1-354.
15. Blencowe H, Cousens S, Oestergaard MZ, Csou D, Moller AB, et al. (2012) National, regional and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *Lancet* 379(9832): 2161-2172.
16. Lefebvre F, Ducarme M (1989) Incidence and duration of lactational performance among mother of low-birth-weight infants. *CMAJ* 140(10): 1159-1164.
17. Boo NY, Goh ES (1999) Predictors of breast feeding in very low birth weight infants at the time of discharge from hospital. *J Trop Pediatr* 45(4): 195-201.
18. Feldman R, Eidelman AI, Sirota L, Waller A (2002) Comparison of skin-to-skin (Kangaroo) and traditional care: Parenting outcome and preterm infant development. *Pediatrics* 110(1-1): 16-26.
19. Hake-Brooks SJ, Anderson GC (2008) Kangaroo care and breastfeeding of mother-preterm infant dyads 0-18 months: A randomized control trail. *Neonatal Netw* 27(3): 151-159.

20. Rao PN, Udani R, Nanavati R (2008) Kangaroo Mother Care for Low birth weight infants: a randomized controlled trial. *Indian pediatr* 45(1): 17-23.
21. Conde-Agudelo A, Diaz Rosselo JL, Belizan JM (2004) Kangaroo Mother Care to reduce mortality and morbidity in low birth weight infants. *Cochrane Database Syst Rev* 22(4): CD002771.
22. Chowdhury RM (2013) Kangaroo Mother Care as compared to standard care for management of preterm low birth weight babies. MD Thesis, BSMMU, Dhaka.
23. Rekha HU, Nandkishor K, Ruchi NN, Alope VR (2013) Kangaroo mother care: A cohort follow up study on impact of duration of KMC on mortality, morbidity, hospital stay and breastfeeding. *Journal of Neonatology* 27(1): 5-10.