

Determination of sexual dimorphism from the area of the triangle

Abstract

A triangular area can be formed from lines connecting the infraorbital foramina (IOFs) and the nasion to each other. Measurements on this triangle may help to determine sexual dimorphism in human skulls. The aim of this study was to determine sexual dimorphism in Brazilian skulls from measurements of the area of the triangle formed between lines joining the IOFs to the nasion. 242 human skulls belonging to the Study and Research Center for Anatomy and Forensic Anthropology of Tiradentes University. It was observed that out of the 242 skulls examined, 148 were male and 94 were female, with ages ranging from 18 to 91 years (mean: 57 years). In the male skulls, the distance between the IOFs ranged from 53 to 58.8 mm (mean: 56 mm). In the female skulls, it ranged from 50.7 to 56.0 mm (mean: 53.7 mm). In the male skulls, the mean distance from the right IOF (RIOF) to the nasion ranged from 44.1 to 49.0 mm (mean: 46.2 mm) and from the left IOF (LIOF) to the nasion, 44 to 48 mm (mean: 46.0 mm). In the female skulls, the distance from the RIOF to the nasion ranged from 42 to 47 mm (mean: 44.7 mm) and from the LIOF to the nasion, 41.8 to 46.1 mm (mean: 44 mm). The area of the triangle formed by the lines connecting the IOFs to the nasion in the male skulls ranged from 944.6 to 1105.3 mm² (mean: 1023.9 mm²). In the female skulls, it ranged from 853.8 to 1025.3 mm² (mean: 946.5 mm²). A sexual dimorphism occurred in all measures and area that formed the triangle between the infraorbital foramen and the nasion that were larger in males.

Keywords: anthropometry, anatomy, sexual dimorphism, infraorbital foramen, nasion

Volume 3 Issue 3 - 2017

José Aderval Aragão,¹ Maria Paula Mendonça Silveira,² Laís Carolina Santos Cisneiros de Oliveira,² Felipe Matheus Sant'Anna Aragão,³ Iapunira Catarina Sant'Anna Aragão,³ Francisco Prado Reis⁴

¹Department of Morphology and Postgraduate Applied Health Science Program, Federal University of Sergipe (UFS), Brazil

²Dentistry student, Federal University of Sergipe (UFS), Brazil

³Medical Student, University Center of Volta Redonda (UNIFOA), Brazil

⁴Titular Professor, Medical School, Tiradentes University (UNIT), Brazil

Correspondence: José Aderval Aragão, Associate Professor of the Federal University of Sergipe and Titular Professor of the Medical School of the Tiradentes University, Rua Aloisio Campos 500, Bairro Atalaia, Aracaju, Sergipe, Brazil, CEP 49035020, Tel +5579991916767, Email jaafelipe@infonet.com.br

Received: November 01, 2016 | **Published:** March 17, 2017

Abbreviations: N, nasion; RIOF, right infraorbital foramen; LIOF, left infraorbital foramen; DRIOF, distance from the right infraorbital foramen to the nasion; DLIOFN, distance from the left infraorbital foramen to the nasion; DIOF, distance between the infraorbital foramina; PIOFN, perimeter formed by lines between the infraorbital foramina and the nasion; ATIOFN, area of the triangle formed by lines between the infraorbital foramina and the nasion

Introduction

The human skull is considered to be the second most useful indicator for determining an individual's sex, after the pelvis.¹ It is also responsible for furnishing important data for identifying the sex of living individuals and even cadavers.^{2,3} For forensic experts to perform these tasks, precise methods and techniques are required.^{4,5}

Male skulls are generally formed by coarser or rougher structures because of the presence of stronger muscle insertions. Moreover, they are also larger than female skulls.^{4,5}

Two methods are used in the process of determining individuals' sex from parts of the craniofacial skeleton: the qualitative or morphological method and the quantitative or metric method. Most studies have used the morphological method, in which characteristics such as the frontal sinuses, teeth, glabella, bone thickness, eyebrow ridge thickness and chin shape have been studied.⁵ On the other hand, the quantitative method makes use of measurements between pre-established points on the skull. This method, which is considered to possibly be more efficient, has contributed greatly towards determining individuals' sex.^{2,6}

Most authors in the literature on skull measurements from quantitative or metric variables have used samples from other countries (i.e. not from Brazil).⁶⁻⁸ This places limitations on the

applicability of these studies to the Brazilian population, given that without knowledge of the parameters for measurements on Brazilian skulls, researchers are obliged to use international tables. This could lead to uncertainty regarding the results.²

Therefore, the present study had the objective of determining sexual dimorphism in Brazilian skulls from measurement of the area of the triangle formed by lines between the infraorbital foramina (IOFs) and the nasion.

Materials and methods

In this study, 242 dry human skulls were used: 148 male skulls and 94 female skulls, ranging in age from 18 to 91 years. The skulls had been obtained in accordance with law no. 8501 of 1992, which deals with use of unclaimed cadavers for use in studies and research. These skulls belong to the Study and Research Center for Anatomy and Forensic Anthropology (CEPAAF) of Tiradentes University.

This was a morphometric study in which the following were measured: distance of the right infraorbital foramen to the nasion (DRIOFN), distance of the left infraorbital foramen to the nasion (DLIOFN), distance between the infraorbital foramina and perimeter and area of the triangle formed by lines between the infraorbital foramina and the nasion (Figure 1). The linear measurements were made with the aid of digital calipers with accuracy to the nearest 0.01 mm.

The data were analyzed both descriptively and analytically. The normality of distribution of the numerical variables was assessed by means of the Shapiro-Wilk test. Variables that met the assumption of normal distribution were presented as the mean (M) and 95% confidence interval. Spearman's linear correlation test was applied and the values obtained were categorized as follows: 0 to 0.39 as weak

correlation; 0.40 to 0.69 as moderate correlation; and 0.70 to 1.00 as strong correlation. To compare morphometric values according to sex, the Mann-Whitney test was applied. To evaluate multivariate differences we apply MANOVA and Lambda Wilk's Test. The statistical significance level was stipulated as 5% ($p \leq 0.05$). For all the analyses, the Statistical Package for the Social Sciences (SPSS 15.0) software was used.




Figure 1 Triangle formed by the infraorbital foramina and the nasion.

N, nasion; RIOF, right infraorbital foramen; LIOF, left infraorbital foramen; DRIOFN, distance from the right infraorbital foramen to the nasion; DLIOFN, distance from the left infraorbital foramen to the nasion; DIOF, distance between the infraorbital foramina; ATIOFN, area of the triangle formed by lines between the infraorbital foramina and the nasion

Results

In the sample of 148 male skulls and 94 female skulls, there were no abnormalities with regard to age. The median age of the male skulls was 57 years (41.75-70) and the median of the female skulls was 56.50 years (41.50-70). There was no statistically significant difference in age between the sexes of the skulls ($p = 0.887$).

In relation to sex, there were statistically significant differences in the measurements of DIOF, DRIOFN, DLIOFN and the perimeter and area of the triangle delimited by the infraorbital foramina and the nasion (Table 1) and a multivariate difference in measurements.

Table 1 Morphometric variables in relation to sex

Variables	Male M(IC95%)	Female M(IC95%)	p
DIOF	56.0(53.0-58.8)	53.7(50.7-56.0)	<0.001*
DRIOFN	46.2(44.1-49.0)	44.7(42.0-47.0)	<0.001*
DLIOFN	46.0(44.0-48.0)	44.0(41.8-46.1)	<0.001*
PIOFN	147.7(142.5-154.0)	142.5(135.0-147.6)	<0.001*
ATIOFN	1023.9(944.6-1105.3)	946.5(853.8-1025.3)	<0.001*

DIOF, distance between the infraorbital foramina; DRIOFN, distance from the right infraorbital foramen to the nasion; DLIOFN, distance from the left infraorbital foramen to the nasion; PIOFN, perimeter formed by lines between the infraorbital foramina and the nasion; ATIOFN, area of the triangle formed by lines between the infraorbital foramina and the nasion Mann-Whitney test; *Significance ($p \leq 0.05$). MANOVA, Lambda Wilk's=0.82, $F_{94,148} = 10.12, p < 0.001$

The results from the comparisons of measurements between the sexes showed that all of them were greater in the male skulls. It could be seen from Spearman's linear correlation that all the correlations between age and all the measurements of distance, perimeter and area were weak or insignificant (Table 2).

Table 2 Correlation analysis between age and all the measurements of distance, perimeter and area

	DIOF	DRIOFN	DLIOFN	PIOFN	ATIOFN
Age	0.049	-0.002	-0.013	0.011	0.009
Rho (P-value)	-0.447	-0.978	-0.836	-0.868	-0.892

DIOF, distance between the infraorbital foramina; DRIOFN, distance from the right infraorbital foramen to the nasion; DLIOFN, distance from the left infraorbital foramen to the nasion; PIOFN, perimeter formed by lines between the infraorbital foramina and the nasion; ATIOFN, area of the triangle formed by lines between the infraorbital foramina and the nasion

Discussion

Determination of sex is considered to be one of the main pillars of human identification protocols. This can be achieved through analysis of cranial measurements and visual evaluation of the skull, among other means.^{1,4,5,7,9-14} Because craniofacial structures are seldom destroyed, they have been widely used by several authors for determining sex.^{1-6,9,10,15,16} The quantitative method was used in the present study to determine sex. Some authors have considered that this method is more reliable when pre-established points are used and have accepted that qualitative methods are subjective and unsafe.^{2-6,11} Almeida Junior et al.¹⁷ using the logistic regression technique, for this type of study, obtained a score of 71.8%.

We did not find any use of the triangular area formed between the infraorbital foramina and the nasion for determining sexual dimorphism, in the literature. Almeida Junior et al.⁵ conducted a similar study, but they used the distance from the infraorbital foramina to the prosthion as craniometric reference points. These authors found that the mean distance between the infraorbital foramina was 60.94 mm for male skulls and 58.25 mm for female skulls. Regarding the distance from the infraorbital foramina to the prosthion, the mean was 31.30 mm for male skulls and 29.47 mm for female skulls. Regarding the area formed by lines between the infraorbital foramina and the prosthion, the mean was 956.68 mm² for male skulls and 860.46 mm² for female skulls. In the present study, the mean DIOF was 56.0 mm for the male skulls and 53.7 mm for the female skulls. Regarding the area of the triangle, the mean obtained was 1023.9 mm² for the male skulls and 946.5 mm² for the female skulls. For all the variables studied, the means were greater in the male skulls.

Badam et al.⁹ and Patil & Mody³ analyzed linear skull measurements obtained via lateral cephalometry. In both studies, the nasion was used as a reference point in order to establish and compare results between the sexes. Like in the present study, the values obtained were greater in the male skulls. In our study, comparisons of the DIOF, DRIOFN, DLIOFN and the perimeter and area of the triangle formed by lines between the infraorbital foramina and the nasion, in relation to the sexes, all showed statistically significant differences, such that all of the measurements were greater in the male skulls. This may be explicable in terms of the anatomical differences between male and female skulls.

Studies involving visual analyses on skulls have indicated differences between the sexes. For example, prominences, bone crests and other anatomical abnormalities are more perceptible in male skulls than in female skulls.^{2,4,5,9,16-19} However, according to Badam et al.,⁹ Patil & Mody,³ Shearer et al.¹³ and Spradley & Jantz,¹ metric analysis makes it possible to obtain results that are more objective, especially for researchers with little experience of determining individuals' sex. Moreover, this facilitates the statistical analysis on the sample and comparison with other studies.

Many studies conducted around the world have used quantitative variables in their analyses.^{7,10,12,13,16-18,20-23} In these analyses, the researchers only used international data because they did not have any measurements on Brazilian skulls. According to Almeida Junior et al.,⁴ Almeida Junior et al.,⁵ Bigoni et al.,¹⁷ Francesquini Junior et al.,² Franklin et al.,¹⁸ Gapert et al.,⁷ Kimmerle et al.¹¹ and Kruger et al.,¹² this may compromise the analysis because of the possibility that a variety of factors such as climate, local geography, diet, socioeconomic conditions and quality of life might interfere with defining an individual's sex. In this regard, Almeida Junior et al.⁴ accepted that difficulties in the process of determining sex might arise through variations in measurements and morphology found in different populations. However, authors such as Ferreira et al.,²⁴ Almeida Junior et al.²⁵ and Lima et al.²⁶ believe in the possibility of using their findings, in Brazilian skulls, in services of Forensic Dentistry and in the practical field of Forensic Anthropology.

Conclusion

The area of the triangle formed by lines between the infraorbital foramina and the nasion, and also all the other variables, were greater in the male skulls, with a statistically significant difference. This means that anatomical characteristics are of fundamental importance in distinguishing between the sexes, given that factors such as climate, local geography, diet, socioeconomic condition and quality of life may interfere in defining an individual's sex.

Acknowledgements

None.

Conflict of interest

Author declares that there is no conflict of interest.

References

1. Spradley MK, Jantz RL. Sex estimation in forensic anthropology: skull versus postcranial elements. *J Forensic Sci.* 2011;56(2):289-296.
2. Francesquini Junior L, Francesquini MA, Cruz BM, et al. Identification of sex using cranial base measurements. *J Forensic Odontostomatol.* 2007;25(1):7-11.
3. Patil KR, Mody RN. Determination of sex by discriminant function analysis and stature by regression analysis: a lateral cephalometric study. *Forensic Sci Int.* 2005;147(2-3):175-180.
4. Almeida Junior E, Reis FP, Galvão LCC, et al. Investigação do sexo e idade por meio de mensurações interforames em crânios secos de adultos. *Rev Ciênc Méd Biol.* 2013;12(1):55-59.
5. Almeida Junior E, Araújo TM, Galvão LCC, et al. Investigação do sexo através de uma área triangular facial formada pela interseção dos pontos: forame infra-orbital direito, esquerdo e o próstio, em crânios secos de adultos. *Rev Ciênc Méd Biol.* 2010;9(1):8-12.
6. Hatipoglu HG, Ozcan HN, Hatipoglu US, et al. Age, sex and mass index in relation to calvarial diploe thickness and craniometric data on MRI. *Forensic Sci Int.* 2008;182(1-3):46-51.
7. Gapert R, Black S, Last J. Sex determination from the foramen magnum: discriminant function analysis in an eighteenth and nineteenth century British sample. *Int J Legal Med.* 2009;123(1):25-33.
8. Konigsberg IW, Algee Hewitt BF, Steadman DW. Estimation and evidence in forensic anthropology: sex and race. *Am J Phys Anthropol.* 2009;139(1):77-90.
9. Badam RK, Manjunath M, Rani MS. Determination of sex by discriminant function analysis of lateral radiographic cephalometry. *Journal of Indian Academy of Oral Medicine and Radiology.* 2011;23(3):179-183.
10. Jiménez Arenas JM, Esquivel JA. Comparing two methods of univariate discriminant analysis for sex discrimination in an Iberian population. *Forensic Sci Int.* 2013;228(1-3):175.e1-e4.
11. Kimmerle EH, Ross A, Slice D. Sexual dimorphism in America: geometric morphometric analysis of the craniofacial region. *J Forensic Sci.* 2008;53(1):54-57.
12. Kruger GC, L'Abbé EN, Stull KE, et al. Sexual dimorphism in cranial morphology among modern South Africans. *Int Legal Med.* 2014;129(4):869-875.
13. Shearer BM, Sholts SB, Garvin HM, et al. Sexual dimorphism in human browridge volume measured from 3D models of dry crania: A new digital morphometrics approach. *Forensic Sci Int.* 2012;222(1-3):400.e1-400.e5.
14. Vidya CS, Shamasundar NM, Manjunatha B, et al. Evaluation of size and volume of maxillary sinus to determine gender by 3D computerized tomography scan method using dry skulls of South Indian origin. *Int J Res Rev.* 2013;5(3):98.
15. Cornélio Neto WL, Bérzin F, Daruge Júnior E, et al. Identificação antropológica: superposição de imagens pelos ossos nasais. *Rev Ciênc Méd Biol.* 2011;10(2):105-111.
16. Green H, Curnoe D. Sexual dimorphism in Southeast Asian crania: morphometric approach. *Homo.* 2009;60(6):517-534.
17. Almeida Júnior E, Reis FP, Galvão LCC, et al. Estimativa do sexo e idade por Meio de mensurações cranianas. *Rev Bahiana Odonto.* 2015;6(2):81-88.
18. Bigoni L, Velemínská J, Bruzek J. Three-dimensional geometric morphometric analysis of cranio-facial sexual dimorphism in a Central European sample of known sex. *Homo.* 2010;61(1):16-32.
19. Franklin D, Cardini A, Flavel A, et al. The application of traditional and geometric morphometric analyses for forensic quantification of sexual dimorphism: preliminary investigation in a Western Australian population. *Int J Legal Med.* 2012;126(4):549-558.
20. Perri RA, Kairaitis K, Wheatley JR, et al. Anthropometric and craniofacial sexual dimorphism in obstructive sleep apnea patients: is there male-female phenotypical convergence? *J Sleep Resp.* 2014;24(1):82-91.
21. Franklin D, Cardini A, Flavel A, et al. Estimation of sex from cranial measurements in a Western Australian population. *Forensic Sci Int.* 2013;229(1-3):158.e1-158.e8.
22. Marinescu M, Panaitescu V, Rosu M, et al. Sexual dimorphism of crania in Romanian population: discriminant function analysis approach for sex estimation. *Rom J Leg Med.* 2014;22:21-26.
23. Steyn M, Íscan MY. Sexual dimorphism in the crania and mandibles of South African whites. *Forensic Sci Int.* 1998;98(1-2):9-16.

24. Ferreira RFA, Neves FS, Almeida Júnior E, et al. Avaliação do dimorfismo sexual por meio de medidas lineares entre os processos mastoideos e a espinha nasal anterior em crânios secos humanos. *J Health Sci Inst.* 2015;33(2):130–134.
25. Almeida Júnior, Reis FP, Galvão LCC, et al. Investigação do sexo e idade por meio de mensurações no palato duro e base de crânios secos de adultos. *Rev Ciênc Méd Biol.* 2016;15(2): 172-177.
26. Lima AIC, da Silva RA, Almeida Júnior E. Análise entre os pontos zigomáticos orbitais e espinha nasal anterior na investigação do sexo e idade em crânios secos de adultos. *J Health Sci Inst.* 2016;34(1):11–16.