

Bioinformatic analysis of glycoside hydrolases in the proteomes of mesophilic and thermophilic *Actinobacteria*

Abstract

Petroleum reserves are rapidly depleting and alternative renewable sources of energy need to be developed to meet the energy demands of the planet. Lignocellulose has been recognized as a highly promising and renewable resource for the development of clean energy. Thermophilic microbes and thermostable enzymes are being sought for biological conversion of lignocellulose into biofuels. The phylum *Actinobacteria* includes several efficient cellulose-degrading microorganisms. Genomes of several *Actinobacteria* have been completely sequenced and deposited in public databases, which are a great resource for uncovering new enzymes and targets for biotechnology. We searched the predicted proteomes of 69 *Actinobacteria* for the homologs of 20 glycoside hydrolase families relevant to lignocellulose degradation and identified 589 glycoside hydrolase homologs. We analyzed (1) the distribution of the glycoside hydrolase homologs across mesophilic and thermophilic *Actinobacteria* (2), the domain architecture of cellulases (from GH5 and GH6 families) and xylanases (from GH10 and GH11 families) from mesophilic and thermophilic *Actinobacteria*, and (3) asymmetric amino acid substitutions between mesophilic and thermophilic glycoside hydrolases. Overall, our data provide new insights into the distribution of different glycoside hydrolases in *Actinobacteria* as well as into the thermostability features of cellulases and xylanases from *Actinobacteria*. Our findings provide a basis for genetic engineering of glycoside hydrolases as well as new targets for biotechnology.

Keywords: thermophiles, enzymes, cellulases, xylanases, biofuel, lignocellulose, genome, proteome

Volume 5 Issue 3 - 2017

Kip A Teegardin, Steven James, Ravi D Barabote

Department of Biological Sciences, University of Arkansas, USA

Correspondence: Ravi D. Barabote, Department of Biological Sciences, University of Arkansas, Fayetteville, AR 72701, USA., Tel (479) 5752475, Fax (479) 5754010, Email barabote@uark.edu

Received: January 30, 2017 | **Published:** March 16, 2017

Abbreviations: GH, glycoside hydrolases; CBM, carbohydrate binding module; OGT, optimal growth temperature

Introduction

Petroleum fuels are finite and non-renewable and they pose a significant concern for global climate, sustainability, and international security.¹ Alternative renewable sources of energy are urgently needed to meet the current global challenges. Plants are the most abundant source of renewable carbon on Earth. Plant cell wall (lignocellulose) can be used for the production of renewable, sustainable, and environmentally -clean biofuels.² Lignocellulose is mainly composed of polymers of sugars (cellulose and hemicellulose) and phenolic units (lignin). While complex lignocellulose can be converted into liquid fuels thermo-chemically, biological transformation of lignocellulosic polysaccharides using microorganisms and microbial enzymes is an economical and environmentally benign process for sustainable production of biofuels.^{3,4} Several microorganisms produce glycoside hydrolase enzymes such as cellulases and xylanases that break down cellulose and xylan (hemicellulose), respectively.⁵ Efficient lignocellulose-degrading microorganisms and catalytically- superior cellulases and xylanases are of very high value in the bioconversion of lignocellulose into biofuels.^{6,7}

Actinobacteria are a phylum of Gram-positive bacteria that are found abundantly in soil.⁸ They include some of the most prolific lignocellulose-degrading bacteria.⁹ *Actinobacteria* include both mesophilic and thermophilic members. Many new *Actinobacteria* continue to be isolated and sequenced in bioprospecting studies aimed

at identifying new biotechnological targets.¹⁰ Growing number of completely sequenced genomes are being steadily deposited in public databases, which provide an expanding resource for discovering novel targets for biotechnology. Systematic bioinformatic mining of the genomes and predicted proteomes of sequenced *Actinobacteria* has the potential to reveal novel insights into lignocellulose-degrading enzymes for bioenergy applications.¹¹

Thermophilic microbes and thermostable enzymes are most useful for the development of cost- effective, industrial scale technologies.¹² Thermostability of enzymes increases their shelf life, reduces reaction times, improves industrial productivity, and lowers manufacturing costs.¹² Thus, enzyme thermostability is a highly desirable property for industrial enzymatic deconstruction of lignocellulose. Valuable insights can be gleaned about factors that contribute to thermostability by performing comparative analysis of amino acid sequences of proteins from mesophilic and thermophilic organisms.¹³ Such insights can be exploited for designing and genetically engineering enhanced enzymes for industrial applications.

In this study, we systematically analyzed the predicted proteomes of 69 *Actinobacteria* for homologs of glycoside hydrolase enzymes that are relevant to lignocellulose degradation. We analyzed the distribution of the homologs across the phylum. We identified homologs from mesophilic and thermophilic *Actinobacteria* and analyzed their domain architecture to decipher thermophilic patterns. Finally, we analyzed the amino acid sequences of cellulases and xylanases from mesophilic and thermophilic *Actinobacteria* and identified asymmetric amino acid substitution patterns in the thermophilic enzymes.

Methodology

Predicted proteomes of known lignocellulose-degrading *Actinobacteria* were obtained from NCBI (ftp://ftp.ncbi.nlm.nih.gov/). Optimal growth temperature (OGT) information was obtained through literature. Organisms were classified as mesophilic (OGT<40°C) or thermophilic (OGT>40°C). Glycoside hydrolase families that contain lignocellulose degradation enzymes were identified from the CAZy database.¹⁴ Representative *Actinobacterial* sequences from the CAZy families were used to identify homologs in the proteomes of the *Actinobacteria* using BLAST.¹⁵ Domains in the glycoside hydrolase proteins were identified using the NCBI's CDD- search tool.¹⁶ Amino acid substitutions between homologs of mesophilic and thermophilic *Actinobacteria* were identified using multiple alignments as described previously.¹⁷ Briefly, for each GH family, orthologs from mesophilic and thermophilic *Actinobacteria* were aligned using CLUSTAL.^{18,19} Each substitution was counted only once per position in the alignment. For each amino acid substitution pair (e.g., AMBT and ATBM where A and B represent amino acids and the subscripts M and T represent mesophilic and thermophilic organisms, respectively), the total number of substitutions over the entire alignment was summed and the percentage of each substitution within the pair was calculated. Statistical significance (*p*-value)

of asymmetric amino acid substitutions between the two groups of organisms was calculated using a binomial function. The asymmetry (i.e., bias) in AMBT and ATBM substitutions was considered significant if their *p*-value was below the threshold.

Results and discussion

Distribution of glycoside hydrolases in *Actinobacteria*

We identified a total of 1133 *Actinobacteria* in the NCBI database. Of these, genomes of only 236 (21%) *Actinobacteria* have been completed sequenced. Within the 236 sequenced *Actinobacteria*, we identified 69 (29%) organisms that have been described in literature to have cellulolytic activity. We analyzed the predicted proteomes of the 69 *Actinobacteria* for the presence of glycoside hydrolases relevant to lignocellulose degradation. In addition, we collected information on their optimal growth temperature for each organism from literature. Using the CAZy database, we identified 20 glycoside hydrolase families that contain enzymes known to hydrolyze various plant cell wall polysaccharides.¹⁴ A total of 589 glycoside hydrolase homologs were identified in the proteomes of the 69 *Actinobacteria* (Table 1). Of the 69 *Actinobacteria*,²⁰ 61 organisms are mesophilic and only 8 are thermophilic. This highlights the need to sequence more thermophilic *Actinobacteria*.

Table 1 Summary of the analysis of *Actinobacteria* used in this study

Organism name	OGT (°C)	# of GH families	Total GH homologs
<i>Acidothermus cellulolyticus</i> I 1B ATCC 43068	55	7	12
<i>Actinosynnema mirum</i> DSM 43827	28	9	30
<i>Amycolatopsis mediterranei</i> S699	26	13	37
<i>Amycolatopsis mediterranei</i> U32	26	12	33
<i>Bifidobacterium adolescentis</i> 15703	50	3	4
<i>Bifidobacterium animalis</i> AD011	37	1	1
<i>Bifidobacterium animalis</i> ATCC 25527	37	1	1
<i>Bifidobacterium animalis</i> B420	37	1	1
<i>Bifidobacterium animalis</i> Bb12	37	2	2
<i>Bifidobacterium animalis</i> Bi-04	37	1	1
<i>Bifidobacterium animalis</i> Bi-07	37	1	2
<i>Bifidobacterium animalis</i> BLC1	37	1	1
<i>Bifidobacterium animalis</i> CNCM I-2494	37	2	3
<i>Bifidobacterium animalis</i> DSM 10140	37	1	1
<i>Bifidobacterium animalis</i> V9	37	1	1
<i>Bifidobacterium bifidum</i> PRL2010	37	1	2
<i>Bifidobacterium bifidum</i> S17	37	1	2
<i>Bifidobacterium breve</i> ACS-071-V-Sch8b	37	2	2
<i>Bifidobacterium breve</i> UCC2003	37	0	0
<i>Bifidobacterium dentium</i> Bd1	29	2	7
<i>Bifidobacterium Longum</i> 157F	34	2	6
<i>Bifidobacterium Longum</i> BBMN68	34	1	3
<i>Bifidobacterium Longum</i> DJO10A	34	1	2
<i>Bifidobacterium Longum</i> F8	34	1	1
<i>Bifidobacterium Longum</i> JCM 1217	34	2	5
<i>Bifidobacterium Longum</i> JCM 1222 ,ATCC 15697	34	2	3

Table Continued...

Organism name	OGT (°C)	# of GH families	Total GH homologs
<i>Bifidobacterium Longum JDM301</i>	34	2	5
<i>Bifidobacterium Longum KACC 91563</i>	34	2	4
<i>Bifidobacterium Longum NCC2705</i>	34	2	5
<i>Cellulomonas fimi ATCC 484</i>	40	11	34
<i>Cellulomonas flavigena DSM 20109</i>	30	9	37
<i>Cellvibrio gilvus ATCC 13127</i>	25	9	20
<i>Clavibacter michiganensis NCPPB 382</i>	37	3	7
<i>Clavibacter michiganensis sepedonicus</i>	37	1	2
<i>Jonesia denitrificans DSM 20603</i>	37	9	13
<i>Micrococcus luteus</i>	37	0	0
<i>Micromonospora aurantiaca ATCC 27029</i>	27	10	22
<i>Modestobacter marinus BC501</i>	28	0	0
<i>Mycobacterium abscessus</i>	30	2	2
<i>Mycobacterium avium 104</i>	37	3	3
<i>Mycobacterium avium K-10</i>	37	3	3
<i>Mycobacterium bovis AF2122/97</i>	35	4	5
<i>Mycobacterium bovis BCG str. Mexico</i>	35	3	3
<i>Mycobacterium bovis Pasteur 1173P2</i>	35	3	3
<i>Mycobacterium bovis Tokyo 172</i>	35	3	3
<i>Mycobacterium gilvum PYR-GCK</i>	30	3	3
<i>Mycobacterium marinum</i>	37	2	2
<i>Mycobacterium smegmatis MC2 155</i>	30	3	4
<i>Rhodococcus erythropolis PR4 PR4 (= NBRC 100887)</i>	25	2	2
<i>Rhodococcus opacus B4</i>	27	2	2
<i>Saccharomonospora glauca</i>	45	0	0
<i>Saccharomonospora viridis DSM 43017</i>	55	2	2
<i>Streptomyces avermitilis MA-4680</i>	32	9	19
<i>Streptomyces bingchenggensis BCW-1</i>	28	12	47
<i>Streptomyces cattleya DSM 46488,8057</i>	34	4	12
<i>Streptomyces clavuligerus</i>	28	0	0
<i>Streptomyces coelicolor A3(2)</i>	28	11	20
<i>Streptomyces flavogriseus ATCC 33331</i>	28	11	19
<i>Streptomyces hygrosopicus jinggangensis 5008</i>	35	9	13
<i>Streptomyces pristinaespiralis</i>	28	0	0
<i>Streptomyces scabiei 87.22</i>	27	10	27
<i>Streptomyces sirex AA3</i>	28	8	9
<i>Streptomyces sviveus</i>	28	1	1
<i>Streptomyces violaceusniger Tu 4113</i>	28	10	22
<i>Streptosporangium roseum DSM 43021</i>	28	9	14
<i>Thermobifida fusca YX</i>	55	8	12
<i>Thermobispora bispora 43833</i>	55	8	10
<i>Thermomonospora curvata 43183</i>	55	2	3
<i>Xylanimonas cellulositytica DSM 15894</i>	30	7	12

OGT (°C), optimal growth temperature (degrees Celsius); GH, glycoside hydrolase

We analyzed the relationship between optimal growth temperature and glycoside hydrolases encoded in the proteomes of the *Actinobacteria* (Figure 1). In general, there was very poor correlation ($R^2 < 0.1$) between optimal growth temperature and glycoside hydrolase content of the proteomes. However, this may be partly due to the overrepresentation of mesophilic *Actinobacteria* in the dataset. The 61 mesophilic *Actinobacteria* encoded between 0 and 13 glycoside hydrolase families with an average of 4.0 ± 3.9 , while they encoded between 0 and 47 homologs of glycoside hydrolases with an average of 8.4 ± 10.9 . The 8 thermophilic *Actinobacteria* encoded between 0 and 11 glycoside hydrolase families with an average of 5.1 ± 3.9 , while they encoded between 0 and 34 homologs of glycoside hydrolases with an average of 9.6 ± 10.9 . There were no statistically significant differences in the distribution of glycoside hydrolases between mesophilic and thermophilic *Actinobacteria*. However, substantially greater numbers of thermophilic *Actinobacteria* need to be sequenced before deciphering any underlying biases between the two groups of *Actinobacteria*.





Figure 1 Relationship between optimal growth temperature and glycoside hydrolases in *Actinobacteria*.

(A) Scatter plot of number of glycoside hydrolases (GH) families versus optimal growth temperature.

(B) Scatter plot of number GH homologs versus optimal growth temperature. Best-fit line with R-squared value is shown.

We analyzed relative abundances of the 20 glycoside hydrolase families across *Actinobacteria* (Figure 2). The data show that GH5 was the most highly represented family in the *Actinobacteria*.

It was the only family that was found in majority (70%) of the organisms analyzed. The GH6 and GH43 families were the next most represented families and were found in 48% of the *Actinobacteria*. The GH5 family is known to contain cellulose- and hemicellulose-degrading enzymes, while the GH6 family contains cellulases and the GH43 family contains hemicellulases.¹⁴ The GH45, GH51, and



GH128 families were not represented in any of the *Actinobacteria* in our dataset. Other GH families showed intermediate representation.

Figure 2 Relative abundance of glycoside hydrolase (GH) families in *Actinobacteria*. Percentage of *Actinobacteria* containing homologs of the different GH families are plotted.

Domain architecture of glycoside hydrolases in *Actinobacteria*

To minimize over-representation of mesophilic *Actinobacteria* in the dataset, we selected one representative species per genus and also retained saprophytic free-living bacteria while removing animal and human pathogens. This yielded a more balanced set of *Actinobacteria* (6 thermophiles and 8 mesophiles). We focused our analysis on four GH families - cellulases from GH5 and GH6 families and xylanases from GH10 and GH11 families. There were 113 glycoside hydrolases from the four families across the 14 *Actinobacteria* (Table 2). There were 77 homologs in the 8 mesophilic bacteria, and 36 homologs in the 6 thermophilic bacteria. Six organisms contained representatives from all four families, while five organisms contained representatives from only three families and two organisms contained homologs from just one family. We analyzed the domain architecture of the 113 glycoside hydrolases using the NCBI's CDD-search tool.¹⁶ At least five different types of carbohydrate binding modules (CBMs - CBM-2, CBM-3, CBM-X2, CBM-9, and CBM-4-9) were found fused to the catalytic domains of glycoside hydrolases (Table 3). Further analysis revealed a bias in the presence and location of certain CBMs. For example, CBM-2 was found fused on the C-terminal side of the catalytic domain in all four glycoside hydrolase families, while it was found on the N-terminal side of the catalytic domain in GH5 and GH6 cellulases. CMB -3 was only found in homologs from thermophilic *Actinobacteria*, and it always occurred C-terminal to the catalytic hydrolase domain. CBM-9 and CMB-4-9 were found attached to only GH10 xylanases. CBM-9 occurred C-terminal to the catalytic domain, while CBM-4-9 was found on the N-terminal side of the hydrolase domain. CBM-X2 was found only in GH5 hydrolases from mesophilic *Actinobacteria* and was found C-terminal to the hydrolase domain. These data suggest that there are positional constraints for CBM domains in glycoside hydrolases. Certain domains may be required for the functioning and stability of the enzymes, while others may be specific to the substrates hydrolyzed by the associated catalytic domains.

Table 2 Distribution of glycoside hydrolases in mesophilic and thermophilic Actinobacteria

Organism name	OGT	GH5	GH6	GH10	GH11
(A) Mesophilic Actinobacteria					
<i>Actinosynnema mirum</i> DSM 43827	28	5	3	4	1
<i>Amycolatopsis mediterranei</i> S699	26	4	2	7	1
<i>Cellvibrio gilvus</i> ATCC 13127	25	2	4	6	0
<i>Jonesia denitrificans</i> DSM 20603	37	0	2	4	1
<i>Micromonospora aurantiaca</i> ATCC 27029	27	4	2	4	1
<i>Streptomyces coelicolor</i> A3(2)	28	1	3	2	2
<i>Streptosporangium roseum</i> DSM 43021	28	1	3	1	0
<i>Xylanimonas cellulosilytica</i> DSM 15894	30	0	2	4	1
(B) Thermophilic Actinobacteria					
<i>Acidothermus cellulolyticus</i> 11B	55	2	2	2	0
<i>Cellulomonas fimi</i> ATCC 484	40	0	6	8	1
<i>Saccharomonospora viridis</i> DSM 43017	55	0	0	1	0
<i>Thermobifida fusca</i> YX	55	2	2	2	1
<i>Thermobispora bispora</i> DSM 43833	55	1	2	2	1
<i>Thermomonospora curvata</i> DSM 43183	55	0	1	0	0

Table 3 Domain architecture of glycoside hydrolases in Actinobacteria

		C-Terminal domain									
(A) Mesophiles		GH5	GH6	GH10	GH11	CBM_2	CBM_3	CBM_X2	CBM_9	CBM_4_9	No CBM
	GH5					8		2			2
	GH6					8					6
	GH10					16			2		7
	GH11					4					1
N-terminal domain	CBM_2	7	8					2			
	CBM_3										
	CBM_X2										
	CBM_9										
	CBM_4_9			4		1			2		
		C-Terminal domain									
(B) Thermophiles		GH5	GH6	GH10	GH11	CBM_2	CBM_3	CBM_X2	CBM_9	CBM_4_9	No CBM
	GH5					3	2				
	GH6					5	1				3
	GH10					8	1		2		3
	GH11					3					
N-terminal domain	CBM_2	1	1								
	CBM_3										
	CBM_X2										
	CBM_9										
	CBM_4_9			2					2		

Asymmetric amino acid substitutions in glycoside hydrolases

We wanted to understand amino acid mesophilic *Actinobacteria*. This would help identify amino acid substitutions that may contribute to thermostability of glycoside hydrolases. For each glycoside hydrolase family, we aligned only the hydrolase domains of orthologs from mesophilic and thermophilic organisms identified earlier (Table 2). We calculated the frequencies of all amino acid substitutions between mesophilic and thermophilic homologs at every position and identified the statistically significant asymmetric amino acid substitutions (Table 4). The data revealed 41 pairs of amino acid substitutions that are

asymmetric between the homologs from thermophilic and mesophilic *Actinobacteria*. Certain amino acid preferences in the thermophiles were specific to the glycoside hydrolase family, while other amino acid preferences were independent of the glycoside hydrolase family. For example, thermophilic enzymes from GH6, GH10, and GH11 families showed preferences for alanine over glycine. Similarly, thermophilic proteins showed preference for aspartate over thermolabile serine and threonine residues. There was also a biased preference for isoleucine over valine in thermostable homologs. Overall, the data provide several new targets for genetically engineering higher thermostability in glycoside hydrolases²⁰ (Table 4).

Table 4 Asymmetric amino acid substitution patterns in glycoside hydrolases

		Amino acid in the thermophilic homologs														
		A	D	E	F	H	I	K	L	N	P	Q	S	T	V	Y
Amino acid in the mesophilic homologs	A		C74	C62	A100	B100					C77	B100	C61, B68			B100
	D	B92								B93						
	E		A62													
	F															C65
	G	B88, C67, D86,	C100					B91								
	K			A86												A100
	L						A62									
	N	B100	D91			D92							D84	D100	C100	
	R			C67					B90			C59				
	S	D86	B77, C67						C100			C64		D85		
	T		A80, C75				B100, C78				B100				B88	D100
	V	B80					A60, C56		D77							
	Y	A100														

Standard single letter amino acid code is used to represent amino acids. Data are represented with a letter followed by a number, where A represents GH5, B represents GH6, C represents GH10, D represents GH11, and numbers represent the percentage of occurrence of the particular substitution. Only statistically significant ($p < 0.1$) asymmetric substitutions are shown.

Conclusion

We analyzed the predicted proteomes of 69 sequenced *Actinobacteria* and identified homologs of 20 glycoside hydrolase families associated with plant cell wall degradation. Some glycoside hydrolase families were well represented across the phylum, while a few families were not represented in any of the *Actinobacteria* we analyzed. The glycoside hydrolases appear to have a constrained domain architecture that likely determines their stability, functioning, and interaction with substrates. Certain carbohydrate binding modules found fused to the glycoside hydrolases were only associated with thermophilic *Actinobacteria*. Finally, glycoside hydrolases from thermophilic *Actinobacteria* showed preferences for certain amino acid substitutions over their mesophilic counterparts. Overall, our data provide new insights into glycoside hydrolases in *Actinobacteria* and provide a basis for genetically enhancing the stability of glycoside hydrolases towards industrial applications.

Acknowledgements

This research was supported by startup funds provided to RDB by the University of Arkansas. KAT acknowledges support from the National Science Foundation Research Experience for Undergraduates program through the University of Arkansas REU Site (DBI-1063067).

Conflict of interest

The author declares no conflict of interest.

References

- Solomon BD. Biofuels and sustainability. *Ann N Y Acad Sci.* 2010;1185:119–134.
- Pothiraj C, Kanmani P, Balaji P. Bioconversion of lignocellulose materials. *Mycobiology.* 2006;34(4):159–165.

3. Peralta–Yahya PP, Keasling JD. Advanced biofuel production in microbes. *Biotechnol J*. 2010;5(2):147–162.
4. Clark JH, Luque R, Matharu AS. Green chemistry, biofuels, and biorefinery. *Annu Rev Chem Biomol Eng*. 2012;3:183–207.
5. Cragg SM, Beckham GT, Bruce NC, et al. Lignocellulose degradation mechanisms across the Tree of Life. *Curr Opin Chem Biol*. 2015;29:108–119.
6. Blumer–Schuette SE, Brown SD, Sander KB, et al. Thermophilic lignocellulose deconstruction. *FEMS Microbiol Rev*. 2014;38(3):393–448.
7. Bhalla A, Bansal N, Kumar S, et al. Improved lignocellulose conversion to biofuels with thermophilic bacteria and thermostable enzymes. *Bioresour Technol*. 2013;128:751–759.
8. Ventura M, Canchaya C, Tauch A, et al. Genomics of *Actinobacteria*: tracing the evolutionary history of an ancient phylum. *Microbiol Mol Biol Rev*. 2007;71(3):495–548.
9. McCarthy AJ, Williams ST. *Actinomycetes* as agents of biodegradation in the environment—a review. *Gene*. 1992;115(1–2):189–192.
10. Ward AC, Bora N. Diversity and biogeography of marine *Actinobacteria*. *Curr Opin Microbiol*. 2006;9(3):279–286.
11. Ohm RA, Riley R, Salamov A, et al. Genomics of wood–degrading fungi. *Fungal Genet Biol*. 2014;72:82–90.
12. Bruins ME, Janssen AE, Boom RM. Thermozyms and their applications: a review of recent literature and patents. *Appl Biochem Biotechnol*. 2001;90(2):155–186.
13. Kumar S, Nussinov R. How do thermophilic proteins deal with heat? *Cell Mol Life Sci*. 2001;58(9):1216–1233.
14. Cantarel BL, Coutinho PM, Rancurel C, et al. The Carbohydrate–Active EnZymes database (CAZy): an expert resource for Glycogenomics. *Nucleic Acids Res*. 2009;37(Database issue):D233–D238.
15. Altschul SF, Gish W, Miller W, et al. Basic local alignment search tool. *J Mol Biol*. 1990;215(3):403–410.
16. Marchler–Bauer A, Derbyshire MK, Gonzales NR, et al. CDD:NCBI’s conserved domain database. *Nucleic Acids Res*. 2015;43(Database issue):D222–226.
17. Takami H, Takaki Y, Chee GJ, et al. Thermodaptation trait revealed by the genome sequence of thermophilic *Geobacillus kaustophilus*. *Nucleic Acids Res*. 2004;32(21):6292–6303.
18. Higgins, Desmond G, Sharp PM. CLUSTAL: A Package for Performing Multiple Sequence Alignment on a Microcomputer. *Gene*. 1988;73(1):237–244.
19. Larkin MA, Blackshields G, Brown NP, et al. Clustal W and Clustal X Version 2.0. *Bioinformatics*. 2007;23(21):2947–2948.
20. Gao SJ, Wang JQ, Wu MC, et al. Engineering hyperthermostability into a mesophilic family 11 xylanase from *Aspergillus oryzae* by in silico design of N–terminus substitution. *Biotechnol Bioeng*. 2013;110(4):1028–1038.