

# Addressing the Challenge of Iodine Deficiency in Developing Countries

## Abstract

Iodine deficiency causes stunted physical and mental growth as well as infertility, lethargy and cognitive impairment. Although iodine deficiency is endemic in most countries and is therefore a public health issue, it is preventable through adequate intake of iodine in diet. Recent global data estimated that 1.88 billion people are at risk of iodine deficiency with 241 million children taking inadequate iodine in diet. Majority of these children with inadequate dietary iodine intake lives in Africa and South/South-East Asia. Gains have been made towards the eradication of iodine deficiency disorders (IDDs) through the universal salt iodisation (USI) programme which has ensured greater coverage of iodine intake throughout the world. In developing countries however, the perennial problem of weaker institutions means that much needs to be done if the goal of eradicating IDDs are to be realized. This mini-review looks at what steps could be undertaken to maximize the odds of eradicating IDD through food fortification, strengthening of regulatory institutions, empowering small-scale salt manufacturers and conscientization of the populace to patronise not only iodised salt but also iodine-fortified foods.

**Keywords:** Iodine deficiency disorder; Universal salt iodisation programme; cretinism; Thyroxin

## Mini Review

Volume 5 Issue 3 - 2017

**Patrick Adu and David Larbi Simpong\***

*Department of Medical Laboratory Sciences, School of Allied Health Sciences, University of Cape Coast, Ghana*

**\*Corresponding author:** David Larbi Simpong, Department of Medical Laboratory Sciences, School of Allied Health Sciences, University of Cape Coast, Ghana, Email: [Dsimpong@ucc.edu.gh](mailto:Dsimpong@ucc.edu.gh)

**Received:** September 30, 2016 | **Published:** March 07, 2017

**Abbreviations:** USI: Universal Salt Iodisation; IDD: Iodine Deficiency Disorder; RTK: Rapid Test Kit; T3: Triiodothyronine; T4: Thyroxin

## Introduction


Iodine is an essential micro-nutrient necessary for the formation of thyroid hormones [T3 (triiodothyronine) and T4 (thyroxin)]. As thyroid hormones regulate proper neuronal development, deficiency of iodine can have grave consequences especially when it occurs during pregnancy as it negatively affects foetal and subsequent neonatal life leading to irreversible cognitive impairment [1-3]. Primarily, iodine deficiency is caused by inadequate dietary supply. The plethoras of disorders caused by iodine deficiency are collectively called iodine deficiency disorder (IDD), the most prominent of which are goitre and cretinism. In line with these devastating and irreversible adverse health implications of iodine deficiency, the WHO recommended the adoption of the universal salt iodisation (USI) programme [4] to help eliminate this nutritional deficiency. In addition to this, most countries have adopted fortification of foods such as oils either as alternative iodine source or to augment the USI initiative [5,6]. However, the salt fortification programme has many advantages as salt is used by almost everybody and therefore overcome the issue of logistics required for the dissemination of other forms of iodine supplementation approaches. Not surprisingly, most developing countries mainly rely on the USI programme to meet iodine needs of the populace. Although empirical evidence shows that effective implementation of the USI programme offers undisputed prospect towards eliminating iodine deficiency, IDD

is still prevalent in most developing countries. For example, the WHO Global database on iodine deficiency show that in some parts of Cote d'ivoire, Egypt, Nigeria, and Ghana, prevalence of iodine insufficiency could be as high as 33.8%, 31.2%, 42% and 71.3% respectively [7]. Furthermore, the WHO recently estimated the regional averages of household consumption of adequately iodised salt to be 50%, 59%, 65, and 69, for least developed countries, Sub-Saharan Africa, West and Central Africa, and South Asia respectively (Figure 1) [8].

This calls for some serious consideration for the evaluation and monitoring of the implementation of the USI programme in these developing countries. Most of these countries have weak institutions with inadequate infrastructures necessary for effectively assessing the iodine levels in the supposedly iodised salts on offer. To our knowledge, ensuring adequacy of salt iodisation may be the critical component in the chain of measures necessary to actualise the goals of the USI programme. This salt iodisation and subsequent storage conditions may need some critical evaluation to ensure that any gains made in the USI initiative are not derailed. Authorities in these developing countries must invest in their respective regulatory agencies tasked with ensuring that quality standards are met by salt producers before these iodised salts are released onto their respective markets. Based on the daily iodine requirement, estimated average salt intake of 10 g/day, as well as estimated iodine losses during production and consumption [9], the WHO recommends the incorporation of 20 to 40 parts per million (ppm) of iodine to salt at the point of production [10]. Thus the existing regulatory agencies must be empowered to ensure that the salts on their respective markets

meet these basic quality standards. Rapid test kits (RTKs) have revolutionized the diagnosis and management of malaria even in remote communities. Although RTKs have also been developed for quantitative iodine measurements [11,12], there is urgent need for optimization of iodine RTKs [13] to provide accurate quantitative estimates of iodine in salts that would empower

regional and district level regulatory agencies to properly monitor the USI programme. It must be pointed out that these monitoring and evaluation processes should be periodic so as to ensure that iodine levels are consistently maintained within safe and optimal limits.



**Figure 1:** Global picture of the household consumption of adequately iodized salt. [Source: Iodized Salt Consumption SOWC; UNICEF (<http://data.unicef.org/nutrition/iodine.html>); Accessed October 2014.

### Way forward

Public education through radio, television, whatsapp and other social network platforms must be given priority so as to properly sensitize the populace with regards to the adverse health implications of IDD. Such public educational campaigns have previously shown successful in countries [14]. Any such educational programmes should also conscientize the populace about the proper storage conditions and use of these iodised salts. In our experience, the iodised salts sold in most developing countries are in polythene bags where consumers cut openings to ease salt dispensation (Simpong & Adu, unpublished observations). As these openings are never closed, it is a feasible supposition that the high ambient temperature and humidity/moisture prevailing in these tropical countries may cause iodine levels in the salts to wane with the passage of time [15,16]. These

unfortunate practices on the part of consumers inadvertently negate any investment made by the iodised salt manufacturers. This could be prevented by encouraging salt manufacturers to package their products in easy to reseal cartons and/or re-sealable plastic containments. Additionally, there may also be the need to include community health nurses and disease control officers to drive these community-based educational programmes. The successes enjoyed by the Global Polio eradication Initiative [17,18] where these health professionals routinely canvas the cooperation of even the remotest communities should be ample evidence that their inclusion in any IDD eradication campaigns may guarantee some meaningful gains. In spite of the many years that has passed since the introduction of the USI programme, many published data indicates that a considerable number of people are not using iodised salt [19,20]. Much as these data are helpful, most of them offer only snapshots of iodine status of

segments of the population at different geographic locations in these IDD regions. Interestingly, Juan et al. [21] demonstrated that to obtain a reliable estimate of iodine status of any population, consideration should be given to the dietary iodine intake from all food sources in addition to the UIC measurements [21]. It is imperative for governments and heads of state of these IDD regions to demonstrate commitment to the IDD eradication by annually devoting portions of national health budgets for continuous surveillance and monitoring of the USI programme. Such funds should be used purposely to encourage a concerted research to provide regional and ultimately national iodine status estimates to inform intervention strategy formulation. These population-based research/surveillance studies could effectively use the UIC [22] to provide credible data on present iodine status to inform intervention strategies as well as attract funding from the WHO and other international agencies (e.g. UNICEF, UNDP, Swedish International Development Authority-SIDA and USAID) to complement local governmental efforts. Even though serum thyroid stimulating hormone, serum thyroglobulin and serum T4 measurements offer alternative ways of assessing iodine status, these alternative approaches are invasive, not feasible in population scale monitoring studies and their usefulness in certain population groups have been questioned [23,24].

Example of such fruitful governmental-international body partnership is the GAIN (Global Alliance for Improved Nutrition)-UNICEF USI Partnership Project in Indonesia that aims to make iodised salt readily available to Indonesians [25]. Moreover, Ecuador provides a clear argument for the effectiveness of joint enterprise between governments and international partners in the fight against IDDs. Although Ecuador had a high prevalence of goitre and cretinism in the 1950s, it has practically eliminated these disorders through such vibrant partnership between the Ecuadorian and Belgian governments [14]. Additionally, portion of these nationally devoted budgets must also be used to build the capacity of small scale salt manufacturers who produce cheap non-iodised salt alternatives especially in countries such as India, Ghana, and Indonesia [25-27]. As these small-scale salt producers do not have the capacity to iodise their salt, these funds should aim to build their infrastructural capacity so as to realise the goal of universal iodisation of all salts on the markets. A recent study in Ghana and Sierra Leone respectively found that 33% and 20% households were not using iodised salt [19,20]. Taking measures to prevent availability of non-iodised salt on the market is crucial as a recent adjusted logistic regression analysis demonstrated a 3% higher odd of stunted growth with non-availability of iodised salt [28]. Empowering the small-scale salt producers in this way would help obviate the lack of adherence to the ban on non-iodised salt usage in certain countries like India [26,29]. Moreover, food fortification programmes must also be given serious consideration by these developing countries to increase the odds of eliminating iodine deficiency scourge. Bread, cooking oils, and rice are staple foods in most of these communities have been successfully iodised in certain countries [14,30]. Countries like Australia and New Zealand have implemented mandatory food fortification with iodine that have significantly improved the iodine levels of people in these countries [31]. In the United States also, cow milk [32], and chicken eggs [33] contain iodine that may

supplement what may be ingested through iodised salt. Therefore, efforts must be made to iodise foods such as bread, cooking oils, rice and even water in these developing countries to offer a multi-faceted approach so as to maximize the chances of eliminating IDDs. A recent study demonstrated that fortification of vegetable oil with other micronutrient such as vitamin A and iron improved the nutritional status in the study participants [34]. An additional advantage offered by fortification of these other vehicles is the prevention of hypertension associated with increased salt intake.

## Conclusion

Eradication of IDD the USI initiative is attainable only through a concerted effort by governments, international funding bodies, and salt producers coupled with sustained education of consumers to motivate them to utilize these iodised salts.

## References

1. Morreale de Escobar G, Obregon MJ, Escobar del Rey F (2004) Role of thyroid hormone during early brain development. *Eur J Endocrinol* 151(suppl 3): U25-37.
2. Dong J, Yin H, Liu W, Wang P, Jiang Y, et al. (2005) Congenital iodine deficiency and hypothyroidism impair LTP and decrease C-fos and C-jun expression in rat hippocampus. *Neurotoxicology* 26(3): 417-426.
3. Glinoe D, Delange F (2000) The potential repercussions of maternal, fetal and neonatal hypothyroxinemia on the progeny. *Thyroid* 10(10): 871-887.
4. (1990) WHO, Prevention and control of iodine deficiency disorders, Geneva, Switzerland.
5. Thilly, Delange F, Golstein-Golaire J, Ermans AM (1973) Endemic goiter prevention by iodized oil: a reassessment. *J Clin Endocrinol Metab* 36(6): 1196-1204.
6. Pretell, Moncloa F, Salinas R, Kawano A, Guerra Garcia R, et al. (1969) Prophylaxis and treatment of endemic goiter in Peru with iodized oil. *J Clin Endocrinol Metab* 29(12): 1586-1595.
7. (2004) WHO, Iodine status Worldwide, Geneva, Switzerland, p. 1-58.
8. UNICEF (2014) Global Database: Household Consumption of adequately Iodised salt, D.a.A. section, WHO.
9. Diosady LL, A J, Mannar MG, (1998) Fitzgerald S Stability of iodine in iodized salt used for correction of iodine-deficiency disorders. *Food Nutr Bull* 19: 240-250.
10. (1996) WHO/UNICEF/ICCIDD/NUT, Recommended iodine levels in salt and guidelines for monitoring their adequacy and effectiveness, Geneva, Switzerland, p. 9.
11. Diosady LL, J O Alberti, S FitzGerald, M G Venkatesh Manna, (1999) Field tests for iodate in salt. *Food Nutr Bull* 20(2): 208-214.
12. Pandav CS, Arora NK, Krishnan A, Sankar R, Pandav S, et al.(2000) Validation of spot-testing kits to determine iodine content of salt. *Bull World Health Organ* 78(8): 975-980.
13. Gorstein J, Van Der Haar F, Codling K, Houston R, Knowles J, et al. (2016) Performance of rapid test kits to assess household coverage of iodized salt. *Public Health Nutr* 19(15): 2712-2724.

14. Christopher P Howson, Eileen T Kennedy, Abraham Horwitz, (1998) Prevention of Micronutrient Deficiencies: Tools for Policymakers and Public Health Workers. National Academies Press, USA.
15. Diosady LL, Alberti JO, Ramcharan K, Mannar MG (2002) Iodine stability in salt double-fortified with iron and iodine. *Food Nutr Bull* 23(2): 196-207.
16. Dasgupta PK, Liu Y, Dyke JV (2008) Iodine nutrition: iodine content of iodized salt in the United States. *Environ Sci Technol* 42(4): 1315-1323.
17. Nkowane AM, Boualam L, Haithami S, El Sayed el TA, Mutambo H, et al. (2009) The role of nurses and midwives in polio eradication and measles control activities: a survey in Sudan and Zambia. *Hum Resour Health* 7: 78.
18. Aylward RB, Jennnifer L (2005) Polio eradication: mobilizing the human resources. *Bulletin of the World Health Organization*: Geneva, Switzerland, pp. 268-273.
19. Rohner F, Wirth JP, Woodruff BA, Chiwile F, Yankson H, et al. (2016) Iodine Status of Women of Reproductive Age in Sierra Leone and Its Association with Household Coverage with Adequately Iodized Salt. *Nutrients* 8(2): 74.
20. Simpong DL, Adu P, Bashiru Morna MT, Yeboah FA, et al. (2016) Assessment of iodine status among pregnant women in a rural community in Ghana - a cross sectional study. *Arch Public Health* 74: 8.
21. Juan W, Trumbo PR, Spungen JH, Dwyer JT, Carriquiry AL, et al. (2016) Comparison of 2 methods for estimating the prevalences of inadequate and excessive iodine intakes. *Am J Clin Nutr* 104(Suppl 3): 888S-897S.
22. (1994) WHO/UNICEF/ICCIDD/NUT, Indicators for assessing iodine deficiency disorders and their control through salt iodination, Geneva, Switzerland.
23. Ristic Medic D, Piskackova Z, Hooper L, Ruprich J, Casgrain A, et al. (2009) Methods of assessment of iodine status in humans: a systematic review. *Am J Clin Nutr* 89(6): 2052S-2069S.
24. Offie P Soldin, Rochelle E Tractenberg, John C Pezzullo (2005) Do Thyroxine and Thyroid-Stimulating Hormone Levels Reflect Urinary Iodine Concentrations?. *Therapeutic Drug Monitoring* 27(2): 178-185.
25. (2008) GAIN, Universal Salt Iodization: "GAIN-UNICEF USI Partnership Project" in Indonesia.
26. Kalra SKB, Sawhney K (2013) Usage of non-iodized salt in North West India. *Thyroid Res Pract* 10: 12-14.
27. Buxton C, Baguune B (2012) Knowledge and practices of people in Bia District, Ghana, with regard to iodine deficiency disorders and intake of iodized salt. *Arch Public Health* 70(1): 5.
28. Kramer M, Roland Kupka, SV Subramanian, Sebastian Vollmer (2016) Association between household unavailability of iodized salt and child growth: evidence from 89 demographic and health surveys. *Am J Clin Nutr*.
29. Priya R, A Kotwal, I Qadeer (2009) Toward an ecosocial epidemiological approach to goiter and other iodine deficiency disorders: a case study of India's technocratic program for universal iodization of salt. *Int J Health Serv* 39(2): 343-362.
30. Kevany J, Fierro-Benitez R, Pretell EA, Stanbury JB, (1969) Prophylaxis and treatment of endemic goiter with iodized oil in rural Ecuador and Peru. *Am J Clin Nutr* 22(12): 1597-1607.
31. (2011) Mandatory folic acid and iodine fortification in Australia and New Zealand: supplement to the baseline report for monitoring, Australian Institute of Health and Welfare, Canberra.
32. Flachowsky G, Franke K, Meyer U, Leiterer M, Schöne F (2014) Influencing factors on iodine content of cow milk. *Eur J Nutr* 53(2): 351-365.
33. Rottger AS, Ingrid Halle, Hubertus Wagner, Gerhard Breves, Dänicke S, et al. (2012) The effects of iodine level and source on iodine carry-over in eggs and body tissues of laying hens. *Arch Anim Nutr* 66(5): 385-401.
34. Rohner F, Raso G, Aké-Tano SO, Tschannen AB, Mascie-Taylor CG, et al. (2016) The Effects of an Oil and Wheat Flour Fortification Program on Pre-School Children and Women of Reproductive Age Living in Cote d'Ivoire, a Malaria-Endemic Area. *Nutrients* 8(3): 148.