

# Exercise, beyond chronic disease prevention- pathways for brain plasticity

## Abstract

Physical inactivity and sedentary lifestyle, can be considered as important health-related challenges which can affect public health. Decrease in level of physical activity may increase the risk of chronic diseases including over weight/obesity, diabetes, cardiovascular disease, cancers and neurodegenerative diseases. Regular physical activity and exercise, regardless the type, intensity, duration and frequency, can bring many health benefits including increase in cardio-respiratory fitness, physical fitness, decrease in risk of chronic diseases and brain health. The purpose of this study, was to assess the effect of exercise on brain health and brain plasticity along with other beneficial effects including weight management and disease prevention and highlight the relation between body organs with emphasize of cardiovascular system and central nervous system.

**Keywords:** exercise, brain plasticity, cardiovascular system, health

Volume 12 Issue 5 - 2022

Kimia Moiniafshari,<sup>1</sup> Farshad kalantari,<sup>2</sup> Hafez Behzadi Nezhad<sup>3</sup>

<sup>1</sup>Department of Land, Environment, Agriculture and Forestry, University of Padua, Padua, Italy

<sup>2</sup>Department of physical education and sport sciences, Islamic Azad University, East Tehran Branch, Tehran, Iran

<sup>3</sup>Department of physical education and sport sciences, University of Tehran, Tehran, Iran

**Correspondence:** Kimia Moiniafshari, Department of Land, Environment, Agriculture and Forestry, University of Padua, Padua, Italy, Email [kimia.moiniafsha@gmail.com](mailto:kimia.moiniafsha@gmail.com)

**Received:** December 03, 2022 | **Published:** December 09, 2022

## Introduction

Recent scientific research have been suggested that physical activity and regular exercise, may improve brain function which can be considered as a preventative strategy for neurodegenerative diseases. The underlying mechanism which exercise can affect the brain, it not clear yet, although some evidences have suggested that an ingredient known as clasterin, can increase the effect of exercise on hippocampus, cell production, memory improvement and decrease in inflammation in brain.<sup>1</sup> Physical activity and exercise can affect different regions in brain, especially hippocampus, which may affect memory, learning and neurodegenerative diseases prevention. Endurance exercise such as running, may increase neurogenesis, synaptic plasticity, increase in neutrophin and neurotransmitters level, angiogenesis and decrease in neural inflammation along with increase in hippocampus mass. It has been indicated that during exercise, adipose tissue and liver, produce Exerkin which can help homeostasis, neural function, synaptic and brain plasticity due to increase in blood flow.<sup>2</sup>

## Physical exercise- a reverse epigenetic clock

Sedentary lifestyle and physical inactivity are common challenges in industrialized societies which can be a leading cause for cardiovascular diseases, metabolic impairment, cancer and neurodegenerative diseases. During recent years, regular exercise, regardless to its type, has been introduced as a component of healthy lifestyle, as it can improve physical function, lipid profile and metabolic status beside its beneficial effects on brain.<sup>3,4,5</sup>

## How can exercise affect brain plasticity?

**Telomers:**-Studies on brain and factors which can affect on brain health, have indicated that exercise can be considered as an external indicator which can inhibit brain aging. Exercise can induce neurovascularization and BDNF (Brain-Derived-Neurotrophic Factor) production which delay the brain aging. Exercise can increase telomerase activity, telomere length and its stability. Moreover, circulating BDNF, fibronectin-3, Irisin and ketone bodies are other factors which may be affected by exercise and can affect various functions on central nervous system which can lead to brain plasticity and functional improvements.<sup>6,7</sup> BDNF is an important indicator for

brain function and synaptic transformation which can be induce by exercise and mRNA expression. As a result, it can lead to neural and brain plasticity along with brain functional and structural adaptation in response to environmental stimuli.<sup>8</sup>

## Why brain plasticity is important?

During past decades, the effect of exercise on brain plasticity has gained a lot of attention. Studies have suggested that lifestyle can improve brain plasticity, brain structure and its function which can decrease the risk of neural-cognitive impairments. It seems that aerobic exercise can affect frontal region of hippocampus by increasing gray matter.<sup>9</sup> Studies using fMRI (functional magnetic resonance imaging) on brain, has shown that brain plasticity will increase in the mentioned region and code-processing will be better along with improvement in brain function. This result, has made researcher to investigate the relation between exercise and brain plasticity. The results from the studies have indicated that there is a relation between cardio-vascular endurance and brain plasticity, which means that individuals with better aerobic endurance, have better brain plasticity and motor networking which can be achieved even by single bout of training. Researcher have then started to assess the effect of resistance training on brain function and brain plasticity which indicated the similar results.<sup>10</sup>

## Molecular pathways for the effect of exercise on brain health

### How can exercise improve brain plasticity?

Regular exercise can improve brain function, synaptic flexibility and BDNF especially in hippocampus. This result has also been achieved by single bouts of training. One suggested mechanism for the effect of exercise on brain plasticity, is due to increase in the level of Endocannabinoids and Andamide along with increase in BDNF level.<sup>11-14</sup> Exercise can affect cell growth and differentiation, which increase the demand for nutrients. In response to cell growth and need for nutrients, vascularization will occur which can provide nutrients which is not only for hippocampus, but also for motor cortex and frontal lobe origin that ultimately lead to learning and brain plasticity.<sup>15,16</sup> Furthermore, exercise can increase the number of synapses in hippocampus and BDNF gene expression.<sup>17</sup>

### Exercise effect on cognitive function

Increase in physical activity status and recruiting regular exercise, may improve cardio-respiratory fitness and capacity, help cognitive function and improve brain plasticity. Studies have suggested that this may be more important in older ages. Regular exercise for 3-6 months can improve cognitive function. Exercise with moderate intensity and/or stretch training for 6 month, 3 sessions/week, 30-45 minutes/ session can bring similar effect on brain plasticity and cognitive function.<sup>18,19</sup> This can be as a result of brain ability to be flexible and modifiable. As aging occurs, the brain tends to become smaller which is considered as atrophy. After the age of 55, both frontal and hippocampus region, will become smaller by 1-2% each year which can lead to decrease in cognitive function. It has been indicated that regular exercise can be an effective and preventative strategy for prevention of gray matter in frontal and temporal regions which can be as a result of aerobic and cardio-vascular capacity improvement.<sup>20,21</sup> In conclusion, physical activity and regular exercise, can bring beneficial health outcomes, not only for obesity, metabolic disorder, cardiovascular disease and other health complications, but can also improve brain plasticity due to the connection of several systems including skeletal muscle (exercise increase the demand for nutrients), cellular mechanisms (cell signaling and cell cascade), cardiovascular system (angiogenesis, vascularization) and central nervous system (neurogenesis, neural plasticity and brain plasticity).

### Acknowledgements

None

### Conflicts of interest

The author declares no conflicts of interest.

### References

- De Miguel Z, Khoury N, Lehallier B, et al. Exercise plasma boosts memory and dampens brain inflammation via clusterin. *Nature*. 2021;600(7889):494–499.
- Rai M, Demontis F. Muscle-to-brain signaling via myokines and myometabolites. *Brain Plasticity*. 2021;8(1):43–63.
- Bangsbo J, Blackwell J, Boraxbekk CJ, et al. Copenhagen Consensus statement 2019: physical activity and ageing. *Br J Sports Med*. 2019;15(14):856–858.
- Amaro-Gahete FJ, De-la OA, Jurado-Fasoli L, et al. Metabolic rate in sedentary adults, following different exercise training interventions: the FIT-AGEING randomized controlled trial. *Clin Nutr*. 2020;39(11):3230–3240.
- Valenzuela PL, Castillo Garcia A, Morales JS, et al. Exercise benefits on Alzheimer's disease: State-of-the-science. *Ageing Res Rev*. 2020;62:101108.
- Lu AT, Xue L, Salfati EL, et al. GWAS of epigenetic aging rates in blood reveals a critical role for TERT. *Nat Commun*. 2018;9(1):387.
- Jachim SK, Sakamoto AE, Zhang X, et al. Harnessing the effects of endurance exercise to optimize cognitive health: fundamental insights from Dr. Mark P. Mattson. *Ageing Res Rev*. 2020;64:101147.
- Xu M, Zhu J, Liu, et al. Roles of physical exercise in neurodegeneration: reversal of epigenetic clock. *Translational Neurodegeneration*, 2021;10(1):1–15.
- Voss MW, Heo S, Prakash RS, et al. The influence of aerobic fitness on cerebral white matter integrity and cognitive function in older adults: Results of a one-year exercise intervention. *Hum Brain Mapp*. 2013;34(11):2972–2985.
- Best JR, Chiu BK, Hsu CL, et al. Long-term effects of resistance exercise training on cognition and brain volume in older women: Results from a randomized controlled trial. *Neuropsychol Soc*. 2015;21(10):745–756.
- Sorrells S F. Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. *Nature* 555. 2018; 377–381.
- Izquierdo I, Bevilacqua LR, Rossato JI, et al. The connection between the hippocampal and the striatal memory systems of the brain: a review of recent findings. *Neurotox Res*. 2006;10(2):113–121.
- Creer D J, Romberg C, Saksida L M, et al. Running enhances spatial pattern separation in mice. *Proc Natl Acad Sci U S A*. 2010;107(5):588–597.
- Chen HI, Lin LC, Liu YF, et al. Treadmill exercise enhances passive avoidance learning in rats: the role of down-regulated serotonin system in the limbic system. *Neurobiol Learn Mem*. 2008;89(4):489–496.
- Creer DJ, Romberg C, Saksida LM, et al. Running enhances spatial pattern separation in mice. *Proc Natl Acad Sci U S A*. 2010;107(5):2367–2372.
- Farmer J, Zhao X, van Praag H, et al. Effects of voluntary exercise on synaptic plasticity and gene expression in the dentate gyrus of adult male Sprague-dawley rats in vivo. *Neuroscience*. 2004;124(1):71–79.
- Vivar C, Potter MC, van Praag H. All about running: Synaptic plasticity, growth factors and adult hippocampal neurogenesis. *Curr Top Behav Neurosci*. 2013;15:189–210.
- Colcombe S, Kramer AF. Fitness effects on the cognitive function of older adults: a meta-analytic study. *Psychol Sci*. 2003;14(2):125–130.
- Kramer AF, Hahn S, Cohen NJ, et al. Ageing, fitness, and neurocognitive function. *Nature*. 1999;400(6743):418–419.
- Colcombe SJ, Erickson KI, Raz N, et al. Aerobic fitness reduces brain tissue loss in aging humans. *J Gerontol A Biol Sci Med Sci*. 2003;58(2):176–180.
- Gordon BA, Rykhlevskaia EI, Brumback CR, et al. Neuroanatomical correlates of aging, cardiopulmonary fitness level, and education. *Psychophysiology*. 2008;45(5):825–838.