

# Influence of biochar on soil nutrient transformations, nutrient leaching, and crop yield

## Abstract

Biochar, a solid carbon rich material and by-product of pyrolysis, has been identified as an amendment to improve soil fertility as well as sequester carbon (C). A growing number of studies have been conducted to test the effect of biochar in soil environment within the past ten years requiring frequent updated reviews and mini-reviews to summarize this rapidly growing body of literature. In this paper, we will summarize and review possible mechanisms of biochar effects on soil nitrogen (N) and phosphorus (P) transformations, nutrient leaching, and crop yield, providing an outlook of biochar implementation in soil environment for future research.

**Keywords:** biochar, soil environment; nutrient transformations, nutrient leaching, crop yield

Volume 4 Issue 5 - 2016

**Si Gao, Thomas H DeLuca**

School of Environmental and Forest Sciences, University of Washington, Seattle, WA, USA

**Correspondence:** Thomas H DeLuca, Professor and Director of School of Environmental and Forest Sciences, University of Washington, Seattle, WA, USA, Email [deluca@uw.edu](mailto:deluca@uw.edu)

**Received:** August 13, 2016 | **Published:** September 26, 2016

## Introduction

Utilization of wood residues to generate energy and byproduct biochar has been identified as a carbon (C) negative solution for agriculture and forest management;<sup>1</sup> however, the rapidly growing body of literature on this topic requires frequent summarization of recent studies to allow researchers to stay current on the topic. Biochar is a C-rich, recalcitrant solid material that is generated from the pyrolysis or thermochemical decomposition of organic material in an oxygen limited environment under controlled conditions. Biochar can serve as an effective soil C sink as it has high proportion of recalcitrant C with thousands of years of stability,<sup>2</sup> thus there is an intense interest in using biochar as a means to mitigate climate change by offsetting C emissions.<sup>3</sup> Besides, the application of biochar to soils has been shown many agronomic benefits in many cases. The physicochemical properties of biochar are key to understanding the function of biochar in soil. The highly porous structure and large surface area of biochar may offer appropriate habitat for beneficial microorganisms to flourish; other physico-chemical properties such as high ion-exchange capacity can also impact a number of processes in the soil N cycle associated with enhanced soil fertility.<sup>4</sup> Improvements in soil fertility by biochar addition have also led to increased crop yield and productivity, the magnitude of response varies with biochar feed stocks, biochar activation or inoculation process, application rates, crop species, soil types and other soil inputs, as well as combination of these factors.<sup>5</sup>

According to Web of Science database, around 2,000 papers has been published based on “biochar and soil” topic since 2012, the overwhelming focus of studies on biochar effects and soil processes has been experimental reports, relatively few new manuscripts have examined potential mechanisms of biochar effects in soil environment, particularly with new studies since 2012. Therefore, a systematic and comprehension review of biochar effect and its mechanisms in soil environment is still a needed area of research currently. Herein, the purpose of this review paper is to summarize and evaluate the mechanisms of biochar effect on

i. soil N and P transformations;

ii. nutrient leaching; and

iii. crop yield.

## Biochar and soil nutrient transformations

### Nitrogen

Nitrogen is the most commonly limiting nutrient in temperature agroecosystems. Most agricultural plants primarily take up inorganic N, which comes from the conversion of organic N to inorganic N (mineralization), although a few crop species were observed to directly take up organic N for energy and growth.<sup>6,7</sup> Biochar has been widely reported to influence N cycling. Several primary N transformation processes associated with biochar addition (N<sub>2</sub> fixation, mineralization, immobilization, gaseous N emissions (i.e. denitrification, ammonia volatilization) are discussed below.

**Nitrogen fixation:** Biological N fixation (BNF) is the primary natural input of N to terrestrial ecosystems, and it plays an essential role in the N cycles of agricultural system.<sup>8</sup> The process of BNF is conducted by bacteria that are either free-living associative, or symbiotic living in an obligate arrangement with host plants (e.g. legumes) or fungal partners (e.g. lichen).<sup>9</sup> In recent years, a couple of agronomic studies have reported that biochar had the capacity to influence BNF in leguminous plants,<sup>10-15</sup> however, mechanisms remain unclear. A possible mechanism related to the biochar-associated increased N<sub>2</sub> fixation could be the effect of nutrient availability. Rondon et al. (2007) conducted a short-term study investigating biochar effect on the BNF of common beans (*Phaseolus vulgaris*) and showed a significant increase in BNF after biochar addition compared to the control.<sup>16</sup> They suggested that the positive result could be attributed to the observed greater availability of trace metals brought by biochar, such as molybdenum (Mo), which is a constituent of the Mo-Fe protein nitrogenase, that can stimulate nodulation.<sup>16</sup> In addition to trace metals, it is also likely that the enhanced BNF is correlated with higher macro- or micro-nutrient availability, such as K,<sup>10</sup> P,<sup>17</sup> Ca and Mg,<sup>18,19</sup> Fe and Mn.<sup>20</sup> However, inhibitory effect has been also observed, since

- i. nodulation is reported more likely to happen under the addition of nutrient-rich biochar<sup>21–23</sup>
- ii. The adsorption of soil signaling compounds to biochar.

Nodule formation in leguminous plants is initiated by the release of signaling compounds (e.g. flavonoid).<sup>24</sup> Gundale et al.,<sup>25</sup> indicated that such polyphenolic compounds could be absorbed by biochar, leading to a reduction of nodulation process. However, it is also important to note that nodule numbers may not represent the activity of N fixation, as Quilliam et al.,<sup>12</sup> found a reduced numbers of nodules, but the mass of nodules and nitrogenase activity were increased.

Compared to the symbiotic N-fixing bacteria that colonize leguminous plants, only a few studies have been conducted to examine the effect of biochar on free-living N fixation bacteria. DeLuca et al. (2015) indicated that one methodology issue might influence the accuracy of the interpretation of N<sub>2</sub> fixation activity, since nitrogenase activity is commonly measured using the acetylene reduction assay, but biochar itself could release ethylene when applied to soil.<sup>26,27</sup> Other than this, biochar could enhance activity of free-living N-fixing bacteria by influencing systematic N availability. Similar to post-fire BNF process, a decrease of N availability through N immobilization could possibly lead to the stimulation of BNF process.<sup>28–30</sup>

**Nitrogen mineralization:** Nitrogen mineralization is defined as the process by which organic N is converted to inorganic forms (primarily  $\text{NH}_4^+\text{-N}$  and  $\text{NO}_3^-\text{-N}$ ). The conversion of organic-N to  $\text{NH}_4^+\text{-N}$  is defined as ammonification. The conversion of  $\text{NH}_4^+\text{-N}$  or organic-N to  $\text{NO}_3^-\text{-N}$  by autotrophic bacteria, archaea or certain fungi is defined as nitrification. Many studies in temperate and boreal forest soils have shown an increased net nitrification rates in forest soils by biochar additions; however, few studies had found out such results in agricultural systems that already accommodate an active nitrifying community<sup>16,27,31</sup> (Table 1).

Ammonification is a key component of the N cycle in agroecosystems and is of unique importance in organic farming systems. This process is driven by a broad consortium of organisms that are capable of enzymatic denaturation of proteins and the removal of amide groups from organic compounds (e.g. amino acids and amino sugars). It is typically measured by extracting  $\text{NH}_4^+\text{-N}$  from soil at different points in time using a high concentration salt solution, typically potassium chloride (KCl), but can be more completely assessed using a combination of ionic resins, isotopic methods, and incubations.<sup>7</sup> The  $\text{NH}_4^+\text{-N}$  present in KCl extracts is used to represent the N mineralized or ammonified from the organic N pool over a given period of time that is free in the soil solution or on cation exchange sites. Generally, the capacity of biochar to hold  $\text{NH}_4^+\text{-N}$  depends on the cation exchange capacity of the biochar. Therefore,  $\text{NH}_4^+\text{-N}$  extracted as a measure of N mineralization may actually represent the cation exchange capacity of biochar, and vice versa.

Studies have shown either an increase, decrease or no change in N mineralization with biochar application to soil (Table 1). Xu et al.<sup>32</sup> recently described an increase of net N mineralization under biochar application in a soil column study. Similarly, but in a field study, Pereira et al.<sup>33</sup> documented an increase in N mineralization nearly two times greater than the control after biochar addition in an organically managed lettuce farm. They also showed that the gross N mineralization rate was positively correlated with biochar H/C ratio and suggested that less recalcitrant biochar with high H/C ratios increased mineralization rates, since they are more likely to

decompose and thereby free up N into the mineral pool. Gundale et al.,<sup>34</sup> found enhanced net soil N mineralization rates and soil  $\text{NH}_4^+\text{-N}$  concentrations two growing seasons after wood biochar application to soil in northern Sweden regardless of the soil mixing treatment. They attributed this difference more to the promotion of net N mineralization rather than the ash input from biochar itself (biochar serves as a modest  $\text{NH}_4^+\text{-N}$  source). However, a recent study by Luo et al.,<sup>35</sup> found reduced N mineralization after biochar application to a coastal wetland soil. Similar to Dempster et al.,<sup>35</sup> they indicated that the decrease of N mineralization was due to a higher C:N ratio of the biochar and that N mineralization potential is likely related to the character of the biochar feedstock.<sup>36</sup> A decrease in total net N mineralization was also observed with biochar addition to an Aridisol from Colorado and an Alfisol from Virginia following 18 days of incubation with switchgrass biochar.<sup>37</sup> This decline in mineralization was attributed to a decline in microbial activity due to the presence of chemicals such as ethylene which is a known nitrification inhibitor. In contrast, both Prommer et al.<sup>38</sup> and Ulyett et al.,<sup>39</sup> found no significant change in N mineralization with low N feedstock biochar application. In summary, these studies suggest that the biochar feedstock, conditions of biochar formation, time since application, capacity of biochar to adsorb  $\text{NH}_4^+$ , and soil type are all factors that needed to be considered in assessing soil N mineralization response to biochar.

**Nitrogen immobilization:** Nitrogen immobilization is defined as the conversion of inorganic N into organic N via microbial uptake and formation of amino acid N. Whether N mineralization or immobilization occurs with organic amendments to soil depends on the C/N ratio of the amendment, if the C/N ratio is high enough (generally more than 25:1), then the N tends to be immobilized.<sup>40</sup> Biochar generated from wood or N-limited feedstock generally has a high C/N ratio, whereas biochar generated from N-rich feedstock (such as agriculture waste) could serve as N source.<sup>41</sup> It is therefore uncertain whether biochar provide enough C to stimulate N immobilization.

Biochar studies have found variable results in terms of N immobilization. Bruun et al.,<sup>42</sup> indicated that application of incompletely pyrolyzed biomass (fast pyrolysis at low temperature) may cause immobilization of soil N, as more N is needed by the developing microorganisms than is provided by the substrate, in other words, low-temperature biochar contain more bioavailable C or surface functional groups that can serve as microbial substrates.<sup>30,43</sup> Sigua et al.,<sup>44</sup> conducted a field study using switch grass biochar and found that biochar additions to soil increased N immobilization and decreased in total inorganic N in soils due to the wide C/N ratio of switch grass. These researchers also observed a significant increase in cumulative and net  $\text{CO}_2$  flux implying biochar simulated switchgrass mineralization and accelerated decomposition of resident soil C.<sup>44,45</sup> Similar observations such as increased respiration rates have been recently reported;<sup>46</sup> however, Jones et al.<sup>47</sup> suggested that increased  $\text{CO}_2$  evolution immediately after biochar addition partially originates from the emission of inorganic C within biochar itself. Therefore, the influence of biochar on N immobilization needs to be further studied with focus on the bioavailable C and direct effects of biochar on microbial activity.

**Gaseous N emissions:** Nitrogen losses from the soil ecosystem occur as a result of leaching, denitrification, volatilization, crop removals, soil erosion and runoff. Among these mechanism, denitrification and  $\text{NH}_3$  volatilization are two primary processes of gaseous N emissions. Denitrification is the process by which bacteria convert nitrate to N

gases that are lost to the atmosphere ( $\text{NO}_3^- \rightarrow \text{NO}_2 \rightarrow \text{N}_2\text{O} \rightarrow \text{N}_2$ ). Many studies have focused on  $\text{N}_2\text{O}$ , because of its importance as a greenhouse gas.<sup>48</sup> Biochar has been reported to influence  $\text{N}_2\text{O}$  flux in many studies (Table 1). Case et al.,<sup>49</sup> reported that biochar suppressed cumulative soil  $\text{N}_2\text{O}$  production by 91% in near-saturated, fertilized soils in a field study. Another recent field study conducted by Ameloot et al.<sup>50</sup> also observed a 50-90% reduction in  $\text{N}_2\text{O}$  emissions seven months after biochar application to a loam, implying that the biochar exerts an indirect physical control over soil denitrification several months after incorporation. Harter et al.,<sup>51</sup> illustrated that biochar addition to soil enhanced microbial nitrous oxide reduction with enhanced transcript copy numbers of the *nosZ*-encoded bacterial  $\text{N}_2\text{O}$  reductase, similar to other studies.<sup>52,53</sup> A meta-analysis done by Cayuela et al.,<sup>54</sup> reported that biochar application to soils reduced soil  $\text{N}_2\text{O}$  emissions by 54% in laboratory and field studies, across 30 studies and 261 experimental treatments during 2007 to 2013. Several explanations and mechanisms were generated to explain the decreased  $\text{N}_2\text{O}$  emissions with biochar addition to soil:

- i. biochar elevates soil pH slightly creating an environment where  $\text{N}_2\text{O}$  reductase activity is more readily promoted;<sup>55,56</sup>
- ii. Enhanced soil aeration inhibiting denitrification due to more oxygen being present;<sup>49,57,58</sup>
- iii. A shortage of available C due to the adsorption of labile soil organic matter (SOM) compounds on biochar may decrease the denitrification potential and lower  $\text{N}_2\text{O}$  emission rates;<sup>56</sup>
- iv. A reduction in the availability of inorganic-N to denitrifying bacteria and archaea thereby reducing denitrification potential.<sup>4,53</sup>

Ammonium volatilization is another process of gaseous N loss to the atmosphere. It is well known that  $\text{NH}_3$  volatilization can be enhanced in soils with a higher pH.<sup>6</sup> It is also been reported that biochar with residual ash can act as a liming agent that can increase soil pH.<sup>59</sup> However, studies have illustrated that the pH increase with biochar is usually not high enough to enhance  $\text{NH}_3$  volatilization.<sup>27</sup> A recent study by Dougherty et al.,<sup>60</sup> showed that  $\text{NH}_3$  volatilization was significantly reduced with the addition of Douglas-fir chip produced biochar, mostly due to the  $\text{NH}_3$  adsorption at oxygen-containing surface functional group or biochar micro pores.<sup>60,61</sup> Table 1 provides a summary of findings on  $\text{NH}_3$  emissions with biochar additions. Taghizadeh-Toosi et al.<sup>62</sup> found a 45% reduction of  $\text{NH}_3$  volatilization after addition of wood-derived biochar; Doydora et al.,<sup>63</sup> found a 56-63% reduction of  $\text{NH}_3$  loss using poultry litter biochar. Studies have also illustrated that biochar could induce ammonium immobilization and nitrification that can reduce  $\text{NH}_3$  volatilization potential.<sup>61,64</sup> Further *in-situ* field trial and adsorption or desorption studies are needed to verify these results and fully understand the dynamics of  $\text{NH}_3$  adsorption and release.

## Phosphorus

Phosphorus is another plant macro-nutrient that is often co-limiting along with N in agricultural systems. Phosphorus exists in soils in organic and inorganic forms. P is reported almost inaccessible to plants in the organic form, thus need to be mineralized into inorganic P (mostly as  $\text{H}_2\text{PO}_4^-$  and  $\text{HPO}_4^{2-}$ ) prior to plants uptake [65]. Inorganic P is negatively charged in most soils, therefore it tends to react readily with positively charged ions to form mineral precipitates such as Ca-P, or strongly sorbed to the mineral phase (e.g. on Fe and Al oxy-hydroxide surfaces) thus will reduce the solubility of P.<sup>27</sup> Until

now, biochar is reported to alter soil available P by three primary mechanisms:

- i. By acting as a P source providing available P for soils and plants;
  - ii. By altering P solubility, through the alteration of soil pH, adsorption of specific chelates or formation of specific compounds, and P solubilizing bacteria, etc.; and
  - iii. By altering the process of P mineralization and phosphatase enzyme activities. Several recent observations of soil P availability in responses to biochar additions are listed in Table 2.
- Biochar serve as a modest source of P, because the volatilization temperature of P is over 700°C,<sup>66</sup> a temperature in excess of most biochar production temperatures resulting in a residual concentration of P of about 0.4% P in biochar.<sup>67</sup> Wang et al.<sup>68</sup> conducted a study to explore the bioavailability of P in biochars associated with feedstocks (dairy manure and biosolids), results showed that P in feedstock was fully recovered in the biochars by 98% to 119%. Therefore, the proportion of different P pools in biochar and total available P levels are highly dependent on original feedstocks. For instance, wood-derived biochar usually has low P concentrations, whereas manure- or biosolid- derived biochar has relatively higher levels of P that is plant available.<sup>69,70</sup> Pyrolysis can cleave the organic P bonds present in the feedstock, therefore pyrolysis can also lead to the formation of a range of mineral P forms which complexes with Fe, Al, Ca and Mg predominate, biochar therefore contains three pools of P:
- i. Free soluble;
  - ii. Strongly bond to Fe and Al;
  - iii. Organically bound as a residue of the original feedstock.<sup>27</sup>

Biochar can alter soil P solubility through several mechanisms. Biochar can influence P precipitation by altering soil pH and thus the strength of ionic P interactions with  $\text{Al}_3^+$ ,  $\text{Fe}_3^+$ , and  $\text{Ca}_2^+$ ; or by adsorbing organic molecules that normally act as chelates (such as phenolic acids, complex proteins and carbohydrates) of metal ions that otherwise precipitate P.<sup>27,71-73</sup> Hydrophobic or charged biochars are more efficient in adsorbing these organic molecules and creating organo-biochar or organo-mineral-biochar complexes over time, leading to an enhanced P solubility, retention and availability.<sup>74,75</sup> Soil microorganisms are also effective in releasing soil P through solubilization processes.<sup>76</sup> For instance, Suksabye et al. (2016) reported that  $\text{PO}_4^{3-}$  solubilizing bacteria *Pseudomonas aeruginosa* and *Bacillus subtilis* are effective in solubilizing considerable amounts of  $\text{Ca}_3(\text{PO}_4)_2$ .<sup>77</sup> Promoted growth of bacteria that correspond to producing P solubilizing compounds in the presence of biochar could influence inorganic P bioavailability.<sup>78</sup>

Phosphorus in organic forms is released by mineralization process involving soil organisms. Biochar can alter the activity and abundance of these microbes thus P availability. Phosphatase is an enzyme that can hydrolyze compounds of organic P and transform them into different forms of inorganic P, which are assimilated by plants.<sup>79</sup> It is been widely illustrated that biochar can enhance phosphatase activity,<sup>80-82</sup> whereas some studies reported no change.<sup>83,84</sup> However, most of these studies are based on observation of change in uptake or availability and may not actually address mechanisms of increased P availability.

## Biochar and soil nutrient leaching

Nutrient leaching and loss is a significant concern in agricultural systems. Nutrient leaching occurs when mobile nutrients are translocated downward in the soil profile with water percolation

below the rooting zone making the nutrients unavailable for plant uptake Major et al.<sup>85</sup> Biochar has been widely reported to reduce nutrient leaching in agricultural systems. A variety of observations from recent lab and field studies related to the influence of biochar on soil nutrient leaching are listed in Table 3.

**Table 1** Studies on soil microbial N cycle responses to biochar additions

Microbial N Cycle Variables	Observations	Type of Study	Biochar Description	Application Rate	Soil Characteristics	Citations
N mineralization	↓	Lab incubation (46 d)	350°C peanut biochar, sieved under 2mm	0, 1%, 3% (w/w)	Sandy loam	120
N mineralization	↑		Straw residues, wood chips	0, 0.5%, 1%, 2% (w/w)	Paddy soil	121
N mineralization	Mineralization to NO <sub>3</sub> <sup>-</sup> ↓, mineralization to NH <sub>4</sub> <sup>+</sup> no change overall	Field (3 years, Mediterranean barley crop)	Pine ( <i>Pinus pinaster</i> + <i>Pinus radiata</i> ) chip gasifier biochar, 600-900°C	0, 12, 50t ha <sup>-1</sup>	Sandy loam	122
N mineralization	Mineralization to NO <sub>3</sub> <sup>-</sup> ↑	Lab	Poultry litter 400, 600°C, swine manure 400, 600°C	0, 2% (w/w)	Sandy, silt-loam soil	123
N mineralization	↓	Lab	Pine chips and poultry litter at 400°C and 500°C	0, 20t ha <sup>-1</sup>	Luvisols	124
N mineralization	↑	Field (wheat and oilseed rape)	Hardwood trees thinnings, slow pyrolysis 400°C, sieved < 2mm	0, 2% (w/w)	Sandy loam	49
N mineralization	↑	Field (boreal forest)	<i>P. sylvestris</i> , wood and bark	0, 10t ha <sup>-1</sup>	Fine sandy Typic Haplocryod	34
N mineralization	↓	Growth chamber	Switchgrass	0, 25, 50, 100t ha <sup>-1</sup>	Aridisol, Alfisol	37
N mineralization	↑	Field (organic lettuce farm)	Douglas-fir wood pyrolyzed at 410°C; Douglas-fir wood pyrolyzed at 510°C, hogwaste wood pyrolyzed between 600-700°C	0, 10t ha <sup>-1</sup>	Loam	33
N mineralization	↑ at the first 4 days, – after 4 days	Lab	Rye grass, pyrolysis at 450°C	0, 13 mg g <sup>-1</sup>	Cambisol (forest)	125
N mineralization	–	Field (barley and sunflower)	Hardwood-derived biochar (mostly beech), 500°C for 2 h	0, 24, 72t ha <sup>-1</sup>	Sandy to loamy silt	38
N mineralization	– organic, ↓ conventional	Field (organic and conventional)	A mix of sycamore, oak, beech, bird cherry, 600°C 16h, crushed to a diameter of less than 15mm	0, 30, 60t ha <sup>-1</sup>	Sandy loam (Luvisol, Cambisol)	39



Table Continued

Microbial N Cycle Variables	Observations	Type of Study	Biochar Description	Application Rate	Soil Characteristics	Citations
N mineralization	—	Field (maize system)	Maize stover, slow pyrolyzed at 600°C	0, 1, 3, 12, 30 t ha <sup>-1</sup>	Kendaia silt loam and Lima loam	17
N mineralization	↓	Greenhouse	<i>Eucalyptus marginata</i> , Pyrolysis 24h 600°C	0, 5, 25 t ha <sup>-1</sup>	Grey Orthic Tenosol	36
N mineralization	↑	Lab	Swine manure, barley stover, carbonized 600–800°C, digest 30 min 320°C, cooled, filtered, dried	0, 2% (w/w)	Ustisols (under paddy or pasture)	80
N mineralization	—, ↓ at 14 days	Field	Commercial horticultural charcoal (coppiced woodlands: beech, oak, hazel, and birch), pyrolysis 500°C	0, 3, 6 kg m <sup>-2</sup>	Silty loam	126
N mineralization	↓	Lab	Four biochars: douglas fir pellets, douglas fir bark, switchgrass straw, animal digested fiber, all pyrolysis at 600°C	0, 9.8, 19.5, 39.0 t ha <sup>-1</sup>	Sand, silt loam	127
N mineralization	↓	Lab	<i>Macadamia integrifolia</i> , flash pyrolysis, 300–800°C	0, 2.5% (w/w)	Ustic kanhaplohumult	95
N mineralization	↑	Field (Scots pine forest, Sweden)	Activated carbon	1000 kg ha <sup>-1</sup>	Typic or Entic Haplocryods	128
Nitrification	↑	Field (wheat and oilseed rape)	Hardwood trees thinnings, slow pyrolysis 400°C, sieved < 2mm	0, 2% (w/w)	Sandy loam	49
Nitrification	↑ at the first 18 days, — after 4 days	Lab	Rye grass, pyrolysis at 450°C	0, 13 mg g <sup>-1</sup>	Cambisol (forest)	125
Nitrification	↑	Field (barley and sunflower)	Hardwood-derived biochar (mostly beech), 500°C for 2 h	0, 24, 72 t ha <sup>-1</sup>	Sandy to loamy silt	38
Nitrification	↓	Greenhouse	<i>Eucalyptus marginata</i> , Pyrolysis 24h 600°C	0, 5, 25 t ha <sup>-1</sup>	Grey Orthic Tenosol	36
Nitrification	—	Field	Commercial horticultural charcoal (coppiced woodlands: beech, oak, hazel, and birch), pyrolysis 500°C	0, 3, 6 kg m <sup>-2</sup>	Silty loam	126
Nitrification	↑	Lab (used forest soils)	Lab biochar, ponderosa pine wood, homogenized, sieved < 2mm	1000 mg charcoal kg <sup>-1</sup> soil	Sandy loam	31
Nitrification	↑	Lab	Activated carbon	2000 kg ha <sup>-1</sup>	Typic or Entic Haplocryods	129

Table Continued

Microbial N Cycle Variables	Observations	Type of Study	Biochar Description	Application Rate	Soil Characteristics	Citations
N immobilization	NH <sub>4</sub> <sup>+</sup> immobilization ↑, NO <sub>3</sub> <sup>-</sup> immobilization ↓	Field (wheat and oilseed rape)	Hardwood trees thinnings, slow pyrolysis 400°C, sieved < 2mm	0, 2% (w/w)	Sandy loam	49
N immobilization	NO <sub>3</sub> <sup>-</sup> immobilization ↑, NH <sub>4</sub> <sup>+</sup> immobilization –	Field (barley and sunflower)	Hardwood-derived biochar (mostly beech), 500°C for 2 h	0, 24, 72 t ha <sup>-1</sup>	Sandy to loamy silt	38
N immobilization	↑ during the 65 days of incubation	Lab	Wheat straw, 525°C, fast pyrolysis	0, 5% (w/w)	Sandy loam	42
N immobilization	↑	Lab (column study)	Pecan shell biochar	0, 0.5%, 1.0%, 2.0% (w/w)	Loamy sand (fine-loamy, kaolinitic, thermic typic Kandiodults)	130
N <sub>2</sub> O evolution	↓	Lab incubation	Swine manure digestate biochar 350, 700°C, willow wood biochar 350, 700°C	0, 10 t ha <sup>-1</sup>	Loam (Alfisol)	50
N <sub>2</sub> O evolution	↓	Field (wheat and oilseed rape)	Hardwood trees thinnings, slow pyrolysis 400°C, sieved < 2mm	0, 2% (w/w)	Sandy loam	49
N <sub>2</sub> O evolution	↓	Lab	Commercial green waste biochar, 700°C	0, 2%, 10% (w/w)	Loamy sand (calcaric leptosol)	51
N <sub>2</sub> O evolution	↓	Lab (column study)	Commercial wheat straw biochar, 450°C, 4.5 h	0, 30 t ha <sup>-1</sup>	Agricultural soil (silt clay), forest soil (loam)	131
N <sub>2</sub> O evolution	↓	Lab incubation	Oil mallee, wheat chaff, and poultry litter biochars, all produced at 500°C	0, 1% (w/w)	Vertisol (clay), Ferrosol (clay), Calcarosol (sandy clay loam) and Tenosol (sand)	53
N <sub>2</sub> O evolution	↓ (pasture soil + barley stover biochar), ↑ (rice paddy soil + swine manure biochar)	Lab	Swine manure, barley stover, carbonized 600–800°C, digest 30 min 320°C, cooled, filtered, dried	0, 2% (w/w)	Udisols (under paddy or pasture)	80
N <sub>2</sub> O evolution	↑ at first 3 days, – after 3 days	Field	Commercial horticultural charcoal (coppiced woodlands: beech, oak, hazel, and birch), pyrolysis 500°C	0, 3, 6 kg m <sup>-2</sup>	Silty loam	126
N <sub>2</sub> O evolution	↓	Lab	Municipal biosolids	0, 10% (w/w)	Loam	57
NH <sub>3</sub> volatilization	↓	Lab incubation	Poultry litter biochar and Macadamia nut shell biochar	0, 5% (w/w)	Mawson Lakes Technology Park soil, Port Sunny Vale soil, Port Wakefield soil, Mount Lofty soil, and Adelaide Hill soil	61
NH <sub>3</sub> volatilization	↓	Lab incubation	Coconut shell biochar followed by steam activation	0, 1.5%, 3% (w/w)	Silty loam	132

Table Continued

Microbial N Cycle Variables	Observations	Type of Study	Biochar Description	Application Rate	Soil Characteristics	Citations
NH <sub>3</sub> volatilization	↑	Chamber	Commercial Miscanthus giganteus biochar, slow pyrolysis at 600°C	0, 3% (w/w)	Silt-loam, loam soil	133
NH <sub>3</sub> volatilization	↑ (agricultural soil), ↓ (forest soil)	Lab (column study)	Commercial wheat straw biochar, 450°C, 4.5 h	0, 30t ha <sup>-1</sup>	Agricultural soil (silt clay), forest soil (loam)	131
NH <sub>3</sub> volatilization	↑ when under low pH (pH=5), ↓ when under medium pH (pH=7-8)	Lab incubation	Green waste biochar	0, 1%, 5%, 10%, 20% (w/w)	Bauxite residue sand	134
NH <sub>3</sub> volatilization	↓	Lab	Monterey Pine biochar, 300, 350, 500°C, sieved < 2mm	0, 2% (w/w)	Temuka silt loam	62
NH <sub>3</sub> volatilization	↓	Lab incubation (21 d)	Pine chips and peanut hulls biochar, slow pyrolysis 400°C, 1 h	0, 5t ha <sup>-1</sup>	Pasture soil (Cecil)	63

↑ = indicates an increase in occurrence

↓ = indicates a decrease in occurrence

**Table 2** Studies on soil P availability responses to biochar additions

Observations	Type of study	Biochar description	Application rate	Soil characteristics	Citations
Wood derived biochar increased soil citrate extractable P (active inorganic P) by 29% after one growing season across 10 organic farms	Field study (10 organic farms)	Douglas fir, white fir, and western red fir mixture derived biochar	0, 20t ha <sup>-1</sup>	Sandy loam, loamy sand	135
Co-pyrolysis of bone and wood decreased available P	Pot study	Hardwood chips and bone meal derived biochar	0, 4.2, 1.2, 0.9 g kg <sup>-1</sup>	Silty clay	136
Biochar increased Fe-P uptake, with arbuscular mycorrhizas related Fe-P uptake increased by 12%	Greenhouse study	Maple-hickory biochar (450°C for 1h, slow pyrolysis)	0, 7.8t ha <sup>-1</sup>	Weathered tropical soils with eroded surface horizons	137
The concentration of total and available P increased with higher biochar and sewage sludge application rates	Incubation	Wood chip biochar in combination with dried sewage sludge	0, 1%, 5% (w/w)	Cambisol, Rendzina	138
Biochar at 1% application rate showed the highest concentration of water-soluble P across 11 experimental fields	Field study (11 experimental fields)	Wheat residue derived biochar	0, 0.5, 1, 2, 4% (w/w)	Silt loam, clay loam, loam soil	139
Phosphatase activity: 1.5% manure biochar decreased acid phosphatase activity by 18.6% and 34.0% for clay loam and silt loam soil, respectively; increased alkaline phosphatase activity by 28.5% and 95.1% for clay loam and silt loam, respectively	Microcosm incubation	Manure-derived biochar	0, 0.5%, 1.5% (w/w)	Clay loam, silt loam	140
No impact on acid phosphatase activity	Incubation	Bamboo, rice straw derived biochar (over 500°C)	0, 1%, 5% (w/w)	Sandy loam	141

Table Continued

Observations	Type of study	Biochar description	Application rate	Soil characteristics	Citations
Biochar applied at 30 and 45t ha <sup>-1</sup> increased soil alkaline phosphatase activity by 198% and 120%, respectively after the second growing season	Field study	Commercial biochar produced of wheat straw	0, 30, 45 th <sup>-1</sup>	Loamy sand	81
Biochar has no impact on acid phosphatase activity, both 20t ha <sup>-1</sup> and 40t h <sup>-1</sup> biochar increased alkaline phosphatase activity	Field study	Wheat straw derived biochar (between 350-550°C)	0, 20, 40t ha <sup>-1</sup>	Aquept	142
20g kg <sup>-1</sup> biochar application rate showed maximum increases in acid phosphatase activity (32%) and alkaline phosphatase activity (22.8%)	Pot study	<i>Eichorinia</i> derived biochar	0, 1,3,5, 10, 20g kg <sup>-1</sup>	Ustorthents	143
Biochar amended with sewage sludge increased acid phosphatase activity	Pot study	Biochar with sewage sludges added	0, 4% (w/w)	Umbrisol, sandy loam	144

Table 3 Studies on soil nutrient leaching responses to biochar additions

Biochar	Type of study	Soils characteristics	Observations	Citations
Corn stalks, 350°C	Lab	Loam with low SOC level (0.79%)	29% decrease in NO <sup>3-</sup> leaching	145
Sewage sludge, 300°C	Lab	Clay loam (Ultisol)	6.8%, 8.5%, 7.9% decrease in NH <sup>4+</sup> , PO <sub>4</sub> <sup>3-</sup> , K <sup>+</sup> leaching, respectively; 0.2% increase in NO <sup>3-</sup> leaching	146
Sewage sludge, 500°C	Lab	Clay loam (Ultisol)	19.4%, 6.4%, 12.9%, 12.1% decrease in NH <sup>4+</sup> , NO <sup>3-</sup> , PO <sub>4</sub> <sup>3-</sup> , K <sup>+</sup> leaching, respectively	146
Sewage sludge, 700°C	Lab	Clay loam (Ultisol)	35.9%, 9.7%, 23.7%, 23.4% decrease in NH <sup>4+</sup> , NO <sup>3-</sup> , PO <sub>4</sub> <sup>3-</sup> , K <sup>+</sup> leaching, respectively	146
Filtercake biochar, 575°C	Lab	Sandy clay loam	No biochar effect on NO <sup>3-</sup> leaching	147
Acacia whole-tree greenwaste biochar, 550°C	Field	Loamy sand	No significant effect on NO <sup>3-</sup> , K <sup>+</sup> leaching, but significantly increased the concentration (34%) and flux (103%) of PO <sub>4</sub> <sup>3-</sup> leaching	148
Pig manure biochar and wood biochar, 600°C	Lab	Sandy loam	24-26% decrease of NO <sup>3-</sup> leaching, no biochar effect on NH <sup>4+</sup> leaching	149
Commercially produced from mixed feedstock of fruit trees, ~500°C	Field	Silty clay loam	72% decrease in NO <sup>3-</sup> leaching, no effect on NH <sup>4+</sup> leaching	150
Maize stover, 600°C	Field	Aeric Endoaquepts, fine-loamy	82% reduction in NO <sup>3-</sup> leaching at 100% recommended fertilization rate; no effect at 50% fertilization rate	17
Peanut hull, 600°C	Lab	Sandy	34 and 14% reduction in NO <sup>3-</sup> and NH <sup>4+</sup> leaching, respectively; 39% increase in P leaching	151



Table Continued

Biochar	Type of study	Soils characteristics	Observations	Citations
Brazilian pepperwood, 600°C	Lab	Sandy	30 and 35% reduction in NO <sup>3-</sup> and NH <sub>4</sub> <sup>+</sup> leaching; 21% reduction in P leaching	152
Locally produced mixed wood, ~500-700°C	Field	Typic Haplustox clay soil	Leaching varied within the rooting zone: at 1.2 m depth Ca <sup>2+</sup> , Mg <sup>2+</sup> , K <sup>+</sup> , NO <sup>3-</sup> and Sr <sup>2+</sup> leaching decreased by 14, 22, 31, 2 and 14%, respectively, while no biochar effect on NH <sub>4</sub> <sup>+</sup> and P	152
Switchgrass at 250°C	Lab	Xeric Haplocalcids loamy soil	27, 27, and 88% reduction in cumulative leaching of Ca, Mg and NO <sup>3-</sup> , respectively; 47% increase in K leaching; no effect on P leaching	153
Switchgrass at 500°C	Lab	Xeric Haplocalcids loamy soil	67% reduction in cumulative leaching of NO <sup>3-</sup> , 267 and 172% increase in K and P, respectively; no effect on Ca and Mg leaching	153
Switchgrass at 250°C	Lab	Xeric Haplocalcids silty soil	32, 28 and 72% reduction in Ca, Mg and NO <sup>3-</sup> , respectively; no effect on K and P leaching	153
Switchgrass at 500°C	Lab	Xeric Haplocalcids silty soil	10, 11 and 152% increase in Mg, K and P leaching, respectively; 37% reduction in NO <sup>3-</sup> leaching	153
Bagasse at 800°C	Lab	Clay soil	5% reduction in NO <sup>3-</sup> leaching	154
Mixed wood at 475°C	Lab	Silty and sandy soils	No effect on P and NO <sup>3-</sup> leaching	155
Bamboo at 600°C	Lab	Sandy silt	15% reduction in NH <sub>4</sub> <sup>+</sup> leaching at the subsurface 10-20 cm depth	156
Mixed wood at ~550°C	Lab	Typic Hapludolls fine loamy soil	74, 14, 28, 35, and 26% increase in leaching of K, Mg, Zn, Ca, and total N, respectively; no effect on P, Cu, Mn, Na, B and Si leaching	157
Pecan shells at 700°C	Lab	Typic Kandiodults fine loamy soil	206 and 110% increase in K and Na leaching, respectively; 35 and 78% decrease in P and Zn leaching; no effect on Ca, Mg and S leaching	130

In general, biochar could affect soil nutrient leaching by these following mechanisms:

- i. Biochar surface chemistry can lead to the retention of nutrients by cation exchange associated with acidic functional groups formed during oxidation process on biochar surfaces; therefore retain most cations like Ca, Mg, K, and Na. The cation exchange capacity (CEC) of biochar has been considered to be one of the most essential surface chemistry properties that can enhance nutrient retention,<sup>4,58,86</sup> and has been reported to increase with biochar time in soil.<sup>87-89</sup>
- ii. Biochar affects soil solution chemistry and soil physical properties, thus altering nutrient retention. Biochar generally has a higher pH value and is known to be used as a liming agent in many agricultural cases, therefore it can indirectly alter soil nutrients solubility through changes in soil pH.<sup>90</sup> Biochar can also affect soil physical properties such as soil bulk density, water retention, soil structure, aggregate stability, and total porosity,<sup>91</sup> thus nutrient retention. A recent study from Andrenelli et al.,<sup>92</sup> reported a significant increase of soil water retention properties with total water stored in soil pores increased up to 18-25%, and a decrease in soil bulk density after pelletized biochar addition, implying a

nutrient retention potential through reduction of water mobility.<sup>92</sup>

- iii. Soil microbial activity as influenced by biochar can alter soil nutrient retention. Studies have illustrated that biochar have greater potential to lead changes in microbial abundance, community structure and activities.<sup>93,94</sup> Pore spaces within biochar structure could provide suitable habitat for soil microorganisms (bacteria, fungi, protozoa).<sup>12,93</sup> The nutrients and DOC that are desorbed from biochar surface are responsible for the microbial growth, and will lead to alterations of nutrient cycling thus nutrient retention.<sup>26,30,95</sup> Biochar may also induce soil N immobilization to some degree as it is N limited and has a high C:N ratio.<sup>27</sup> A biochar pot experiment on soil bacterial community structure from Anderson et al.,<sup>78</sup> indicated that, the addition of biochar could potentially enhance the growth of organisms that will produce NH<sub>4</sub><sup>+</sup>-N from NO<sub>3</sub><sup>-</sup>-N that can then be adsorbed to biochar.<sup>78</sup> However, further studies related to direct evidences for the impact on microbial processes are needed.

### Biochar effect on plant growth and crop yield

A large number of studies have focused on the influence of biochar on crop yield under both greenhouse and field environments.<sup>2</sup>

The response varies with biochar application rates, crop types, soil types, biochar types including feedstock and pyrolysis conditions, and combinations of these factors.<sup>96</sup> Generally, increasing biochar application rate (within 5–150 t ha<sup>-1</sup>) led to a greater increase in crop production or yields; however, this trend has only been observed in short-term studies (generally within a year),<sup>5</sup> indicating that extra attention should be paid when interpreting these results. From 60 studies that are associated with biochar and crop production, commercial crops such as rice, wheat, maize and soybean all showed significantly higher crop production after biochar additions.<sup>5</sup> However, more field studies for specific species, either short- or long-term, are needed to increase the persuasiveness of this evidence and the accuracy for further reference. Besides, studies conducted on acidic soils or coarse textured soils tended to have greater biochar effect on crop productivity, suggesting liming effect and enhanced soil water storage are the two main reasons improving crop nutrient availability and thus yields.<sup>28,96–98</sup>

Enhanced crop production with biochar additions may be observed as change in plant growth, nutrient uptake and crop yields.<sup>5,99,100</sup> First, biochar can alter soil nutrient pools and availability. Biochar itself can serve as a source of nutrients,<sup>67</sup> and its structure and surface chemistry can enhance the capacity to hold nutrient ions thus increase availability.<sup>101</sup> Second, plant-soil water storage and status may be altered by biochar addition to soil.<sup>102</sup> Biochar can alter the pore size distribution of soil in a long term due to its porous structure,<sup>92,103</sup> thus the addition of biochar may help improve topsoil water holding capacity and storage by the plant delivery of ground water to the topsoil through root hydraulic conductivity.<sup>99,104,105</sup> Although people are arguing that the pores in biochar are too small (usually less than 0.2 μm) for water molecule to percolate or stay,<sup>106</sup> the micro-pores of biochar can still be the source of water vapor that can move within the soil under different temperatures, especially for sandy soils in arid environments.<sup>99</sup>

In addition to soil water storage, biochar itself has been reported to release volatile organic compounds (VOCs) that may promote plant growth.<sup>107–109</sup> However, opposite views also exist on this topic.<sup>110,111</sup> As a common VOC, ethylene produced from biochar may count for another possible reason improving plant growth.<sup>26</sup> Ethylene (C<sub>2</sub>H<sub>4</sub>) is a natural product of plant metabolism,<sup>112</sup> and it has been found to impact the soil microbial and plant processes, for instance, fine root hair growth, increased seed germination, leaf and flower senescence, and increased crop yield in some cases.<sup>26,112–114</sup> Spokas et al. (2010) observed an increase of ethylene production following incorporation of biochar into soils compared to a no biochar control, while the rate of ethylene production varied with biochar production temperature and source materials.<sup>26</sup>

Additionally, biochar can alter plant growth and nutrient uptake by altering the growth of roots and rhizosphere microbial activities.<sup>99</sup> Joseph et al. (2010) indicated that plant roots or root hairs could enter the water-filled macropores or bond onto the biochar surface, causing a wide range of reactions that help the uptake of nutrient.<sup>115</sup> However, the diameter of typical root hairs (5–20 μm) may not match the size of large macro-pores of biochar (wood-derived biochar: 10 μm or more, cellulosic straws-derived biochar: 1–10 μm), limiting the habitat of root hairs in biochar particles.<sup>99</sup> In contrast, fungal hyphae may have more access to biochar, and influence plant nutrient uptake through participation in mycorrhizal functioning.<sup>116</sup>

Somewhat outside of the scope of this review article, but of relevance, biochar has been reported to increase plant protection

against some soil borne plant diseases<sup>117</sup> and induce systemic plant resistance responses to foliar fungal pathogens thereby increasing plant productivity.<sup>118,119</sup> However, further studies are needed to more thoroughly address this topic. Deeper exploration of biochar effect on plants and associated plant pathogens is essential in understanding the potential value of biochar in modifying productivity.<sup>120–136</sup>

## Conclusion

The rapidly growing body of literature on biochar addition to soils has generally indicated that biochar has the potential to alter soil N, P dynamic transformations, soil nutrient leaching, crop growth and yield. Results from these recent studies using a diverse range of types of biochar have mostly focused on outcomes with few studies addressing specific mechanisms for altered nutrient transformations or uptake.<sup>137–156</sup> However, potential explanations for the positive influence of biochar on the soil environment do exist and can be summarized as follows:

- i. Biochar can enhance N<sub>2</sub> fixation in legumes by stimulating nodulation process or providing greater nutrient availability to the host plant;
- ii. Biochar effects on soil N mineralization and immobilization are varied and are related to factors including biochar production feedstock, conditions of biochar production, total adsorption capacity, and soil type;
- iii. Biochar effects on gaseous soil N emissions are dependent upon soil physicochemical properties and changes in the abundance and diversity of related microbial community followed by biochar addition to soil;
- iv. Biochar application to soil may influence P availability and uptake by altering P solubility, altering P mineralization processes and by the biochar serving as a source of P in low P soils;
- v. The surface chemistry of biochar is greatly responsible for adsorption potential and thus the potential to alter nutrient leaching;
- vi. Enhanced soil fertility, changes in soil-water status, and specific stimulation compound within biochar can further improve plant growth and crop yield. This mini-review was intended to simply serve as an interim update on the effect of biochar on soil nutrient transformations, losses and influence on plant productivity. Future research must seek to elucidate mechanisms and processes to allow for a better understanding of how biochar influences the soil environment and plant productivity.

## Acknowledgements

None.

## Conflict of interest

The author declares no conflict of interest.

## References

1. Lehmann J. Bio-energy in the black. *Frontiers in Ecology and the Environment*. 2007;5(7):381–387.
2. Lehmann J, Joseph S. *Biochar for Environmental Management: science, technology and implementation*. 2nd ed. Routledge; 2015.
3. Woolf D, Amonette JE, Perrott SFA, et al. Sustainable biochar to mitigate global climate change. *Nat Commun*. 2010;1(5):1–9.

4. Clough TJ, Condon LM, Kamman C, et al. A review of biochar and soil nitrogen dynamics. *Agronomy*. 2013;3(2):275–293.
5. Jeffery S, Abalos D, Spokas KA, et al. Biochar effects on crop yield. In: Johannes Lehmann, et al. editors. *Biochar for Environmental Management: Science, Technology and Implementation*. 2nd ed. Routledge; 2015. p. 301–325.
6. Stevenson FJ. *Cycles of Soils: carbon, nitrogen, phosphorus, sulfur, micronutrients*. 2nd ed. John Wiley & Sons; 1999. 448 p.
7. Schimel JP, Bennett J. Nitrogen mineralization: challenges of a changing paradigm. *Ecology*. 2004;85(3):591–602.
8. Vitousek PM, Cassman K, Cleveland C, et al. Towards an ecological understanding of biological nitrogen fixation. *Biogeochemistry*. 2002;57(1):1–45.
9. Robertson GP, Groffman PM. *Nitrogen transformations*. In: Paul EA editor. New York, USA: Springer; 2015. p. 421–446.
10. Mia S, Groenigen VJW, Voorde VTFJ, et al. Biochar application rate affects biological nitrogen fixation in red clover conditional on potassium availability. *Agriculture, Ecosystems & Environment*. 2014;191:83–91.
11. Güereña DT, Lehmann J, Thies JE, et al. Partitioning the contributions of biochar properties to enhanced biological nitrogen fixation in common bean (*Phaseolus vulgaris*). *Biology and Fertility of Soils*. 2015;51(4):479–491.
12. Quilliam RS, DeLuca TH, Jones DL. Biochar application reduces nodulation but increases nitrogenase activity in clover. *Plant and Soil*. 2013;366(1):83–92.
13. Zwieten VL, Rose T, Herridge D, et al. Enhanced biological N<sub>2</sub> fixation and yield of faba bean (*Vicia faba* L.) in an acid soil following biochar addition: dissection of causal mechanisms. *Plant and Soil*. 2015;395(1):7–20.
14. Rose TJ, Julia CC, Shepherd M, et al. Faba bean is less susceptible to fertilizer N impacts on biological N<sub>2</sub> fixation than chickpea in monoculture and intercropping systems. *Biology and Fertility of Soils*. 2016;52(2):271–276.
15. Mollinedo J, Schumacher TE, Chintala R. Biochar effects on phenotypic characteristics of “wild” and “sickle” *Medicago truncatula* genotypes. *Plant and Soil*. 2016;400(1–2):1–14.
16. Rondon MA, Lehmann J, Ramírez J, et al. Biological nitrogen fixation by common beans (*Phaseolus vulgaris* L.) increases with bio-char additions. *Biology and Fertility of Soils*. 2007;43(6):699–708.
17. Güereña D, Lehmann J, Hanley K, et al. Nitrogen dynamics following field application of biochar in a temperate North American maize-based production system. *Plant and Soil*. 2013;365(1):239–254.
18. Major J, Rondon M, Molina D, et al. Maize yield and nutrition during 4 years after biochar application to a Colombian savanna oxisol. *Plant and Soil*. 2010;333(1):117–128.
19. Backer RGM, Schwinghamer TD, Whalen JK, et al. Crop yield and SOC responses to biochar application were dependent on soil texture and crop type in southern Quebec, Canada. *Journal of Plant Nutrition and Soil Science*. 2016;179(3):399–408.
20. Hass A, Gonzalez JM, Lima IM, et al. Chicken manure biochar as liming and nutrient source for acid appalachian soil. *J Environ Qual*. 2012;41(4):1091–1106.
21. Tagoe SO, Horiuchi T, Matsui T. Effects of carbonized and dried chicken manures on the growth, yield, and N content of soybean. *Plant and Soil*. 2008;306(1):211–220.
22. Wurst S, Beersum VS. The impact of soil organism composition and activated carbon on grass-legume competition. *Plant and Soil*. 2009;314(1–2):1–9.
23. Egamberdieva D, Wirth S, Behrendt U, et al. Biochar treatment resulted in a combined effect on soybean growth promotion and a shift in plant growth promoting rhizobacteria. *Front Microbiol*. 2016;7:209.
24. Koes RE, Quattrocchio F, Mol JNM. The flavonoid biosynthetic pathway in plants: Function and evolution. *Bio Essays*. 1994;16(2):123–132.
25. Gundale MJ, DeLuca TH. Temperature and substrate influence the chemical properties of charcoal in the ponderosa pine/Douglas-fir ecosystem. *For Ecol Manage*. 2006;231(1–3):86–93.
26. Spokas KA, Baker JM, Reicosky DC. Ethylene: potential key for biochar amendment impacts. *Plant and Soil*. 2010;333(1–2):443–452.
27. DeLuca TH, Gundale MJ, MacKenzie MD, et al. Biochar effects on soil nutrient transformations. Johannes Lehmann, et al. editors. *Biochar for Environmental Management: Science, Technology and Implementation*. 2nd ed. Routledge; 2015. p. 421–454.
28. Lehmann J, da Silva JP, Steiner C, et al. Nutrient availability and leaching in an archaeological Anthrosol and a Ferralsol of the Central Amazon basin: fertilizer, manure and charcoal amendments. *Plant and Soil*. 2003;249(2):343–357.
29. Steiner C, Teixeira WG, Lehmann J, et al. Long term effects of manure, charcoal and mineral fertilization on crop production and fertility on a highly weathered Central Amazonian upland soil. *Plant and Soil*. 2007;291(1):275–290.
30. Nelissen V, Rütting T, Huygens D, et al. Maize biochars accelerate short-term soil nitrogen dynamics in a loamy sand soil. *Soil Biology and Biochemistry*. 2012;55:20–27.
31. DeLuca TH, MacKenzie MD, Gundale MJ, et al. Wildfire-produced charcoal directly influences nitrogen cycling in ponderosa pine forests. *American Society of Agronomy*. 2006;70(2):448–453.
32. Xu N, Tan G, Wang H, et al. Effect of biochar additions to soil on nitrogen leaching, microbial biomass and bacterial community structure. *European Journal of Soil Biology*. 2016;74:1–8.
33. Pereira EIP, Suddick EC, Mansour I, et al. Biochar alters nitrogen transformations but has minimal effects on nitrous oxide emissions in an organically managed lettuce mesocosm. *Biology and Fertility of Soils*. 2015;51(5):573–582.
34. Gundale MJ, Nilsson M, Pluchon N, et al. The effect of biochar management on soil and plant community properties in a boreal forest. *GCB Bioenergy*. 2015;8(4):777–789.
35. Luo X, Chen L, Zheng H, et al. Biochar addition reduced net N mineralization of a coastal wetland soil in the Yellow River Delta, China. *Geoderma*. 2016;282:120–128.
36. Dempster DN, Gleeson DB, Solaiman ZM, et al. Decreased soil microbial biomass and nitrogen mineralisation with *Eucalyptus* biochar addition to a coarse textured soil. *Plant and Soil*. 2012;354(1):311–324.
37. Kelly CN, Calderon FC, Acosta MV, et al. Switchgrass biochar effects on plant biomass and microbial dynamics in two soils from different regions. *Pedosphere*. 2015;25(3):329–342.
38. Prommer J, Wanek W, Hofhansl F, et al. Biochar decelerates soil organic nitrogen cycling but stimulates soil nitrification in a temperate arable field trial. *PLoS One*. 2014;9(1):e86388.
39. Ulyett J, Sakrabani R, Kibblewhite M, et al. Impact of biochar addition on water retention, nitrification and carbon dioxide evolution from two sandy loam soils. *European Journal of Soil Science*. 2014;65(1):96–104.
40. Robertson GP, Groffman PM. *Nitrogen transformations*. In: Paul EA editor. *Soil Microbiology, Ecology and Biochemistry*. 3rd ed. Springer; 2007. p. 341–364.
41. Lehmann J, Gaunt J, Rondon M. Bio-char sequestration in terrestrial ecosystems—a review. *Mitigation and Adaptation Strategies for Global Change*. 2006;11(2):395–419.

42. Bruun EW, Ambus P, Egsgaard H, et al. Effects of slow and fast pyrolysis biochar on soil C and N turnover dynamics. *Soil Biology and Biochemistry*. 2012;46:73–79.
43. Budai A, Rasse DP, Lagomarsino A, et al. Biochar persistence, priming and microbial responses to pyrolysis temperature series. *Biology and Fertility of Soils*. 2016;52(6):749–761.
44. Sigua GC, Novak JM, Watts DW, et al. Impact of switchgrass biochars with supplemental nitrogen on carbon–nitrogen mineralization in highly weathered Coastal Plain Ultisols. *Chemosphere*. 2016;145:135–141.
45. Novak JM, Busscher WJ, Watts DW, et al. Short-term CO<sub>2</sub> mineralization after additions of biochar and switchgrass to a Typic Kandiuult. *Geoderma*. 2010;154(3–4):281–288.
46. Ouyang W, Geng X, Huang W, et al. Soil respiration characteristics in different land uses and response of soil organic carbon to biochar addition in high-latitude agricultural area. *Environ Sci Pollut Res Int*. 2016;23(3):2279–2287.
47. Jones DLL, Murphy DVV, Khalid M, et al. Short-term biochar-induced increase in soil CO<sub>2</sub> release is both biotically and abiotically mediated. *Soil Biology and Biochemistry*. 2011;43(8):1723–1731.
48. Cayuela ML, Sánchez MMA, Roig A, et al. Biochar and denitrification in soils: when, how much and why does biochar reduce N<sub>2</sub>O emissions? *Scientific Report*. 2013;3:56.
49. Case SDCC, McNamara NP, Reay DS, et al. Biochar suppresses N<sub>2</sub>O emissions while maintaining N availability in a sandy loam soil. *Soil Biology and Biochemistry*. 2015;81(2):178–185.
50. Ameloot N, Maenhout P, De Neve S, et al. Biochar-induced N<sub>2</sub>O emission reductions after field incorporation in a loam soil. *Geoderma*. 2016;267:10–16.
51. Harter J, Krause HM, Schuettler S, et al. Linking N<sub>2</sub>O emissions from biochar-amended soil to the structure and function of the N-cycling microbial community. *ISME J*. 2013;8(10):660–674.
52. Jones DL, Rousk J, Edwards Jones G, et al. Biochar-mediated changes in soil quality and plant growth in a three year field trial. *Soil Biol Biochem*. 2012;45:113–124.
53. L Van Zwieten, BP Singh, SWL Kimber, et al. An incubation study investigating the mechanisms that impact N<sub>2</sub>O flux from soil following biochar application. *Agric Ecosyst Environ*. 2014;191:53–62.
54. Cayuela ML, van Zwieten L, Singh BP, et al. Biochar's role in mitigating soil nitrous oxide emissions: A review and meta-analysis. *Agric Ecosyst Environ*. 2014;191:5–16.
55. Šimek M, Jiřová L, Hopkins DW. What is the so-called optimum pH for denitrification in soil? *Soil Biol Biochem*. 2002;34(9):1227–1234.
56. Van Zwieten L, Singh B, Joseph S, et al. Biochar and emissions of non-CO<sub>2</sub> greenhouse gases from soil. *Biochar Environ Manag Sci Technol*. 2009. p. 227–250.
57. Yanai Y, Toyota K, Okazaki M. Effects of charcoal addition on N<sub>2</sub>O emissions from soil resulting from rewetting air-dried soil in short-term laboratory experiments. *Soil Sci Plant Nutr*. 2007;53(2):181–188.
58. Van Zwieten AL, Kimber S, Morris S, et al. Influence of biochars on flux of N<sub>2</sub>O and CO<sub>2</sub> from Ferrosol. *Aust J Soil Res*. 2010;48(6):555–568.
59. Yuan J-H, Xu R-K, Zhang H. The forms of alkalis in the biochar produced from crop residues at different temperatures. *Bioresour Technol*. 2011;102(3):3488–3497.
60. Dougherty BW. *Biochar as a cover for dairy manure lagoons: reducing odor and gas emissions while capturing nutrients*. Doctoral dissertation, Oregon State University; 2016.
61. Mandal S, Thangarajan R, Bolan NS, et al. Biochar-induced concomitant decrease in ammonia volatilization and increase in nitrogen use efficiency by wheat. *Chemosphere*. 2016;142:120–127.
62. Taghizadeh-Toosi A, Clough TJ, Sherlock RR, et al. A wood based low-temperature biochar captures NH<sub>3</sub>-N generated from ruminant urine-N, retaining its bioavailability. *Plant Soil*. 2012;353(1):73–84.
63. Doydora SA, Cabrera ML, Das KC, et al. Release of nitrogen and phosphorus from poultry litter amended with acidified biochar. *Int J Environ Res Public Health*. 2011;8(12):1491–1502.
64. Steiner C, Das KCC, Melear N, et al. Reducing nitrogen loss during poultry litter composting using biochar. *J Environ Qual*. 2010;39(4):1236–1242.
65. Ryan PR, Delhaize E, Jones DL. Function and mechanism of organic anion exudation from plant roots. *Annu Rev Plant Biol*. 2001;52(1):527–560.
66. Knoepp JD, Reynolds BC, Crossley DA, et al. Long-term changes in forest floor processes in southern Appalachian forests. *For Ecol Manage*. 2005;220(1–3):300–312.
67. Ippolito JA, Spokas KA, Novak JM, et al. Biochar elemental composition and factors influencing nutrient retention. In: Johannes Lehmann and Steven Joseph editors. *Biochar for Environmental Management: Science, Technology and Implementation*. 2nd ed. Routledge; 2015. p. 139–163.
68. Wang T, Camps-Arbestain M, Hedley M, et al. Predicting phosphorus bioavailability from high-ash biochars. *Plant Soil*. 2012;357(1–2):173–187.
69. Gaskin JW, Steiner C, Harris K, et al. Effect of low-temperature pyrolysis conditions on biochar for agricultural use. *Trans ASABE*. 2008;51(6):2061–2069.
70. Jin Y, Liang X, He M, et al. Manure biochar influence upon soil properties, phosphorus distribution and phosphatase activities: A microcosm incubation study. *Chemosphere*. 2016;142:128–135.
71. Gundale MJ, DeLuca TH. Charcoal effects on soil solution chemistry and growth of *Koeleria macrantha* in the ponderosa pine/Douglas-fir ecosystem. *Biol Fertil Soils*. 2007;43(3):303–311.
72. Soinne H, Hovi J, Tammeorg P, et al. Effect of biochar on phosphorus sorption and clay soil aggregate stability. *Geoderma*. 2014;219:162–167.
73. Madiba OF, Solaiman ZM, Carson JK, et al. Biochar increases availability and uptake of phosphorus to wheat under leaching conditions. *Biol Fertil Soils*. 2016;52:439.
74. Joseph S, Graber ER, Chia C, et al. Shifting paradigms: development of high-efficiency biochar fertilizers based on nano-structures and soluble components. *Carbon Manag*. 2013;4(3):323–343.
75. Yang F, Zhao L, Gao B, et al. The interfacial behavior between biochar and soil minerals and its effect on biochar stability. *Environ Sci Technol*. 2016;50(5):2264–2271.
76. Rodríguez H, Fraga R. Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol Adv*. 1999;17(4–5):319–339.
77. Suksabye P, Pimthong A, Dhurakit P, et al. Effect of biochars and microorganisms on cadmium accumulation in rice grains grown in Cd-contaminated soil. *Environ Sci Pollut Res*. 2016;23(2):962–973.
78. Anderson CR, Condon LM, Clough TJ, et al. Biochar induced soil microbial community change: implications for biogeochemical cycling of carbon, nitrogen and phosphorus. *Pedobiologia*. 2011;54(5):309–320.
79. Amador JA, Glucksman AM, Lyons JB, et al. Spatial distribution of soil phosphatase activity within a riparian forest I. *Soil Sci*. 1997;162(11):808–825.
80. Yoo G, Kang H. Effects of biochar addition on greenhouse gas emissions and microbial responses in a short-term laboratory experiment. *J Environ Qual*. 2012;41(4):1193–1202.



81. Oleszczuk P, Joško I, Kuśmierz M, et al. Microbiological, biochemical and ecotoxicological evaluation of soils in the area of biochar production in relation to polycyclic aromatic hydrocarbon content. *Geoderma*. 2014;213:502–511.
82. Bhaduri D, Saha A, Desai D, et al. Restoration of carbon and microbial activity in salt-induced soil by application of peanut shell biochar during short-term incubation study. *Chemosphere*. 2016;148:86–98.
83. Mackie KA, Marhan S, Ditterich F, et al. The effects of biochar and compost amendments on copper immobilization and soil microorganisms in a temperate vineyard. *Agric Ecosyst Environ*. 2015;201:58–69.
84. Pandey V, Patel A, Patra DDD. Biochar ameliorates crop productivity, soil fertility, essential oil yield and aroma profiling in basil (*Ocimum basilicum* L.). *Ecol Eng*. 2016;90:361–366.
85. Major J, Steiner C, Downie A. Biochar effects on nutrient leaching. In: Johannes Lehmann and Steven Joseph editors. *Biochar for environmental management: Science and technology*. 1st ed. Earthscan; 2009. p. 271–288.
86. Takaya CAA, Fletcher LAA, Singh S, et al. Phosphate and ammonium sorption capacity of biochar and hydrochar from different wastes. *Chemosphere*. 2016;145:518–527.
87. Cheng CH, Lehmann J, Thies JE, et al. Oxidation of black carbon by biotic and abiotic processes. *Org Geochem*. 2006;37(11):1477–1488.
88. Mukherjee A, Zimmerman AR, Hamdan R, et al. Physicochemical changes in pyrogenic organic matter (biochar) after 15 months of field aging. *Solid Earth*. 2014;5:693–704.
89. Heitkötter J, Marschner B. Interactive effects of biochar ageing in soils related to feedstock, pyrolysis temperature, and historic charcoal production. *Geoderma*. 2014;245:56–64.
90. Rogovska N, Laird DA, Karlen DL. Corn and soil response to biochar application and stover harvest. *F Crop Res*. 2016;187:96–106.
91. Sun F, Lu S. Biochars improve aggregate stability, water retention, and pore-space properties of clayey soil. *J Plant Nutr Soil Sci*. 2014;177(1):26–33.
92. Andrenelli MC, Maienza A, Genesio L, et al. Field application of pelletized biochar: Short term effect on the hydrological properties of a silty clay loam soil. *Agric Water Manag*. 2016;163:190–196.
93. Gul S, Whalen JK, Thomas BW, et al. Physico-chemical properties and microbial responses in biochar-amended soils: Mechanisms and future directions. *Agric Ecosyst Environ*. 2015;206:46–59.
94. Jaafar NM, Clode PL, Abbott LK. Biochar–soil interactions in four agricultural soils. *Pedosphere*. 2015;25(5):729–736.
95. Deenik JL, McClellan T, Uehara G, et al. Charcoal volatile matter content influences plant growth and soil nitrogen transformations. *Soil Sci Soc Am J*. 2010;74(4):1259–1270.
96. Jeffery S, Verheijen FGA, van der Velde M, et al. A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agric Ecosyst Environ*. 2011;144(1):175–187.
97. Chan KY, Van Zwieten L, Meszaros I, et al. Agronomic values of greenwaste biochar as a soil amendment. *Soil Res*. 2008;45(8):629–634.
98. Gaskin JW, Speir RA, Harris K, et al. Effect of peanut hull and pine chip biochar on soil nutrients, corn nutrient status, and yield. *Agron J*. 2010;102(2):623–633.
99. Kammann C, Graber ER. Biochar effects on plant ecophysiology. In: Johannes Lehmann, et al. editors. *Biochar for Environmental Management: Science, Technology and Implementation*. 2nd ed. Routledge; 2015. p. 391–419.
100. Thies JE, Rillig MC, Graber ER. Biochar effects on the abundance, activity and diversity of the soil biota. In: Johannes Lehmann, et al. editors. *Biochar Environmental Management Science Technology and implementation*. 2nd ed. Routledge; 2015. p. 327–389.
101. Kleber M, Hockaday W, Nico PS. Characteristics of biochar: macro-molecular properties. In: Johannes Lehmann, et al. editors. *Biochar for Environmental Management: Science, Technology and Implementation*. Routledge; 2015. p. 111–137.
102. Kammann CI, Linsel S, Gößling JW, et al. Influence of biochar on drought tolerance of *Chenopodium quinoa* Willd and on soil–plant relations. *Plant and Soil*. 2011;345(1):195–210.
103. Yu X, Wu C, Fu Y, et al. Three-dimensional pore structure and carbon distribution of macroaggregates in biochar-amended soil. *European Journal of Soil Science*. 2016;67(1):109–120.
104. Abel S, Peters A, Trinks S, et al. Impact of biochar and hydrochar addition on water retention and water repellency of sandy soil. *Geoderma*. 2013;202–203:183–191.
105. Hansen V, Nielsen HH, Petersen CT, et al. Effects of gasification biochar on plant-available water capacity and plant growth in two contrasting soil types. *Soil and Tillage Research*. 2016;161:1–9.
106. Sun H, Hockaday WC, Masiello CA, et al. Multiple controls on the chemical and physical structure of biochars. *Ind Eng Chem Res*. 2012;51(9):3587–3597.
107. Spokas KA, Novak JM, Stewart CE, et al. Qualitative analysis of volatile organic compounds on biochar. *Chemosphere*. 2011;85(5):869–882.
108. Bailly A, Weisskopf L. The modulating effect of bacterial volatiles on plant growth: current knowledge and future challenges. *Plant Signal Behav*. 2012;7(1):79–85.
109. Hofmann NR. Volatile organic compounds: A bacterial contribution to plant sulfur nutrition. *Plant Cell*. 2013;25(7):2381.
110. Buss W, Mašek O. Mobile organic compounds in biochar – A potential source of contamination – Phytotoxic effects on cress seed (*Lepidium sativum*) germination. *Journal of Environmental Management*. 2014;137:111–119.
111. Dutta T, Kwon E, Bhattacharya SS, et al. Polycyclic aromatic hydrocarbons (PAHs) and volatile organic compounds (VOCs) in biochar and biochar-amended soil: A review. *GCB Bioenergy*. 2016.
112. Beyer EM. A Potent inhibitor of ethylene action in plants. *Plant Physiol*. 1976;58(3):268–271.
113. Eplee RE. Ethylene: A witchweed seed germination stimulant? *weed science*. 1975;23(5):433–436.
114. Abeles FB, Morgan PW, Saltveit MEJ. *Ethylene in plant biology*. 2nd ed. Academic press; 1992.
115. Joseph SD, Arbestain CM, Lin Y, et al. An investigation into the reactions of biochar in soil. *Soil Research*. 2010;48(7):501–515.
116. Warnock DD, Lehmann J, Kuyper TW, et al. Mycorrhizal responses to biochar in soil – concepts and mechanisms. *Plant and Soil*. 2007;300(1):9–20.
117. Graber ER, Frenkel O, Jaiswal AK, et al. How may biochar influence severity of diseases caused by soilborne pathogens? *Carbon Management*. 2014;5(2):169–183.
118. Elad Y, Cytryn E, Harel YM, et al. The biochar effect: plant resistance to biotic stresses. *Phytopathol Mediterr*. 2011;50:335–349.
119. Bonanomi G, Ippolito F, Scala F. A “black” future for plant pathology? Biochar as a new soil amendment for controlling plant diseases. *Journal of Plant Pathology*. 2015;97(2):223–234.



120. Chang J, Luo X, Li M, et al. Short-term influences of peanut-biochar addition on abandoned orchard soil organic N mineralization in north china. *Polish Journal of Environmental studies*. 2016;25(1):67–72.
121. Li M, Liu M, Li Z, et al. Soil N transformation and microbial community structure as affected by adding biochar to a paddy soil of subtropical China. *Journal of Integrative Agriculture*. 2016;15(1):209–219.
122. Marks EAN, Mattana S, Alcañiz JM, et al. Gasifier biochar effects on nutrient availability, organic matter mineralization, and soil fauna activity in a multi-year Mediterranean trial. *Agriculture Ecosystem Environment*. 2016;215:30–39.
123. Subedi R, Taupe N, Pelissetti S, et al. Greenhouse gas emissions and soil properties following amendment with manure-derived biochars: Influence of pyrolysis temperature and feedstock type. *Journal of Environmental Management*. 2016;166:73–83.
124. Ameloot N, Sleutel S, Das KC, et al. Biochar amendment to soils with contrasting organic matter level: Effects on N mineralization and biological soil properties. *GCB Bioenergy*. 2015;7(1):135–144.
125. Maestrini B, Herrmann AM, Nannipieri P, et al. Ryegrass-derived pyrogenic organic matter changes organic carbon and nitrogen mineralization in a temperate forest soil. *Soil Biology and Biochemistry*. 2014;69:291–301.
126. Castaldi S, Riondino M, Baronti S, et al. Impact of biochar application to a Mediterranean wheat crop on soil microbial activity and greenhouse gas fluxes. *Chemosphere*. 2011;85(9):1464–1471.
127. Streubel JD, Collins HP, Perez GM, et al. Influence of contrasting biochar types on five soils at increasing rates of application. *Soil Science Society of American Journal*. 2011;75(4):1402–1413.
128. DeLuca TH, Nilsson M, Zackrisson O. Nitrogen mineralization and phenol accumulation along a fire chronosequence in northern Sweden. *Oecologia*. 2002;133(2):206–214.
129. Berglund LM, DeLuca TH, Zackrisson O. Activated carbon amendments to soil alters nitrification rates in Scots pine forests. *Soil Biology and Biochemistry*. 2004;36(12):2067–2073.
130. Novak JM, Busscher WJ, Laird DL, et al. Impact of biochar amendment on fertility of a southeastern coastal plain soil. *Soil Science*. 2009;174(2).
131. Sun L, Li L, Chen Z, Wang J, et al. Combined effects of nitrogen deposition and biochar application on emissions of N<sub>2</sub>O, CO<sub>2</sub> and NH<sub>3</sub> from agricultural and forest soils. *Soil Science and Plant Nutrition*. 2014;60(2):254–265.
132. Jordan G, Predotova M, Ingold M, et al. Effects of activated charcoal and tannin added to compost and to soil on carbon dioxide, nitrous oxide and ammonia volatilization. *Journal of Plant Nutrition Soil Science*. 2014;178(2):218–228.
133. Subedi R, Kammann C, Pelissetti S, et al. Does soil amended with biochar and hydrochar reduce ammonia emissions following the application of pig slurry? *European Journal of Soil Science*. 2015;66(6):1044–1053.
134. Chen CR, Phillips IR, Condon LM, et al. Impacts of green waste biochar on ammonia volatilisation from bauxite processing residue sand. *Plant and Soil*. 2013;367(1):301–312.
135. Gao S, Krull HK, Bidwell AL, et al. Locally produced wood biochar increases nutrient retention and availability in agricultural soils of the San Juan Islands, USA. *Agriculture, Ecosystem & Environment*. 2016;233:43–54.
136. Zwetsloot MJ, Lehmann J, Bauerle T, et al. Phosphorus availability from bone char in a P-fixing soil influenced by root-mycorrhizae-biochar interactions. *Plant and Soil*. 2016. p. 1–11.
137. Frišták V, Soja G. Effect of wood-based biochar and sewage sludge amendments for soil phosphorus availability. *Nov Biotechnol Chim*. 2015;14(1):104–115.
138. Vanek SJ, Lehmann J. Phosphorus availability to beans via interactions between mycorrhizas and biochar. *Plant and Soil*. 2015;395(1):105–123.
139. Parvage MM, Ulén B, Eriksson J, et al. Phosphorus availability in soils amended with wheat residue char. *Biology and Fertility of Soils*. 2013;49(2):245–250.
140. Jin Y, Liang X, He M, et al. Manure biochar influence upon soil properties, phosphorus distribution and phosphatase activities: A microcosm incubation study. *Chemosphere*. 2016;142:128–135.
141. Yang X, Liu J, McGrouther K, et al. Effect of biochar on the extractability of heavy metals (Cd, Cu, Pb, and Zn) and enzyme activity in soil. *Environ Sci Pollut Res*. 2016;23(2):974–984.
142. Chen J, Liu X, Zheng J, et al. Biochar soil amendment increased bacterial but decreased fungal gene abundance with shifts in community structure in a slightly acid rice paddy from Southwest China. *Applied Soil Ecology*. 2013;71:33–44.
143. Masto RE, Kumar S, Rout TKK, et al. Biochar from water hyacinth (*Eichornia crassipes*) and its impact on soil biological activity. *Catena*. 2013;71:64–71.
144. Ferreira PJ, Gascó G, Gutiérrez B, et al. Soil biochemical activities and the geometric mean of enzyme activities after application of sewage sludge and sewage sludge biochar to soil. *Biology and Fertility of Soils*. 2012;48(5):511–517.
145. Kanthle AK, Lenka NK, Lenka S, et al. Biochar impact on nitrate leaching as influenced by native soil organic carbon in an Inceptisol of central India. *Soil and Tillage Research*. 2016;157:65–72.
146. Yuan H, Lu T, Wang Y, et al. Sewage sludge biochar: Nutrient composition and its effect on the leaching of soil nutrients. *Geoderma*. 2016;267:17–23.
147. Eykelbosh AJ, Johnson MS, Couto EG. Biochar decreases dissolved organic carbon but not nitrate leaching in relation to vinasse application in a Brazilian sugarcane soil. *J Environ Manage*. 2015;149:9–16.
148. Hardie MA, Oliver G, Clothier BE, et al. Effect of biochar on nutrient leaching in a young apple orchard. *J Environ Qual*. 2015;44(4):1273–1282.
149. Troy SM, Lawlor PG, O Flynn CJ, et al. The impact of biochar addition on nutrient leaching and soil properties from tillage soil amended with pig manure. *Water Air & Soil Pollution*. 2014;225:1900.
150. Ventura M, Sorrenti G, Panzacchi P, et al. Biochar reduces short-term nitrate leaching from a horizon in an apple orchard. *J Environ Qual*. 2013;42(1):76–82.
151. Yao Y, Gao B, Zhang M, et al. Effect of biochar amendment on sorption and leaching of nitrate, ammonium, and phosphate in a sandy soil. *Chemosphere*. 2012;89(11):1467–1471.
152. Major J, Rondon M, Molina D, et al. Nutrient leaching in a colombian savanna oxisol amended with biochar. *J Environ Qual*. 2012;41(4):1076–1086.
153. Ippolito JA, Novak JM, Busscher WJ, et al. Switchgrass biochar affects two Aridisols. *J Environ Qual*. 2012;41(4):1123–1130.
154. Kameyama K, Miyamoto T, Shiono T, et al. Influence of sugarcane bagasse-derived biochar application on nitrate leaching in calcareous dark red soil. *J Environ Qual*. 2012;41(4):1131–1137.
155. Borchard N, Wolf A, Laabs V, et al. Physical activation of biochar and its

- meaning for soil fertility and nutrient leaching—a greenhouse experiment. *Soil Use and Management*. 2012;28(2):177–184.
156. Ding Y, Liu YX, Wu WX, et al. Evaluation of biochar effects on nitrogen retention and leaching in multi-layered soil columns. *Water, Air & Soil Pollution*. 2010;213(1):47–55.
157. Laird D, Fleming P, Wang B, et al. Biochar impact on nutrient leaching from a Midwestern agricultural soil. *Geoderma*. 2010;158(3):436–442.