

Optimization of factors affecting *Agrobacterium*-mediated genetic transformation in groundnut (*Arachis hypogaea* L.)

Abstract

The efficiency of *Agrobacterium*-mediated genetic transformation of groundnut is dependent on various factors such as type and age of explants, antibiotics used for selection, Acetosyringone (AS), duration of co cultivation and infection period. Hence, we have attempted to standardize these parameters to improve the *Agrobacterium* mediated genetic transformation efficiency in groundnut cv ICG 13942. To evaluate the optimization of factors affecting *Agrobacterium* mediated genetic transformation in which we used three different explants (Deembryonated cotyledon-DC, Leaflet-LL, Cotyledonary node-CN) and studied, the effect of inoculum concentration (0.5, 1.0 and 1.5 OD₆₀₀ of *A. tumefaciens*), pre culture period (1-11 days), co cultivation period (1-7 days), Acetosyringone concentration (25-200mg/L) and antibiotic (Kan) concentration (25-150mg/L) on the efficiency of genetic transformation. The transformation was verified by GUS staining and by means of PCR amplification of the *uidA* and *nptII* genes. The optimized factors for *A. tumefaciens* LBA 4404 harboring pBAL2 binary vector mediated transformation in groundnut cv ICG 13942 are: Pre culture period-3 days, Concentration of *A. tumefaciens*-1.0 at OD₆₀₀, Co-cultivation period-4 days, Kan concentration-100mg/L for DC &CN, 75mg/L for LL explants, Concentration of AS-100mg/L during infection and co cultivation and Cefotaxime concentration-250mg/L. Thus, the optimized protocol showed the enhanced transient transformation frequencies such as 18.13% for DC, 15.33% for LL and 12.32% for CN explants.

Keywords: *agrobacterium tumefaciens*, *arachis hypogaea*, GUS expression, *npt-II*, polymerase chain reaction

Volume 8 Issue 3 - 2018

Rajinikanth Marka, Rama Swamy Nanna

Department of Biotechnology, Kakatiya University, Plant Biotechnology Research Laboratory, India

Correspondence: Rama Swamy Nanna, Plant Biotechnology Research Laboratory, Department of Biotechnology, Kakatiya University, Warangal-506009, Telangana, India, Email swamynr.dr@gmail.com

Received: October 24, 2017 | **Published:** June 04 2018

Abbreviations: KAN, Kanamycin; AS, Acetosyringone; CN, Cotyledonary node; LL, Leaflet; DC, De embryonated cotyledon; nptII, Neomycin phosphotransferase; NAA, α -Naphthalene acetic acid; BAP, 6-Benzylaminopurine

Introduction

Groundnut/Peanut (*Arachis hypogaea* L.) has great potential in third world countries to reduce hunger and malnutrition.¹ It is widely grown in the semi-arid tropics and India ranks second by 12.5% in world production.² The production of groundnut is constrained by different biotic and abiotic factors by causing decrease in yield.³ To develop stress tolerance through conventional breeding takes longer time and passes through various incompatibility barriers. To overcome these, the biotechnological tools play a great role for developing stress tolerance through introduction of alien genes into elite germplasm of crop plants through genetic transformation technology.^{4,6} The various factors that affect significant differences in T-DNA delivery and transformation included explant type, duration of pre-culture of explant, incubation period in bacterial suspension, the concentration of *A. tumefaciens* during infection, the use and concentration of inducers such as Acetosyringone during infection, antibiotics used for selection and the extent of the co-culture period were optimized to establish a transformation protocol of common bean (*Phaseolus vulgaris*).⁷ and in chickpea.⁸ Legumes in general are recalcitrant to tissue culture and *Agrobacterium* mediated transformation is also highly genotype specific.⁹ Several reports have been published on efficient regeneration from diverse explants of peanut,^{10,11} but less

success has been recorded with genetic transformation of *A. hypogaea*. This is due to lack of efficient protocols to induce adventitious shoot buds after the transformation.¹²⁻¹⁴ Recently we have developed the reproducible regeneration protocol in groundnut cv ICG 13942.¹⁵⁻¹⁷ The aim of this work was to develop a reproducible and efficient method of *A. tumefaciens* mediated genetic transformation and to evaluate the factors that affect the genetic transformation efficiency by using marker genes (*nptII*; *uidA*) and later the same was used for *Agrobacterium* mediated transformation with the *Tc Chitinase-I* gene to develop the resistance to Tikka disease in cv ICG 13942 for the first time. Therefore, in the present report we describe the successful transformation and regeneration of putatively transformed plants in groundnut cv ICG 13942.

Material and methods

Plant material, culture initiation and maintenance

The mature seeds of groundnut cv ICG 13942, obtained from the germplasm bank of ICRISAT, Patancheru, Hyderabad, Telangana, India were used. Dried seeds were washed under running tap water for 10-15 min followed by treating with Tween-20 (5%-v/v) for 5 min. Later these were rinsed twice in sterile distilled water. Afterwards the seeds were surface sterilized with 0.1% (w/v) HgCl₂ for 8 min followed by rinsing in sterilized distilled water for 3-4 times under aseptic conditions and soaked for 24 h in sterile distilled water. The experiments were conducted on modified MS¹⁸ medium (MMS medium) containing MS basal salts and B₅ vitamins,¹⁹ 100 mg/L myo-inositol and 30 g/L sucrose. The pH of the medium was adjusted to

5.8±0.02 with either 0.1N NaOH or 0.1N HCl before addition of agar. The medium was solidified with 0.8% Difcobacto-agar and autoclaved at 121°C under 15 psi for 15-20 min. All the cultures were incubated at 25±2°C under cool white fluorescent light (1600lux) with a 16/8 h (light/dark) photoperiod.

Explants preparation

Three different kinds of explants i.e. de-embryonated cotyledon (DC), leaf let (LL) and cotyledonary node (CN) were used for transformation. Mature dry seeds were imbibed in sterile distilled water for 24 h after surface sterilization and kept for germination on autoclaved filter paper bridge/boat in ½ strength liquid MMS medium in culture tubes. In the case of CN explants, they were excised from 7days old *in vitro* grown seedlings and removed the cotyledons, radical, shoot tip parts, LL explants from axenic seedlings (2 weeks

old) were cut into 0.8-1.0 cm² size and DC explants were isolated from the soaked (24 hrs) seeds and these were inoculated on the following medium: CN explants on MMS+20 mg/L BAP (SIM),¹⁵ DC explants on MMS+0.5mg/L IAA+15 mg/L TDZ (SIM)¹⁶ and Leaflet explants on MMS+2mg/L AgNO₃+10mg/L BAP+0.5mg/L IAA (SIM).¹⁷

Agrobacterium strain and plasmid vector

The *Agrobacterium tumefaciens* strain LBA 4404 harboring the binary vector pBAL2 (18.8Kb) was used for optimization of transformation protocol in groundnut cv13942. The binary vector pBAL2 consists of reporter gene (β -glucuronidase-*gus/uidA*) by the CaMV 35S promoter and terminator sequences and the selectable marker gene (*npt II*) driven by the nopaline synthase (NOS) promoter and terminator sequences respectively (Figure 1).




Figure 1 Linear diagram of T-DNA portion of pBAL2.

Transformation and regeneration of transformants

The *A. tumefaciens* strain LBA4404 harboring pBAL2 was grown on YEM agar medium containing 50mg/L of Kan and 10mg/L Rif. A single bacterial colony was inoculated in 50ml liquid YEM medium containing the same antibiotics and grown overnight at 28°C in an incubator shaker at 120rpm. 5ml of this overnight culture was re inoculated in 50 ml fresh YEM liquid medium containing 50mg/L Kan and grown overnight. The bacterial cell density was determined at OD₆₀₀ of 0.5 to 1.5. Later the bacterial suspension was pelleted at 5,000rpm for 10 min and re suspended in 25ml hormone-free liquid half-strength MS medium. The explants, DC, LL and CN were pre-cultured for 1-11 days on shoot induction medium (SIM) prior to infection with bacteria. The trimmed explants were immersed in *Agrobacterium* suspension containing 25-200mg/L AS for 10-30 minutes and subjected to manual shaking. Later, the explants were dried on sterile tissue paper and co-cultivated for 1-7 days on SIM along with 100mg/L AS and incubated in the dark at 28±2°C. Afterwards the explants were washed with 250mg/l cefotaxime for 5 min followed by rinsing with sterile distilled water thrice for 5 min and dried on sterile tissue paper. All these explants were cultured on selection medium containing SIM+75mg/L Kan for leaflet and 100mg/L Kan for DC and CN explants and 250mg/l cefotaxime for 2-3 weeks. The explants with Kan^R shoots were sub cultured on SIM with 50mg/L Kan for further proliferation of shoots. Later each micro-shoot was excised and cultured on MMS+50 mg/L Kan+0.5 mg/L BAP for shoot elongation. After elongation, the micro-shoots were transferred on to root induction medium (RIM) containing 1mg/L NAA+50mg/L Kan. All the cultures were incubated at 25±2 °C with a 16/8 hr photoperiod.

These Kan^R plantlets were shifted to plastic cups containing the soil mix, sand: soil: vermiculite (1:1:1) and acclimatized in the growth chamber. Established plantlets were transferred to green house and allowed to grow in an earthenware pots for maturity.

Histochemical GUS analysis

The commercial kit “ β -Glucuronidase Reporter Gene Staining Kit” was performed, following the protocol recommended by the supplier (Sigma Aldrich), after which it was incubated at 37°C for 24h. The shoots were then rinsed twice with ethanol at 70% and once with acetone at 50%, according to the recommendations of Jefferson et al.²⁰ Finally, the number of explants presenting blue coloring and their percentage per treatment was registered by stereoscopic observation.

PCR analysis of putative transformants

The leaf genomic DNA from T₀ plants of cv ICG 13942 was isolated by Cetyl Trimethyl Ammonium Bromide (CTAB) method and used for molecular characterization of putative transgenic by PCR using *nptII* and *uidA* gene specific primers. The *npt II* primer sequences were, (F) 5'-GCT TGG GTG GAG AGG GCT ATT-3' (R) 5'-AGA ACT CGT CAA GAA GGC GA -3' and *uidA* gene specific primer sequences were, (F) 5'- TTT AAC TAT GCC GGG ATC CAT CGC - 3' (R) 5'- CCA GTC GAG CAT CTC TTC AGC GT - 3'. The PCR for *nptII* was carried out by initial denaturation at 94°C for 5 min followed by 35 cycles of 94°C for 1 min, 54°C for 1 min and 72°C for 1.30 min and final extension at 72°C for 10 min. A PCR programme for *uidA* gene was carried out by initial denaturation at 94°C for 5 min followed by 35 cycles of 94°C for 1 min, 64°C for 1.30 min and 72°C for 2 min and final extension at 72°C for 10 min. Plants

confirming positive with PCR were taken to maturity and their seeds were collected. The genomic DNA from the untransformed control plants and PCR +ve plants for *nptII* and *uid A* genes were used as negative and positive controls, respectively. The amplified fragments were separated by electrophoresis on a 0.8% agarose gel, using TE buffer. Gels were stained with ethidium bromide, photographed by using gel documentation system (Biorad).

Data analysis

The transformation efficiency was calculated in different explants.

Results and discussion

The effect of different parameters on the transformation was carried out. The parameters include pre culture of explants, bacterial cell density, bacterial infection period, co cultivation period, concentration of AS and cefotaxime. Hence, all the experiments were carried out with *Agrobacterium tumefaciens* strain LBA 4404 harboring the binary plasmid pBAL2 contains *nptII* and *uidA* marker genes and co cultivation on MMS medium supplemented with appropriate PGRs (SIM) with different explants (DC, LL and CN) of groundnut cv ICG 13942.

Factors affecting genetic transformation

Effect of different OD at 600nm

The pre cultured (3 days) DC, LL and CN explants were submerged in bacterial suspension at different OD₆₀₀ values of 0.5, 1.0, 1.5 bacterial suspension for 20 min and were cultured on SIM containing 100 mg/L AS for 4 days. The bacterial concentration of 1.0 value at OD₆₀₀ resulted the highest percentage of regeneration with transformation efficiency in DC (82.60%), LL (74.25%), CN (61.13%) explants of cv ICG 13942 (Figure 3a).

Effect of pre-culture period (days)

To test whether the pre culture period has influence on transformation, the DC, LL and CN explants were cultured on SIM for 1-11 days before the selection on Kan containing medium and also prior to the infection with *Agrobacterium* strain. Of these, the pre cultured explants for three days on SIM have shown the maximum percentage of response in all the three explants of peanut cv used (Figure 3b). Gradually the percentage of transformation was found to be reduced after 3rd day of pre culture because the explants pre cultured for 5th, 7th, 9th and 11th days were unable to receive the *Agrobacterium* due to the excessive growth of the explant. Among the explants, the DC (69.11%) and LL (66.23%) explants have shown the maximum transformation efficiency followed by CN (64.92%) explants of cv ICG 13942.

Effect of co-cultivation period

The pre cultured (3 days) DC, LL and CN explants were submerged in bacterial suspension for 20 min and were co cultivated on SIM containing 100 mg/L AS for 1-7 days. Four days of co-cultivation resulted the highest percentage of regeneration with transformation efficiency in DC (66.22%), LL (60.14%), CN (57.23%) explants of cv ICG 13942 (Figure 3c).

Effect of infection period

The pre cultured explants viz., DC, LL and CN were submerged

in bacterial suspension for 10, 15, 20, 30 min, and they were co cultivated on SIM containing 100 mg/L AS for 4 days. 20 min infection period had shown the highest percentage of transformation efficiency in DC (58.13%), LL (52.20%), CN (48.62%) explants of cv ICG 13942 (Figure 3d). For more than 20 min infection period, led to the explant abortion caused by bacterial contamination by reducing the transformation frequency.

Effect of antibiotics sensitivity (Kan)

The pre cultured DC, LL and CN explants were inoculated on SIM with different concentrations (25-150mg/L) of Kan (Figure 3e). The percentage of shoot induction response along with multiple shoots development/explants was found to be reduced as the concentration of Kan increased in all the explants tested. More than 90% of shoot induction response was inhibited at 100mg/L Kan in DC, CN explants (Figure 2a–f) and more than 85% at 75mg/L Kan in LL explants (Figure 2g–i). Beyond that level of Kan, showed the bleaching and necrosis of the explants.




Figure 2a-i Determination of antibiotics sensitivity (Kan) in three different types of explants (DC, CN, LL) of groundnut cv ICG 13942.

a-c) Shoot buds proliferation on SIM without antibiotics (control), Shoot buds formation at 100 mg/L Kan, Inhibition of shoot buds development at 125mg/L Kan (Note the necrosis of explants) in DC explants respectively, d-f) Shoot buds proliferation on SIM without antibiotics (control), Shoot buds formation at 100 mg/L Kan, Inhibition of shoot buds development at 125mg/L Kan (Note the necrosis of explants) in CN explants respectively, g-i) Shoot buds proliferation on SIM without antibiotics (control), Shoot buds formation at 75 mg/L Kan, Inhibition of shoot buds development at 100 mg/L Kan (Note the necrosis of explants) in LL explants respectively.

Effect of Acetosyringone (AS)

Different levels of AS (25- 200 mg/L) was tested and at 100 mg/L AS was found to be the best for enhancing the transformation efficiency in groundnut cv ICG 13942. The maximum percentage


(76.54%) of GUS positive explants was observed in DC explants, followed by 71.78% in LL and 68.21% in CN explants (Figure 3f).

Effect of explant age


Seven days old seedlings and 24 hours soaked seeds exhibited more GUS expression by showing maximum percentage of transformation efficiency in all the explants used. Transformation efficiency was thus affected by the age of the explant source with optimal transformation efficiencies of 85.13%, 73.13% and 61.09% for DC, LL and CN explants respectively (Figure 3g-h).

Effect of explant type


Susceptibility to *Agrobacterium* was dependent on the type of explant. In order to improve the protocol of genetic transformation, LL, CN explants were excised from 7-days old axenic seedlings and DC explants were excised from 24 hours soaked seeds and compared for their transformation efficiencies. The difference in transformation efficiencies was recorded. The highest transformation frequency was observed for DC (18.13%) followed by LL (15.33%) and CN (12.32%) explants (Figure 3i).




a) Effect of different values of O.D at 600nm




b) Effect of pre-culture period




c) Effect of cocultivation period




d) Effect of infection period




e) Effect of antibiotics sensitivity




f) Effect of Acetosyringone concentration



g) Effect of age of explant



h) Effect of soaking period



i) Effect of explant type

Figure 3a-i Transient transformation efficiency with different parameters.

Effect of cefotaxime concentration

After four days of co-cultivation, the explants (DC, LL & CN) were washed with sterile half-strength MS liquid medium containing 250mg/l Cefotaxime to prevent the over growth of *Agrobacterium* in the infected explants. Subsequently these infected explants were cultured on the selection medium containing SIM+100mg/L Kan+250mg/L Cefotaxime for DC & CN, SIM+75mg/L Kan+250mg/L Cefotaxime for LL explants for 2 weeks. Cefotaxime totally arrested the overgrowth of *Agrobacterium* which causes the death of the explants.

Confirmation of transformants

All these standardized parameters were used for *A. tumefaciens* LBA 4404 mediated transformation in groundnut cv ICG 13942. After co cultivation for 4 days, the DC, LL and CN were cultured on their respective selection medium for 2-3 weeks. The Kan^R shoots were developed from the transformed cells (Figure 4a-c). The explants with Kan^R shoots were sub cultured on SIM+50mg/L Kan for further proliferation. The individual micro-shoot was elongated on MMS+50mg/L Kan+0.5mg/L BAP. These elongated micro-shoots were rooted on RIM+1.0mg/L NAA+50 mg/L Kan. The Kan^R plantlets (T₀) were successfully acclimatized and maintained in the green house (Figure 4d-i).

GUS assay

To compare transient and stable T-DNA transformation, GUS

analysis was done in explants immediately after co-cultivation as well as in young growing tissues of T₀ generation of putative transformants. Transient GUS expression was not consistent, however, stable GUS expression was observed in young T₀ shoots of DC, LL and CN explants (Figure 4F-H). No GUS activity was observed in non-transformed (control) explants (Figure 4E). Explants without pre culture and mechanical injuries, at the site of regeneration, upon co cultivation with *Agrobacterium* showed no staining (DC, LL and CN). Whereas, pre cultured explants with mechanical injuries showed GUS activity at the wounded regions of the explants. Thus, we have found the typical expression pattern of the CaMV 35S promoter regulated *uidA* gene in transformed tissues. These results show the presence of GUS expression only in transformed tissues, clearly demonstrate the stability of the integrated gene in the putatively transformed plants of cv 13942 of groundnut.

Molecular confirmation

In the present study, molecular analysis was carried out by using PCR technique in T₀ plants. The amplification of the presence of *nptII* and *uidA* (*gus*) genes in Kan^R putatively transformed plants (T₀) was confirmed. These putatively transformed plants were obtained after three cycles of culture on selection medium after co cultivation with *A. tumefaciens* LBA 4404 strain harboring the binary vector pBAL2. The gene specific primers for *nptII* and *uid 'A'* genes were used for PCR amplification and the amplified DNA fragments were found 750bp and 1.9 Kb respectively (Figure 6) (Figure7). The transformation

efficiency was estimated on the basis of PCR +ve plants (*NPT-II* and *GUS*) developed in the present investigations. Out of 915 Kan^R plants developed only 141 plants showed the PCR +ve (*nptII* & *gus*) with 15.40% of transformation efficiency in three different types of explants (DC, LL and CN) used in cv ICG 13942 (Figure 5). Thus, the presence of the *uidA* and *nptII* genes in the putatively transformed plants (T₀) of groundnut cvs was confirmed by PCR analysis. PCR analysis revealed the presence of 1.9 kb *uidA* and 750bp *nptII* DNA fragments amplified from the genomic DNA of all T₀ transgenic plants. However, PCR products of *uidA* and *nptII* genes were not seen in untransformed (control - WT) plants (Figures 6,7), Lane C). All the samples of transgenic plants of cv 13942 gave the predicted size of DNA fragment of *uidA* gene (1.9Kb) (Figure 6, Lane 1-7). The band was not detected in the DNA sample from an untransformed (control) plant (Figure 6, Lane C). The DNA product with the size of 750bp *nptII* gene was amplified from total genomic DNA of the T₀ transgenic plants of cv 13942 of groundnut (Figure 7, Lane 1-7). This DNA fragment was not detected in the untransformed (control-WT) plants (Figure 7, Lane C). As a positive control, the *nptII* gene product was also amplified from the pBAL2 plasmid (Figure 7), Lane P). Based on the development of Kan^R plants, GUS assay and PCR detection, the presence of both *uidA* and *nptII* genes were confirmed in the T₀ transgenic groundnut plants.




Figure 4a-i *Agrobacterium* mediated transformation and GUS Assay of transgenic shoots (Kan^R) developed from DC, CN and LL explants of cv ICG 13942 by using the binary vector pBAL2. a-c Induction of Kan^R shoots on selection medium after 4 weeks of incubation from DC, CN and LL respectively, d) *In vitro* rooting of shoots on RIM supplemented with 1.0 mg/L NAA+ 50mg/L Kan (Note profuse rhizogenesis), e) Non transformed shoots (control-WT), f-h) Transgenic shoots developed from DC, LL and CN explants respectively, i) Plants are shifted to plastic pots containing soil mix and maintained in the green house.




Figure 5 Transformation efficiency in different explants of groundnut cv ICG 13942.




Figure 6 PCR screening of T₀ plants for the presence of the *uidA* gene using gene-specific primers in cv ICG 13942 of groundnut.

Lane M, DNA marker; Lane C, Control plant DNA (-ve control-WT); Lane P, Plasmid pBAL2 (+ve control); Lanes 1-3, T₀ plants developed from DC; Lanes 4-5, T₀ plants developed from LL; Lanes 6-7, T₀ plants developed from CN explants.




Figure 7 PCR screening of T₀ plants for the presence of the *nptII* gene using gene-specific primers in cv ICG 13942 of groundnut.

Lane M, DNA Marker; Lane C, Control plant DNA (-ve control-WT); Lane P, Plasmid pBAL2 (+ve control); Lanes 1-3, T₀ transgenic plants developed from DC; Lanes 4-5, T₀ transgenic plants developed from LL; Lanes 6-7, T₀ transgenic plants developed from CN explants.

Discussion

During the present study, we have attempted for optimization of factors affecting *Agrobacterium* mediated genetic transformation from individual explant of DC, LL, CN and found best transformation response in DC (18.13% Transient transformation efficiency) of groundnut cv. ICG 13942. There are various parameters which affect transformation efficiency as explored in groundnut.²¹ The highest transient transformation efficiency was observed for DC (18.13%) followed by LL (15.33%) and CN (12.32%) explants (Figure 3I) in the present investigation. Thus, the transformation efficiency is dependent upon the type of explant in groundnut and similar observations were also made in this crop: 2%,²² 16.5%²³ when DC was used as explants, 2.2%,²³ 11.3%²³ when LL was used as explants and 5.2%,²⁴ 8.5%²⁴ when used as CN explants.

Thus, we have achieved the increased level of transformation efficiencies in DC, LL and CN explants in comparison to earlier reports. The density of *Agrobacterium* suspension plays an important parameter. In the case of groundnut cv ICG 13942, the O.D₆₀₀ 1.0 gave maximum transient transformation efficiency 82.60% in DC, 74.25% in LL and 63.23% in CN explants while an O.D of 1.5 resulted in decreased transient transformation efficiency 67.43 % in DC, 58.63% in LL and 52.12% in CN explants (Figure 3A). Higher bacterial density in the medium may lead to severe necrosis of tissue.²⁵ Pre-culture is an important parameter for genetic transformation of plants because in this step, to reduce the wound stress and enhance the attachment of bacterial cells at the wound site by increasing the entry of T-DNA part into the host cells. Present study showed that, the Transient transformation efficiency (T.T.E) was high 69.11% in DC, 66.22% in LL, 64.92 in CN explants after 3 days of pre-culture period while it decreased after 5-11 days of pre culture. Similarly, the importance of pre-culture during transformation studies was also reported in peanut.²⁶ Co-cultivation is another important parameter for genetic transformation of plants because in this step, T-DNA is incorporated into plant genomic DNA.²⁷ Our results showed that, the T.T.E was high 62.22% in DC, 60.14% in LL and 57.23 in CN after 4 days while it decreased after 5-7 days of co cultivation (Figure 3c). Co-cultivation for 4 days was shown to be acceptable in groundnut for better transformation while the longer for co-cultivation does not help to increase transformation efficiency in different legumes.²⁸ It is reported that, secretion of Acetosyringone during wounding in plant cells induces the transcription of the virulence genes of *Agrobacterium*. Addition of Acetosyringone can improve transformation efficiency, in various monocots and some dicot plants which fail to secrete Acetosyringone.²⁷

However in our study, addition of 100 mg/L Acetosyringone showed that, the T.T.E was high at 76.54% in DC, 71.78% in LL, 68.21% in CN while decreased at 125-200mg/L. According to our observations 20 min infection period had shown the highest percentage of transformation efficiency DC (58.13%), LL (52.20%) CN (48.62%) while it decreased after 30 min (Figure 3d). An extension of infection period for more than 20 min, led to the explant abortion caused by bacterial contamination which reduced the transformation frequency.²⁹ *Agrobacterium tumefaciens* harboring pBAL2 plasmid carrying *nptIII* and *uidA* genes were used for genetic transformation of groundnut. The present instigation showed more than 90% of shoot induction response inhibition at 100 mg/L Kan in DC, CN explants and more than 85% at 75 mg/L Kan in LL explants (Figure 3e). Beyond that level of Kan, showed the bleaching and necrosis of the explants.

Similar findings were also reported in chickpea.³⁰ For the effect of cefotaxime 250mg/L shows the inhibiting effect of *Agrobacterium* in DC, LL and CN in groundnut cv ICG 13942. Coherent to our results, in the case of *Jatropha curcas*, 250mg/L Cefotaxime supported the callus induction and shoot regeneration in cotyledon explants and therefore, cefotaxime was used to inhibit *Agrobacterium* growth after co-cultivation with cotyledon explants.^{31,32} In castor also similar results were found that overgrowth of *Agrobacterium* was controlled by 250mg/l Cefotaxime during transformation of embryo axes and shoot apices.³³ According to our literature survey, this is the first report of *Agrobacterium* mediated genetic transformation of groundnut cv ICG 13942. Thus, the protocol developed during the present investigation can be used for genetic transformation experiments to introduce any novel gene into peanut.

Acknowledgements

We are grateful to University Grants Commission, New Delhi for financial assistance as Fellow under UGC-BSR-RFSMS (F.4-1/2006(BSR)/7-211/2009(BSR) dt.26-02-2013.). We thank Dr. HD. Upadhyaya ICRISAT, Patancheru, Hyderabad (TS), India for the germplasm (seeds) of cv ICG 13942 and Prof. N. Jayabalan Department of Plant Science, Bharatidasan University, Thiruchurappally, Tamil Nadu, India, for the gift of pBAL2.

Conflict of interest

The author declares there is no conflict of interest.

References

1. Enserink M. The peanut butter debate. *Science*. 2008;322(5898):36–38.
2. United States Department of Agriculture (USDA) Foreign Agricultural Service. Table 13 peanut area, yield, and production; USA; 2014.
3. Jaiwal PK, Singh RP. *Improvement strategies for Leguminosae Biotechnology*, 10th ed. *Focus on biotechnology*. Kluwer Academic Publisher, Dordrecht; 2003.
4. Popelka JC, Terryn N, Higgins TJV. Gene technology for grain legumes: Can it contribute to the food challenge in developing countries. *Plant Science*. 2004;167(2):195–206.
5. Dita MA, Rispaill N, Prats E, et al. Biotechnology approaches to overcome biotic and abiotic stress constraints in legumes. *Euphytica*. 2006;147(1-2):1–24.
6. Eapen S. Advances in development of transgenic pulse crops. *Biotechnology Advances*. 2008;26(2):162–168.
7. Zhang Z, Dermot CP, Mitra A. Factors affecting *Agrobacterium*-mediated transformation of common Bean (*Phaseolus vulgaris*). *J Amer Soc Hort Sci*. 1997;122(3):300-305.
8. Akbulut M, Yucel M, Oktem HA. Analysis and optimization of DNA delivery into Chickpea (*Cicer arietinum* L.) seedlings by *Agrobacterium tumefaciens*. *African Journal of Biotechnology*. 2008;7(8):1011-1017.
9. Somers DA, Samac DA, Olhoft M. Recent advances in legume transformation. *Plant Physiol*. 2003;131(3):892–899.
10. Kanyand M, Peterson CM, Prakash CS. The differentiation of emergences in to adventitious shoots in peanut (*Arachis hypogaea* L.). *Plant Sci*. 1997;126:87–95.
11. Venkatachalam P, Geeta N, Khandelwal A, et al. *Agrobacterium* - mediated genetic transformation and regeneration of transgenic plants from cotyledon explants of groundnut (*Arachis hypogaea* L.) via somatic embryogenesis. *Curr Sci*. 2000;78(9):1130–1136.

12. Livingstone DM, Birch RG. Plant regeneration and micro projectile mediated gene transfer in embryonic leaflets of peanut (*Arachis hypogaea* L.). *Australian Journal of Plant Physiology*. 1995;22(4):585–591.
13. Singsit C, Adang MJ, Lynch RE, et al. Expression of a *Bacillus thuringiensis* cry1A gene in transgenic peanut plants and its efficacy against lesser cornstalk borer. *Transgenic Res*. 1997;6(2):169–176.
14. Yang H, Singsit C, Wang A, et al. Transgenic peanut plants containing a nucleocapsid protein gene of tomato spotted wilt virus show divergent levels of gene expression. *Plant Cell Rep*. 1998;17:693–699.
15. Rajinikanth M, Srinivas P, Samatha T, et al. *In vitro* multiplication potential of cotyledonary node explants in Peanut (*Arachis hypogaea* L.). *Int J Agricult Scie and Res*. 2013;3(1):189–196.
16. Rajinikanth M, Ebrahimzadeh MH, Archana K, et al. *In vitro* direct organogenesis and plantlet formation from de-embryonated cotyledon explants of peanut (*Arachis hypogaea* L.) *Int J Pharma and Bio Sciences*. 2014;5(3)(B):481–490.
17. Rajinikanth M, Mahitha B, Rama Swamy N. Influence of silver nitrate on leaflet based direct regeneration in *Arachis hypogaea* L. *Vegetos*. 2015;28(1):63–69.
18. Murashige T, Skoog I. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiol*. 1962;15:473–497.
19. Gamborg OL, Miller RA, Ojima K. Nutrient requirements of suspension cultures of soya bean root cells. *Exp Cell Res*. 1968;50(1):151–158.
20. Jefferson RA, Kavanagh TA, Bevan MW. Gus fusion: β -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J*. 1987;6(13):3901–3907.
21. Venkatachalam P, Geetha N, Jayabalan N, et al. *Agrobacterium* mediated genetic transformation of groundnut (*Arachis hypogaea* L.) An assessment of factors affecting regeneration of transgenic plants. *J Plant Res*. 1998;111(4):565–577.
22. Freitas VGD, Lacorte C, Martins GS, et al. Identification of competent cells for *Agrobacterium* transformation and *in vitro* regeneration in peanut leaf and cotyledon explants. *R Bras Fisiol Veg*. 1997;157–167.
23. Mehta R, Radhakrishnan T, Kumar A, et al. Coat protein-mediated transgenic resistance of peanut (*Arachis hypogaea* L.) to peanut stem necrosis disease through *Agrobacterium*-mediated genetic transformation. *Indian J Virol*. 2013;24(2):205–213.
24. Beena MR, Tuli R, Gupta AD, et al. Transgenic peanut (*Arachis hypogaea* L.) plants expressing cry1EC and rice chitinase cDNA (Chi11) exhibit resistance against insect pest *Spodoptera litura* and fungal pathogen *Phaeoisariopsis personata*. *Transgenic Plant Journal*. 2008;2(2):157–164.
25. Li JF, Park E, von Arnim AG, et al. The FAST technique: a simplified *Agrobacterium*-based transformation method for transient gene expression analysis in seedlings of *Arabidopsis* and other plant species. *Plant Methods*. 2009;5:6.
26. Vasudevan A, Ganapathi A, Selvaraj N, et al. Factors influencing GUS expression in Cucumber (*Cucumis sativus* Linn). *Indian Journal of Biotechnology*. 2002;1:344–349.
27. Han J, Wang H, Ye H, et al. High efficiency of genetic transformation and regeneration of *Artemisia annua* L. Via *Agrobacterium tumefaciens*-mediated procedure. *Plant Science*. 2005;168(1):73–80.
28. Geetha N, Venkatachalam P, Rao GR. Factors influencing production of *Agrobacterium* mediated genetically transformed calli and plant regeneration in black gram (*Vigna mungo* L. Hepper). *Plant Tiss Cult*. 1997;7:149–152.
29. Bidney D, Seelonge C, Martich J, et al. Micro projectile bombardment of plant tissues increases transformation frequency by *Agrobacterium tumefaciens*. *Plant Molecular Biology*. 1992;18(2):301–313.
30. Kar S, Johnson TM, Nayak P, et al. Efficient transgenic plant regeneration through *Agrobacterium* mediated genetic transformation of chickpea (*Cicer arietinum* L.). *Plant Cell Rep*. 1996;16(1–2):32–37.
31. Li MR, Li HQ, Wu GJ. Study on factors influencing *Agrobacterium*-mediated transformation of *Jatropha curcas*. *Fen Zi Xi Bao Sheng Wu Xue Bao*. 2006;39(1):83–87.
32. Li LC, Qin GJ, Tsuge T, et al. Sporocyteless modulates YUCCA expression to regulate the development of lateral organs in *Arabidopsis*. *New Phytol*. 2008;179(3):751–764.
33. Sujatha M, Sailaja M. Stable genetic transformation of castor (*Ricinus communis* L.) via *Agrobacterium tumefaciens* - mediated gene transfer using embryo axes from mature seeds. *Plant Cell Rep*. 2005;23(12):803–810.