

Major insect pests of soybean - studies on population dynamics

Abstract

The investigation was carried out during *kharif* 2010-2011 and 2011-2012 in order to know the population dynamics of insect pests of soybean. The population of *Gesoniagemma*, *Spodopteralitura*, *elicoverpaarmigera*, *Aproaeremamodicella*, *Obereopsis brevis* and *Melanagromyza sojajae* were ranged from 0-4, 0-2.2, 0-2.4, 0-0.3, 5.20-19.66, and 14.9-27.2 and it was 3.9-20.4, 6.8-26.2, 3.2-16.0, 0-5.5, 2.9-24.7 and 10-25.7 during 2011-12, respectively. In conclusion leaf miner (*A. modicella*), defoliators, stem fly (*M. sojajae*) and girdle beetle (*O. brevis*) were found to be major pests of soybean.

Keywords: Soybean, Insect pests, population dynamics, weather factors, semilooper, spodoptera, helicoverpa, leaf miner, stem pests

Volume 9 Issue 1 - 2019

AA Motaphale, BA Aglave, BB Bhosale

Vasant Rao Naik Marathwada Agricultural University, India

Correspondence: Motaphale AA, Vasant Rao Naik Marathwada Agricultural University, Vasmat Road, Kali Kaman, Parbhani, MS 431402, India, Email ashvini@gmail.com

Received: September 24, 2018 | **Published:** January 07, 2018

Introduction

The population dynamics of major defoliator and stem pests, infesting soybean in relation to weather parameters were envisaged to have insight into the predisposing ecological factors for occurrence of the pest. The climatic factors such as temperature, rainfall and humidity usually act in a density independent manner influencing insect to a greater or lesser extent. The experiment was conducted in pesticide free conditions. The population dynamics of pests of soybean was studied in different meteorological weeks. The population dynamics of leaf miner (*Aproaeremamodicella* Deventer), stemfly (*Melanagromyza sojajae*, Zehntner), girdle beetle (*Obereopsis Brevis* Swedenbord), and some defoliators was worked out by recording their observations weekly.

Material and methods

Field experiments were conducted during *kharif* 2010-2011 and 2011-12, in the field of Department of Agricultural Entomology, College of Agriculture, Marathwada Krishi Vidyapeeth, Parbhani (Maharashtra). The line sowing of seed (MAUS-71) was done by maintaining 45cm distance between two rows and 5cm between two plants. The number of defoliating larvae/meter row length (mrl) was recorded at three places on weekly basis and leaf damage due to defoliators was worked out in terms of per cent defoliation. Number of larvae/plant in 10 plants, total number of leaflets and damage leaflets due to leaf miner was recorded and percentage was worked out. Total number of plants and number of plants succumbed to stem fly infestation/mrl at 3 places per plot on 7 and 10 days after sowing were recorded and expressed in per cent. Plant height and length of stem tunneled in 10 plants at physiological maturity was observed and expressed in percentage. One meter row length area was marked at five places and total number of plants and girdled plants by girdle beetle per mrl were recorded.

Result and discussion

The population of *Gesoniagemma* (0.8 and 7.4/mrl) initiated in 29th and 30th MW with its peak (4.0 and 20.40 larvae per mrl) in 35th and 34th MW during 2010-11 and 2011-12, respectively Table 1 & Table 2. At the time of its peak incidence, the weather parameters *viz.*, rainfall, rainy days, maximum temperature, minimum temperature,

before noon relative humidity, afternoon relative humidity were 91.4 and 61.3mm, 5°C and 4.0 days, 30.6 and 29.9°C, 22.7°C and 21.9°C, 92 and 92 per cent, 73 and 80 per cent, respectively. The incidence of semilooper was below ETL throughout the *kharif* season and a maximum of 2.66 larvae/mrl was observed in last week of August. Population of green semi looper was high during *kharif* 2011-2012, the incidence started in 35th MW by recording 19.33 larvae/mrl and recorded a peak in next week with 28 larvae/mrl.¹ *Spodopteralitura* (Fabricius) incidence was low (0 to 2.2 larvae/mrl) due to heavy rainfall during 2010-2011 and from 6.80 to 26.2 larvae per mrl during 2011-2012. The highest level of population was observed in 36th and 34th MW when weather factors *viz.*, rainfall, rainy days, maximum temperature, minimum temperature, before noon relative humidity, afternoon relative humidity were 51.1 and 61.3 mm, 5.0 and 4.0 days, 29.0 and 29.9°C, 22.3°C and 21.9°C, 94 and 92 per cent, 76 and 80 per cent, respectively. More or less similar results were observed by Chechani et al.,² who noticed tobacco caterpillar during 34th meteorological week (20 to 26 August, 1997). They observed two distinct peaks, the first peak with a mean population of 1.53 larvae/plant in 36th meteorological week (3 to 9 September 1997) and the second peak with a mean population of 1.53 larvae/plant during 38th meteorological week (17 to 23 September 1997), when average temperature and relative humidity were 26.3°C and 26.65°C and 77 and 77.5 per cent, respectively. The pest population showed a positive correlation with temperature and relative humidity. Sreenivas et al.,³ also reported peak incidence during 35th MW (27th August to 2nd September, 2002) in case of tobacco caterpillar, supporting the present results. The abundance of *Helicoverpaarmigera* (Hubner) was observed to be 2.40 and 16.0 larvae/mrl on soybean in 34th MW during both the seasons, when rainfall, rainy days, maximum temperature, minimum temperature, before noon relative humidity, afternoon relative humidity were 27.8 and 61.3mm, 3.0 and 4 days, 30.4 and 29.9°C, 23.1 and 21.9°C, 90 and 92 per cent, 68 and 80 per cent, respectively. The population of *Aproaeremamodicella* (Devender) was recorded on soybean between 27th to 35th (0 to 0.3 larvae per plant) and 30th to 37th meteorological weeks (0 to 5.5 larvae per plant) during 2010-2011 and 2011-2012, respectively. At the maximum level of larval population of *A. modicella*, rainfall, rainy days, maximum temperature, minimum temperature, before noon relative humidity, afternoon relative humidity were 91.4mm and 50mm, 5.0 days and

2.0 days, 30.6°C and 32.0°C, 22.7°C and 22.2°C, 92 per cent and 87 per cent, 73 and 66 per cent during 2010-2011 and 2011-2012, respectively. These results are in line with the results reported by Magar⁴ who reported the highest population of *A. modicella* to the extent of 19.8 larvae per quadrat on soybean in 38th meteorological week when the maximum temperature, minimum temperature, before noon relative humidity, afternoon relative humidity, rainfall and number of rainy days were 27.85°C, 20.92°C, 77.00 per cent, 88.83 per cent, 40.73mm and 3 days, respectively. Behera et al.,⁵ recorded

that the larval activity of leaf-miner started in third week of July and reached its peak during the first week of August, when the mean larval population ranged from 1.0 to 12.3 larvae per five plants. The population of leaf miner maintained from third week of August to third week of September with slight fluctuation of 1.4 to 2.1 larvae per five plants. Bidgire⁶ recorded the first appearance of leaf miner (0.60 larvae/plant) in the last week of July (30th MW) on groundnut crop during *kharif* 1988.

Table 1 Population dynamics of insect pests damaging soybean in relation to weather parameters during kharif 2010-11

MW	Period	No. of <i>G. gemma</i> larvae/mrl	No. of <i>S. litura</i> larvae/mrl	No. of <i>H. armigera</i> larvae/mrl	No. of <i>A. modicella</i> larvae/plant	% Infestation due to <i>O. brevis</i>	% Infestation due to <i>M. sojae</i>
27	02-08 July	0	0	0	0	5.2	12.7
28	09-15 July	0	0	0	0	7.9	14.9
29	16-22 July	0.8	0.8	0.2	0.1	10.8	19.2
30	23-29 July	1.4	0.6	0.2	0.3	12.4	19.9
31	30-05 Aug	1.8	0.6	0.4	0	12	18.8
32	06-12 Aug	2	0.2	1	0	15.7	21.8
33	13-19 Aug	0.8	0	1.6	0.2	17.9	23
34	20-26 Aug	1.8	0.2	2.4	0.2	18	25.9
35	27-02 Sept	4	0.8	1.8	0.3	18.2	26
36	03-09 Sep	2.8	2.2	1.2	0.2	18.4	27.2
37	10-16 Sep	2.2	1.8	3.1	0.1	18.6	25.1
38	17-23 Sep	1	0.8	0.1	0.2	19	25.3
39	24-30 Sep	0.8	0.6	0	0.1	19.6	23.3
40	01-07 Oct.	0.2	0.4	0	0.1	19.66	23

mrl, meter row length

Table 2 Population dynamics of insect pests damaging soybean in relation to weather parameters during kharif 2011-12

MW	Period	No. of <i>G. gemma</i> larvae/mrl	No. of <i>S. litura</i> larvae/mrl	No. of <i>H. armigera</i> larvae/mrl	No. of <i>A. modicella</i> larvae/plant	% Infestation due to <i>O. brevis</i>	% Infestation due to <i>M. sojae</i>
30	23-29 July	7.4	15.9	4.9	0	2.9	10.9
31	30-05 Aug	12.9	11.3	5.7	1.7	8.2	15.9
32	06-12 Aug	5.6	9.9	3.2	1	3.3	10
33	13-19 Aug	7.8	14.9	10.2	4.1	5.9	18.8
34	20-26 Aug	20.4	26.2	16	4.3	15.8	25.7
35	27-02 Sept	17.9	19.8	12.1	3.7	16	21.9
36	03-09 Sep	15	18.8	7.9	4	14.9	21.4
37	10-16 Sep	19.9	20.4	9.9	5.5	20.7	19.7
38	17-23 Sep	10	19.7	10	4.3	20.1	20.4
39	24-30 Sep	8	16.5	5.8	3.3	22.2	18.8
40	01-07 Oct.	3.9	6.8	4.8	2.8	24.7	20.2

mrl, meter row length

The leaf miner population lasted up to second week of September (36th MW) and recorded maximum population (1.15 larvae/plant) in the middle of August. Sayyad⁷ observed first incidence of leaf miner

(0.33 larvae per plant) in fourth week of July (29th MW). The highest population (2.83 larvae per plant) was observed in the third week of August (33rd MW). Later on population of leaf miners decreased slowly

and was 0.23 larvae per plant in third week of September (37th MW). Number of infested plants due to *Melanagromyza sojiae* (Zehntner) was recorded on soybean between 27th to 38th MW (12.70 to 27.2 per cent) and 30th and 40th MW from 10.90 to 25.70 per cent during 2010-2011 and 2011-2012, respectively. At the maximum level of infestation, rainfall, rainy days, maximum temperature, minimum temperature, before noon relative humidity, afternoon relative humidity were 51.4 and 16.4mm, 5 days and 1 day, 29.0°C and 33.1°C, 22.3°C and 20.8°C, 94 and 84 per cent, 76 and 44 per cent, respectively during 2010-2011 and 2011-2012. Ali⁸ reported higher infestation of *Melanagromyza* sp. on soybean during *kharif* than rabi season. Verma et al.,⁹ reported the infestation of *M. sojaeonurdbeanto* to an extent of 22 per cent in 3rd week of July to 70 per cent in 3rd week of August. The infestation of stemfly was noticed from last week of August and from middle of September, the damage was above the ETL throughout the season with 31.40 per cent damaged plants in 3rd week of October.¹ The maximum population of *Oberea brevis* (Sweden bord) (19.66 per cent and 24.7 per cent) was recorded on soybean in 33rd and 40th meteorological week during 2010-2011 and 2011-12, respectively, when rainfall, rainy days, maximum temperature, minimum temperature, before noon relative humidity, afternoon relative humidity were 1.4 and 16.4 mm, 0 and 1 days, 32.8°C and 33.1°C, 21.1°C and 20.8°C, 81 and 84 per cent, 48 and 44 per cent, respectively. Earlier results reported by Rai and Patel¹⁰ revealed that *O. brevis* appeared on soybean when maximum temperature, minimum temperature and relative humidity were 27.60°C, 24.70°C and 84.50 per cent, respectively. Incidence of girdle beetle started in 1st week of August, increased gradually and reached to peak of 16.43 per cent during last week of October, crossing the ETL. During 2011 incidence of girdle beetle was more. It started in 32nd MW (0.83 per cent), increased gradually and reached to peak of 35.92 per cent in 40th and 41st MW.

Acknowledgments

None.

Conflicts of interest

The authors declared there are no conflicts of interest.

References

1. Anonymous. [http://www.google.com/soy/stat/world soybean production](http://www.google.com/soy/stat/world%20soybean%20production); 2010.
2. Chechani VK, Joshi FL, Sharma US, et al. Incidence and chemical control of stemfly *Melanagromyza sojiae* (Zehntner) on soybean. *Indian J Appl Ent.* 2002;14:9–12.
3. Sreenivas K, Dhurve SB, Bisane KD, et al. Biology of *Spilosoma obliqua* Walker and seasonal incidence of soybean pest. *J Oilseeds Res.* 2007;24(1):218–220.
4. Magar SS. *Field life-tables and population dynamics of major pests of soybean*, M.Sc. thesis submitted to Marathwada Agricultural University, Parbhani: 2006.
5. Behera PK, Patnaik HP, Senapati B. Leaf miner *Bilobatasubsecivell* Zehnter (*Gelechiidae*: Lepidoptera) incidence on soybean cultivars in North Orisa. *Orissa J Agric Res.* 1989;2(3-4):227–229.
6. Bidgire US. *Management of some important insect pests of groundnut*. Ph.D. Thesis, MAU, Parbhani (M.S.), India: 1995.
7. Sayyad HS. Efficacy of some newer insecticides against major insect pests of soybean, *Glycine max* (L.) Merrill. M.Sc.(Agri.), Thesis, MAU, Parbhani: 2001.
8. Ali MI. Field observation on the attack of stemfly on different cultivars of soybean (Bangladesh). *Bangladesh J agric Sci* 1981;8(1):83–84.
9. Verma SN, Singh OP. Rice bean, *Vigna umbellata* a new host of girdle beetle (*Obereopsis brevis* S.). *Indian J Plant Prot.* 1989;19:104.
10. Rai RK, Patel RK. Girdle beetle, *Obereopsis brevis* Swed incidence in *kharif* soybean. *Orissa Journal of Agricultural Research.* 1990;3(2):163–165.