

On modeling of lifetimes data using exponential and lindley distributions

Abstract

In this paper, firstly the nature of exponential and Lindley distributions have been studied using different graphs of their probability density functions and cumulative distribution functions. The expressions for the index of dispersion for both exponential and Lindley distributions have been obtained and the conditions under which the exponential and Lindley distributions are over-dispersed, equi-dispersed, and under-dispersed has been given. Several real lifetimes data-sets has been fitted using exponential and Lindley distributions for comparative study and it has been shown that in some cases exponential distribution provides better fit than the Lindley distribution whereas in other cases Lindley distribution provides better fit than the exponential distribution.

Keywords: exponential distribution, lindley distribution, index of dispersion, estimation of parameter, goodness of fit

Volume 2 Issue 5 - 2015

Rama Shanker,¹ Hagos Fesshaye,² Sujatha Selvaraj³

¹Department of Statistics, Eritrea Institute of Technology, Eritrea

²Department of Economics, College of Business and Economics, Eritrea

³Department of Banking and Finance, Jimma University, Ethiopia

Correspondence: Rama Shanker, Department of Statistics, Eritrea Institute of Technology, Asmara, Eritrea, Email shankerrama2009@gmail.com

Received: June 15, 2015 | **Published:** June 26, 2015

Introduction

The time to the occurrence of some event is of interest for some populations of individuals in every field of knowledge. The event may be death of a person, failure of a piece of equipment, development of (or remission) of symptoms, health code violation (or compliance). The times to the occurrences of events are known as “lifetimes” or “survival times” or “failure times” according to the event of interest in the fields of study. The statistical analysis of lifetime data has been a topic of considerable interest to statisticians and research workers in areas such as engineering, medical and biological sciences. Applications of lifetime distributions range from investigations into the endurance of manufactured items in engineering to research involving human diseases in biomedical sciences.

There are a number of continuous distributions for modeling lifetime data such as exponential, Lindley, gamma, lognormal and Weibull. The exponential, Lindley and the Weibull distributions are more popular in practice than the gamma and the lognormal distributions because the survival functions of the gamma and the lognormal distributions cannot be expressed in closed forms and both require numerical integration. Both exponential and Lindley distributions are of one parameter and the Lindley distribution has advantage over the exponential distribution that the exponential distribution has constant hazard rate and mean residual life function whereas the Lindley distribution has increasing hazard rate and decreasing mean residual life function.

In this paper, firstly the nature of exponential and Lindley distribution has been studied by drawing different graphs for probability densities and cumulative distribution functions for the same values of parameter. Several examples of lifetimes data-sets from different fields of knowledge has been considered and an attempt has been made to study the goodness-of-fit for both exponential and Lindley distributions to see the superiority of one over the other.

Exponential and lindley distributions

Exponential distribution

The exponential distribution was the first widely used lifetime

distribution model in areas ranging from studies on the lifetimes of manufactured items¹⁻³ to research involving survival or remission times in chronic diseases.⁴ The main reason for its wide applicability as lifetime model is partly because of the availability of simple statistical methods for it² and partly because it appeared suitable for representing the lifetimes of many things such as various types of manufactured items.¹

Let T be a continuous random variable representing the lifetimes of individuals in some population and following exponential distribution. The probability density function (p.d.f.), cumulative distribution function (c.d.f.), survival function, hazard function, and mean residual life function of T , respectively, are given by

$$f(t) = \theta e^{-\theta t}; \quad \theta > 0, t > 0$$

$$F(t) = 1 - e^{-\theta t}; \quad \theta > 0, t > 0$$

$$S(t) = 1 - F(t) = e^{-\theta t}$$

$$h(t) = \frac{f(t)}{1 - F(t)} = \frac{f(t)}{S(t)} = \theta$$

$$m(t) = \frac{1}{\theta}$$

Lindley distribution

Lindley distribution is a mixture of exponential (θ) and gamma ($2, \theta$) distributions with mixing proportion $\frac{\theta}{\theta+1}$ and is given by Lindley (1958) in the context of Bayesian Statistics as a counter example of fiducial Statistics. Let T be a continuous random variable representing the lifetimes of individuals in some population and following Lindley distribution. The probability density function (p.d.f.), cumulative distribution function (c.d.f.), survival function, hazard function, and mean residual life function of T , respectively, are given by

$$f(t) = \frac{\theta^2}{\theta+1} (1+t) e^{-\theta t}; \theta > 0, t > 0$$

$$F(t) = 1 - \frac{\theta+1+\theta t}{\theta+1} e^{-\theta t}; \theta > 0, t > 0$$

$$S(t) = 1 - F(t) = \frac{\theta+1+\theta t}{\theta+1} e^{-\theta t}$$

$$h(t) = \frac{f(t)}{1-F(t)} = \frac{f(t)}{S(t)} = \frac{\theta^2(1+t)}{\theta+1+\theta t}$$


$$m(t) = \frac{\theta+2+\theta t}{\theta(\theta+1+\theta t)}$$

The Lindley distribution has been extensively studied and

generalized by many researchers such as⁵⁻¹² are among others. A discrete version of the Lindley distribution has been obtained by¹³ and¹⁴ has obtained the Lindley mixture of Poisson distribution.

The graphs of the probability densities functions of exponential and Lindley distributions are presented for different values of parameter and shown in Figure 1. The graphs of the cumulative distribution functions of exponential and Lindley distributions are presented for different values of parameter and are shown in Figure 2.

The expressions for coefficient of variation (C.V.), coefficient of Skewness ($\sqrt{\beta_1}$), coefficient of Kurtosis (β_2), and index of dispersion of exponential and Lindley distributions are summarized in the following Table 1. It can be easily verified that the Lindley distribution is over- dispersed ($\mu < \sigma^2$), equi-dispersed ($\mu = \sigma^2$) and under-dispersed ($\mu > \sigma^2$) for $\theta < (=) > \theta^* = 1.170086487$ respectively, whereas as exponential distribution is over- dispersed ($\mu < \sigma^2$), equi-dispersed ($\mu = \sigma^2$) and under- dispersed ($\mu > \sigma^2$) for $\theta < (=) > \theta^* = 1$ respectively







Figure 1 Graphs of the p.d.f. of exponential and Lindley distributions (left hand side graphs are for exponential and right hand side graphs are for Lindley).






Figure 2 Graphs of the c.d.f. of exponential and Lindley distributions (left hand side graphs are for exponential and right hand side graphs are for Lindley).

Applications

The exponential and Lindley distribution has been fitted to a number of real lifetime data - sets to tests their goodness of fit. Goodness of fit tests for fifteen real lifetime data- sets has been presented here.

In order to compare exponential and Lindley distributions, $-2 \ln L$, AIC (Akaike Information Criterion), AICC (Akaike Information Criterion Corrected), BIC (Bayesian Information Criterion), K-S Statistics (Kolmogorov-Smirnov Statistics) for all fifteen real lifetime data- sets have been computed. The formulae for computing AIC, AICC, BIC, and K-S Statistics are as follows:

$$AIC = -2 \ln L + 2k, AICC = AIC + \frac{2k(k+1)}{(n-k-1)},$$

$$BIC = -2 \ln L + k \ln L \quad \text{and} \quad D = \text{Sup}_x |F_n(x) - F_0(x)|$$

, where k the number of parameters, n is the sample size and $F_n(x)$ is the empirical distribution function. The best distribution corresponds to lower $-2 \ln L$, AIC, AICC, BIC, and K-S statistics.

The fittings of exponential and Lindley distributions are based on maximum likelihood estimates (MLE). Let t_1, t_2, \dots, t_n be a random sample of size n from exponential distribution. The likelihood function L , and the log likelihood function $\ln L$ of exponential distribution are given by $L = \theta^n e^{-n\theta\bar{T}}$ and $\ln L = n \ln \theta - n\theta\bar{T}$. The MLE $\hat{\theta}$ of the parameter θ of exponential distribution is the solution of the equation $\frac{d \ln L}{d\theta} = 0$ and is given by $\hat{\theta} = \frac{1}{\bar{T}}$, where \bar{T} is the sample mean.

Table 1

Exponential Distribution	Lindley Distribution
$C.V. = \frac{\sigma}{\mu_1} = 1$	$C.V. = \frac{\sigma}{\mu_1} = \frac{\sqrt{\theta^2 + 4\theta + 2}}{\theta + 2}$
$\sqrt{\beta_1} = 2$	$\sqrt{\beta_1} = \frac{2(\theta^3 + 6\theta^2 + 6\theta + 2)}{(\theta^2 + 4\theta + 2)^{3/2}}$
$\beta_2 = 9$	$\beta_2 = \frac{3(3\theta^4 + 24\theta^3 + 44\theta^2 + 32\theta + 8)}{(\theta^2 + 4\theta + 2)^2}$
Index of dispersion $\gamma = \frac{\sigma^2}{\mu_1'} = \frac{1}{\theta}$	Index of dispersion $\gamma = \frac{\sigma^2}{\mu_1'} = \frac{\theta^2 + 4\theta + 2}{\theta(\theta^2 + 3\theta + 2)}$

Let t_1, t_2, \dots, t_n be a random sample of size n from Lindley distribution. The likelihood function, L and the log likelihood function, $\ln L$ of Lindley distribution are given by $L = \left(\frac{\theta^2}{\theta+1}\right)^n \prod_{i=1}^n (1+t_i)e^{-n\theta\bar{T}}$ and $\ln L = n \ln \left(\frac{\theta^2}{\theta+1}\right) + \sum_{i=1}^n \ln(1+t_i) - n\theta\bar{T}$. The MLE $\hat{\theta}$ of the parameter θ of Lindley distribution is the solution of the equation $\frac{d \ln L}{d\theta} = 0$ and is given by $\hat{\theta} = \frac{-(\bar{T}-1) + \sqrt{(\bar{T}-1)^2 + 8\bar{T}}}{2\bar{T}}$; $\bar{T} > 0$, where \bar{T} is the sample mean. It was shown by Ghitany *et al.*,⁵ that the estimator $\hat{\theta}$ of Lindley distribution is positively biased, consistent and asymptotically normal.

From above table it is obvious that the fittings of Lindley distribution is better than the exponential distribution in Datasets 1-6,12,14,15. Whereas the fittings of exponential distribution is better than the Lindley distribution in Datasets 7-11,13 (Table 2).

Conclusion

In this paper we have tried to find the suitability of exponential and Lindley distributions for modeling real lifetimes data. It has been observed that neither exponential distribution nor Lindley distribution is appropriate for modeling real lifetime data in all cases. As per the nature of the data related with over-dispersion, equi-dispersion, and under-dispersion, in some cases exponential is better than Lindley while in other cases Lindley is better than exponential. Further, the decision about the suitability of exponential and Lindley for modeling real lifetime data depends on the nature of the data. Of course, Lindley is more flexible than exponential but exponential has some advantage over Lindley due to its simplicity.

Data Set 1 The data set represents the strength of 1.5cm glass fibers measured at the National Physical Laboratory, England. Unfortunately, the units of measurements are not given in the paper, and they are taken from Smith and Naylor¹⁵

0.55	0.93	1.25	1.36	1.49	1.52	1.58	1.61	1.64	1.68	1.73	1.81	2.00
0.74	1.04	1.27	1.39	1.49	1.53	1.59	1.61	1.66	1.68	1.76	1.82	2.01
0.77	1.11	1.28	1.42	1.50	1.54	1.60	1.62	1.66	1.69	1.76	1.84	2.24
0.81	1.13	1.29	1.48	1.50	1.55	1.61	1.62	1.66	1.70	1.77	1.84	0.84
1.24	1.30	1.48	1.51	1.55	1.61	1.63	1.67	1.70	1.78	1.89		

Data Set 2 The data is given by Birnbaum and Saunders¹⁶ on the fatigue life of 6061 –T6 aluminum coupons cut parallel to the direction of rolling and oscillated at 18 cycles per second. The data set consists of 101 observations with maximum stress per cycle 31,000 psi. The data () are presented below (after subtracting 65)

5	25	31	32	34	35	38	39	39	40	42	43	43
43	44	44	47	47	48	49	49	49	51	54	55	55
55	56	56	56	58	59	59	59	59	59	63	63	64
64	65	65	65	66	66	66	66	66	67	67	67	68
69	69	69	69	71	71	72	73	73	73	74	74	76
76	77	77	77	77	77	77	79	79	80	81	83	83
84	86	86	87	90	91	92	92	92	92	93	94	97
98	98	99	101	103	105	109	136	147				

Data Set 3 The data set is from Lawless.¹⁷ The data given arose in tests on endurance of deep groove ball bearings. The data are the number of million revolutions before failure for each of the 23 ball bearings in the life tests and they are:

17.88	28.92	33.00	41.52	42.12	45.60	48.80	51.84	51.96	54.12	55.56	67.80
68.44	68.64	68.88	84.12	93.12	98.64	105.12	105.84	127.92	128.04	173.40	

Data Set 4 The data is from Picciotto¹⁸ and arose in test on the cycle at which the Yarn failed. The data are the number of cycles until failure of the yarn and they are:

86	146	251	653	98	249	400	292	131	169	175	176	76
264	15	364	195	262	88	264	157	220	42	321	180	198
38	20	61	121	282	224	149	180	325	250	196	90	229
166	38	337	65	151	341	40	40	135	597	246	211	180
93	315	353	571	124	279	81	186	497	182	423	185	229
400	338	290	398	71	246	185	188	568	55	55	61	244
20	284	393	396	203	829	239	236	286	194	277	143	198
264	105	203	124	137	135	350	193	188				

Data Set 5 This data represents the survival times (in days) of 72 guinea pigs infected with virulent tubercle bacilli, observed and reported by Bjerkedal¹⁹

12	15	22	24	24	32	32	33	34	38	38	43	44
48	52	53	54	54	55	56	57	58	58	59	60	60
60	60	61	62	63	65	65	67	68	70	70	72	73
75	76	76	81	83	84	85	87	91	95	96	98	99
109	110	121	127	129	131	143	146	146	175	175	211	233
258	258	263	297	341	341	376						

Data Set 6 This data is related with behavioral sciences, collected by N Balakrishnan et al.,²⁰ The scale “General Rating of Affective Symptoms for Preschoolers (GRASP)” measures behavioral and emotional problems of children, which can be classified with depressive condition or not according to this scale. A study conducted by the authors in a city located at the south part of Chile has allowed collecting real data corresponding to the scores of the GRASP scale of children with frequency in parenthesis, which are:

19(16)	20(15)	21(14)	22(9)	23(12)	24(10)	25(6)	
26(9)	27(8)	28(5)	29(6)	30(4)	31(3)	32(4)	
33	34	35(4)	36(2)	37(2)	39	42	44

Data Set 7 The data set reported by Efron²¹ represent the survival times of a group of patients suffering from Head and Neck cancer disease and treated using radiotherapy (RT)

6.53	7	10.42	14.48	16.10	22.70	34	41.55	42	45.28	49.40	53.62	63
64	83	84	91	108	112	129	133	133	139	140	140	146
149	154	157	160	160	165	146	149	154	157	160	160	165
173	176	218	225	241	248	273	277	297	405	417	420	440
523	583	594	1101	1146	1417							

Data Set 8 The data set reported by Efron²¹ represent the survival times of a group of patients suffering from Head and Neck cancer disease and treated using a combination of radiotherapy and chemotherapy (RT+CT)

12.20	23.56	23.74	25.87	31.98	37	41.35	47.38	55.46	58.36	63.47	68.46	78.26
74.47	81.43	84	92	94	110	112	119	127	130	133	140	146
155	159	173	179	194	195	209	249	281	319	339	432	469
519	633	725	817	1776								

Data set 9 This data set represents remission times (in months) of a random sample of 128 bladder cancer patients reported in Lee and Wang.²²

0.08	2.09	3.48	4.87	6.94	8.66	13.11	23.63	0.20	2.23	3.52	4.98	6.97
9.02	13.29	0.40	2.26	3.57	5.06	7.09	9.22	13.80	25.74	0.50	2.46	3.64
5.09	7.26	9.47	14.24	25.82	0.51	2.54	3.70	5.17	7.28	9.74	14.76	6.31
0.81	2.62	3.82	5.32	7.32	10.06	14.77	32.15	2.64	3.88	5.32	7.39	10.34
14.83	34.26	0.90	2.69	4.18	5.34	7.59	10.66	15.96	36.66	1.05	2.69	4.23
5.41	7.62	10.75	16.62	43.01	1.19	2.75	4.26	5.41	7.63	17.12	46.12	1.26
2.83	4.33	5.49	7.66	11.25	17.14	79.05	1.35	2.87	5.62	7.87	11.64	17.36
1.40	3.02	4.34	5.71	7.93	11.79	18.10	1.46	4.40	5.85	8.26	11.98	19.13
1.76	3.25	4.50	6.25	8.37	12.02	2.02	3.31	4.51	6.54	8.53	12.03	
20.28	2.02	3.36	6.76	12.07	21.73	2.07	3.36	6.93	8.65	12.63	22.69	

Data Set 10 This data set is given by Linhart and Zucchini,²³ which represents the failure times of the air conditioning system of an airplane

23	261	87	7	120	14	62	47	225	71	246	21	42
20	5	12	120	11	3	14	71	11	14	11	16	90
1	16	52	95									

Data Set 11 This data set used by Bhaumik et al.,²⁴ is vinyl chloride data obtained from clean upgradient monitoring wells in mg/l

5.1	1.2	1.3	0.6	0.5	2.4	0.5	1.1	8	0.8	0.4	0.6	0.9
0.4	2	0.5	5.3	3.2	2.7	2.9	2.5	2.3	1	0.2	0.1	0.1
1.8	0.9	2	4	6.8	1.2	0.4	0.2					

Data set 12 This data set represents the waiting times (in minutes) before service of 100 Bank customers and examined and analyzed by Ghitany et al.,⁵ for fitting the Lindley²⁵ distribution

0.8	0.8	1.3	1.5	1.8	1.9	1.9	2.1	2.6	2.7	2.9	3.1	3.2
3.3	3.5	3.6	4.0	4.1	4.2	4.2	4.3	4.3	4.4	4.4	4.6	4.7
4.7	4.8	4.9	4.9	5	5.3	5.5	5.7	5.7	6.1	6.2	6.2	6.2
6.3	6.7	6.9	7.1	7.1	7.1	7.1	7.4	7.6	7.7	8	8.2	8.6
8.6	8.6	8.8	8.8	8.9	8.9	9.5	9.6	9.7	9.8	10.7	10.9	11
11	11.1	11.2	11.2	11.5	11.9	12.4	12.5	12.9	13	13.1	13.3	13.6
13.7	13.9	14.1	15.4	15.4	17.3	17.3	18.1	18.2	18.4	18.9	19	19.9
20.6	21.3	21.4	21.9	23.0	27	31.6	33.1	38.5				

Data Set 13 This data is for the times between successive failures of air conditioning equipment in a Boeing 720 airplane, Proschan²⁶

74	57	48	29	502	12	70	21	29	386	59	27	153
26	326											

Data set 14 This data set represents the lifetime's data relating to relief times (in minutes) of 20 patients receiving an analgesic and reported by Gross and Clark²⁷

1.1	1.4	1.3	1.7	1.9	1.8	1.6	2.2	1.7	2.7	4.1	1.8	1.5
1.2	1.4	3	1.7	2.3	1.6	2						

Data Set 15 This data set is the strength data of glass of the aircraft window reported by Fuller et al.,²⁸

18.83	20.8	21.657	23.03	23.23	24.05	24.321	25.5	25.52	25.8	26.69	26.77
26.78	27.05	27.67	29.9	31.11	33.2	33.73	33.76	33.89	34.76	35.75	35.91
36.98	37.08	37.09	39.58	44.045	45.29	45.381					

Table 2 MLE's, -2lnL,AIC,AICC, BIC and K-S Statistics of the fitted distributions of data sets 1-15

	Model	Parameter estimate	-2ln L	AIC	AICC	BIC	K-S Statistic
Data 1	Lindley	0.996116	162.56	164.56	164.62	166.7	0.371
	Exponential	0.663647	177.66	179.66	179.73	181.8	0.402
Data 2	Lindley	0.028859	983.11	985.11	985.15	987.71	0.242
	Exponential	0.014635	1044.87	1046.87	1046.91	1049.48	0.357
Data 3	Lindley	0.027321	231.47	233.47	233.66	234.61	0.149
	Exponential	0.013845	242.87	244.87	245.06	246.01	0.263
Data 4	Lindley	0.00897	1251.34	1253.34	1253.38	1255.95	0.098
	Exponential	0.004505	1280.52	1282.52	1282.56	1285.12	0.19
Data 5	Lindley	0.019841	789.04	791.04	791.1	793.32	0.133
	Exponential	0.010018	806.88	808.88	808.94	811.16	0.198
Data 6	Lindley	0.077247	1041.64	1043.64	1043.68	1046.54	0.448
	Exponential	0.04006	1130.26	1132.26	1132.29	1135.16	0.525
Data 7	Lindley	0.008804	763.75	765.75	765.82	767.81	0.245
	Exponential	0.004421	744.87	746.87	746.94	748.93	0.166
Data 8	Lindley	0.00891	579.16	581.16	581.26	582.95	0.219
	Exponential	0.004475	564.02	566.02	566.11	567.8	0.145
Data 9	Lindley	0.196045	839.06	841.06	841.09	843.91	0.116
	Exponential	0.106773	828.68	830.68	830.72	833.54	0.077
Data 10	Lindley	0.033021	323.27	325.27	325.42	326.67	0.345
	Exponential	0.016779	305.26	307.26	307.4	308.66	0.213
Data 11	Lindley	0.823821	112.61	114.61	114.73	116.13	0.133
	Exponential	0.532081	110.91	112.91	113.03	114.43	0.089

Table Continued

	Model	Parameter estimate	-2ln L	AIC	AICC	BIC	K-S Statistic
Data 12	Lindley	0.186571	638.07	640.07	640.12	642.68	0.058
	Exponential	0.101245	658.04	660.04	660.08	662.65	0.163
Data 13	Lindley	0.01636	181.34	183.34	183.65	184.05	0.386
	Exponential	0.008246	173.94	175.94	176.25	176.65	0.277
Data 14	Lindley	0.816118	60.5	62.5	62.72	63.49	0.341
	Exponential	0.526316	65.67	67.67	67.9	68.67	0.389
Data 15	Lindley	0.062988	253.99	255.99	256.13	257.42	0.333
	Exponential	0.032455	274.53	276.53	276.67	277.96	0.426

Acknowledgement

None.

Conflict of interest

No conflict of interest

References

- Davis DJ. An analysis of some failure data. *Journal of American Statistical Association*. 1952;47(258):113–150.
- Epstein B, Sobel M. Life testing. *Journal of American Statistical Association*. 1953;48(263):486–502.
- Epstein B. The exponential distribution and its role in Life –testing. *Ind Qual Control*. 1958;15:2–7.
- Feil P, Zelen M. Estimation of exponential survival probabilities with concomitant information. *Biometrics*. 1965;21(4):826–838.
- Ghitany ME, Atieh B, Nadarajah S. Lidley distribution and its Applications. *Mathematics Computing and Simulation*. 2008;78(4): 493–506.
- Zakerzadah H, Dolati A. Generalized Lindley distribution. *Journal of Mathematics Extension*. 2010;3(2):13–25.
- Mazucheli J, Achcar JA. The Lindley distribution applied to competing risks lifetime data. *Comput Methods Programs Biomed*. 2011;104(2):188–192.
- Bakouch SH, Al-Zahrani BM, Al-Shomrani AA, et al . An extended Lindley distribution. *Journal of Korean Statistical Society*. 2011; 41(1):75–85.
- Shanker R, Mishra A. A quasi Lindley distribution. *African journal of Mathematics and Computer Science Research*. 2013;6(4):64–71.
- Shanker R, Mishra A. A two-parameter Lindley distribution. *Statistics in transition new series*. 2013;14(1):45–56.
- Shanker R, Sharma S, Shanker R. A two-parameter Lindley distribution for modeling waiting and survival times data. *Applied Mathematics*. 2013;4(2):363–368.
- Shanker R, Amanuel AG. A new quasi Lindley distribution. *International Journal of Statistics and Systems*. 2013;8(2):143–156.
- Deniz EG, Ojeda EC (2011) The discrete Lindley distribution-Properties and Applications. *Journal of Statistical Computation and Simulation* 81(11): 1405-1416.
- Sankaran M. The discrete Poisson-Lindley distribution. *Biometrics*. 1970;26(1):145–149.
- Smith RL, Naylor JC. A comparison of Maximum likelihood and Bayesian estimators for the three parameter Weibull distribution. *Journal of the Royal Statistical Society*. 1987;36(3):358–369.
- Birnbaum ZW, Saunders SC. Estimation for a family of life distributions with applications to fatigue. *Journal of Applied Probability*. 1969; 6(2):328–347.
- Lawless JF. *Statistical models and methods for lifetime data*. John Wiley and Sons, New York, USA; 1982.
- Piccioletto R. Tensile fatigue characteristics of a sized polyester/viscose yarn and their effect on weaving performance, Master thesis, University of Raleigh, North Carolina State, USA; 1970.
- Bjerkedal T. Acquisition of resistance in guinea pigs infected with different doses of virulent tubercle bacilli. *Am J Hyg*. 1960;72:130–148.
- Balakrishnan N, Victor L, Antonio S. A mixture model based on Birnbaum-Saunders Distributions, A study conducted by Authors regarding the Scores of the GRASP (General Rating of Affective Symptoms for Preschoolers), in a city located at South Part of the Chile. 2010.
- Efron B. Logistic regression, survival analysis and the Kaplan-Meier curve. *Journal of the American Statistical Association*. 1988;83(402):414–425.
- Lee ET, Wang JW. *Statistical methods for survival data analysis*. (3rd edn), John Wiley and Sons, New York, USA; 2003.
- Linhart H, Zucchini W. *Model Selection*, John Wiley, New York, USA; 1986.
- Bhaumik DK, Kapur K, Gibbons RD. Testing Parameters of a Gamma Distribution for Small Samples. *Technometrics*. 2009;51(3):326–334.
- Lindley DV. Fiducial distributions and Bayes’ Theorem. *Journal of the Royal Statistical Society*. 1958;20(1):102–107.
- Proschan F. Theoretical explanation of observed decreasing failure rate. *Technometrics*. 1963;5(3):375–383.
- Gross AJ, Clark VA. *Survival Distributions: Reliability Applications in the Biometrical Sciences*, John Wiley, New York, USA; 1975.
- Fuller EJ, Frieman S, Quinn J, et al. Fracture mechanics approach to the design of glass aircraft windows: A case study. *SPIE Proceedings*. 1994:419–430.