

Assessing the role of age specific fertility rate on downturn in maternal mortality ratio in bangladesh

Abstract

Specific mechanism needs to be developed for proper monitoring of maternal mortality with a view to advocating for policy making and tracking progresses. Prior to meeting this need, impact of factors associated with maternal mortality ratio (MMR) needs to be evaluated. Hence, this article endeavors to decompose the overall decline in the MMR in Bangladesh observed between 1991 and 2011 into two components which are namely decline attributable to fertility decline and decline attributable to safe motherhood programs. In this study, age specific fertility rate (ASFR) has been employed as fertility indicator. The results disclose that fertility decline in Bangladesh between 1991 and 2011 has played vital role in reducing MMR. About 44 percent of the estimated number of maternal lives saved in 2011 was attributable to safe motherhood programs. The estimates also show that the MMR in Bangladesh was declined by 53.81% in the last 20 years (1991-2011) on account of fertility reduction merely during the given period. About 41 percent decline was observed in actual birth from projected births owing to fertility decline. Fertility rate decline caused about 49 percent decline in actual maternal deaths (AMD) observed between 1991 and 2011 whereas the remaining portion of decline in AMD was attributed to safe motherhood programs.

Keywords: maternal mortality ratio, fertility; age specific fertility rate, maternal deaths

Abbreviations: MMR, maternal mortality ratio; ASFR, age specific fertility rate; AMD, actual maternal deaths; MDG 5, millennium development goal 5; CBR, crude birth rate

Introduction

The death of a woman during or after (within 42 days) pregnancy or childbirth caused by severe bleeding, infections, high blood pressure during pregnancy (pre-eclampsia and eclampsia), complications from delivery or unsafe abortion is termed as maternal death. It poses a great challenge to global health systems. Albeit several measures have been taken to control but these are not supposed to be sufficient to intensify policy intervention for maternal mortality. An International Conference held in 1994 on Population and Development bolstered international commitment to reproductive health.¹ Facts regarding rates and trends in maternal mortality are imperative for achieving Millennium Development Goal 5 (MDG 5) since its target is 75% reduction in maternal mortality ratio (MMR) from 1990 to 2015.²

MMR refers to the number of maternal deaths per 100,000 live births.³ It is the ratio of maternal deaths and live births during a year reflecting the probability of death resulting from causes associated with pregnancy or childbirth among those who became pregnant. The global average MMR is 226 per 100,000 live births⁴ which is still lower than the proposed annual decline of 5.5% required to achieve the Millennium Development Goal 5.⁵ In accordance with regional-level analysis of trends in MMR decline, progress in sub-Saharan Africa has been relatively slow in comparison with that in South America and Asia.^{5,6} Three South Asian countries-India, Pakistan and Bangladesh accounted for 64% of the global decline of 188,000 annual maternal deaths between 1990 and 2008.³ Unfortunately, in Bangladesh, almost 71% of total births are delivered at home while twenty-nine percent of births are delivered at a health facility including 15 percent in a private facility, 12 percent in a public facility, and 2 percent in an NGO facility.⁷ Therefore a large proportion of births in Bangladesh cannot recognize these obstetric complications and thus may not

refer mother in time to health facilities that provide comprehensive emergency obstetric-care services. MMR can decline when fewer women die because of causes related to pregnancy.

Safe motherhood programs comprising several initiatives were introduced in 1987 with a view to declining maternal mortality ratio.⁸ The main objective of these programs is to make childbirth safer. As part of that these programs paid heed to decline MMR by ensuring healthcare facilities at the time of delivery or involving adroit birth attendants if deliveries take place at home or laying on improved obstetric care facilities.

Albeit all these initiatives have performed reasonably well, nevertheless the progress in maternal mortality has not been found or it has been slow either in many places.^{9,10} The proxy indicators of the effects of safe motherhood programs indicate progress over time; nevertheless it is not able to explain decline in MMR altogether. Hence, in this study we have deemed another component of MMR, i.e. fertility decline along with safe motherhood programs. It is playing role efficaciously in declining MMR. It also helps reduce the number of annual maternal deaths, expected lifetime risk of women's death resulting from pregnancy, childbirth associated causes. A study conducted by Jain³ revealed that around 39 percent decline in MMR between 1990 and 2008 is attributable to fertility reduction while about 61 percent attributable to safe motherhoods program. In that study, crude birth rate (CBR) was used as an indicator of fertility and latter it was used to calculate annual number of live births.

The objective of this study is to measure the effect of fertility decline mainly on overall decline in MMR between 1991 and 2011. We have employed the method proposed by Jain³ and considered age specific fertility rate (ASFR) as an indicator of fertility to calculate annual number of live births.

Data and Methods

Data used in this study were collected from the Population and

Housing Census 2011. The decomposition method used in this includes four steps:

- i. Calculation of projected and actual births in 2011
- ii. Estimation of the potential number of maternal deaths in 2011
- iii. Decomposition of potential number of maternal lives saved in 2011 and
- iv. Estimation of the level of MMR implied by fertility decline between 1991 and 2011. The details of these steps are given below

Calculation of projected and actual births in 2011

In this study we have used ASFR as a fertility indicator. To estimate the projected births in 2011 assuming constant fertility since 1991, we have proceeded to the following approach. Firstly, we have calculated women population in the i^{th} reproductive age group at 2011 with constant annual growth since 1991 $(_{1991}^i W_{\text{projected}})$.

$$_{2011}^i W_{\text{projected}} = _{1991}^i W \times e^{\left\{ _{1991}^i r \times (2011-1991) \right\}}, \quad i = 1, 2, \dots, 7$$

where, $_{1991}^i W$ = women population in the i^{th} reproductive age group at 1991

$_{1991}^i r$ = Average annual growth rate for female at i^{th} age group, 1991-2001

Secondly, using ASFR of the women in the i^{th} reproductive age at 1991 $(_{1991}^i B_{\text{projected}})$ we have calculated projected births for the women belong to that age group at 2011 assuming constant fertility since 1991 $(_{2011}^i B_{\text{projected}})$.

$$_{2011}^i B_{\text{projected}} = \frac{\left(_{1991}^i ASFR \times _{2011}^i W_{\text{projected}} \right)}{1000}$$

Total Number of projected births, $B_{\text{projected}}^{2011} = \sum_{i=1}^7 _{2011}^i B_{\text{projected}}$

We also calculated the actual number of births for the women belong to i^{th} reproductive age group at 2011 $(_{2011}^i B_{\text{Actual}})$ using ASFR of the women in the i^{th} reproductive age at 2011 $(_{2011}^i ASFR)$.

$$_{2011}^i B_{\text{Actual}} = \left(\frac{_{2011}^i ASFR \times _{2011}^i W_{\text{Actual}}}{1000} \right)$$

where, $_{2011}^i W_{\text{Actual}}$ = women population in the i^{th} reproductive age group at 2011.

Total Number of Actual births at 2011,

$$B_{\text{Actual}}^{2011} = \sum_{i=1}^7 _{2011}^i B_{\text{Actual}}$$

Estimation of the potential number of maternal deaths in 2011

To estimate the number of maternal deaths that were averted in 2011, the actual number of maternal deaths occurred in 2011 is first

estimated by using the levels of fertility and the MMR observed in 2011, factoring in the decline observed between 1991 and 2011 in both MMR and fertility. The age specific fertility is used as an indicator of fertility. The annual number of live births given by mothers of a certain age group within reproductive period calculated by multiplying the ASFR of that age group by the total number of female at the age group. Four counterfactual exercises are then conducted to estimate the potential number of maternal deaths expected in 2011 (D_{2011}) under four situations regarding changes in fertility and the MMR (M) between 1991 and 2011:

Situation 1: Neither fertility nor MMR declined

$$Death_{2011}^2 = Birth_{\text{Actual}}^{2011} \times M_{1991}.$$

Situation 2: Fertility declined but MMR did not decline

$$Death_{2011}^2 = Birth_{\text{Actual}}^{2011} \times M_{1991}.$$

Situation 3: Fertility did not decline but MMR declined

$$Death_{2011}^3 = Birth_{\text{projected}}^{2011} \times M_{2011}.$$

Situation 4: Both fertility and MMR declined

$$Death_{2011}^4 = Birth_{\text{Actual}}^{2011} \times M_{2011}.$$

Decomposition of potential number of maternal lives saved in 2011

Using potential number of maternal deaths in 2011 estimated under different situations regarding changes of fertility and MMR we have decomposed the potential number of maternal lives saved in 2011.

The Gross effect of MMR decline on maternal lives saved,

$$B = D_{2011}^1 - D_{2011}^2$$

The Gross effect of Fertility decline on maternal lives saved,

$$B = D_{2011}^1 - D_{2011}^2$$

The Gross effect of decline in both Fertility and MMR on maternal lives saved, $C = D_{2011}^1 - D_{2011}^4$

The Joint effect of decline in both Fertility and MMR on maternal lives saved, $A \cap B = A + B - C$

This joint effect calculation allows us to estimate the net effect of fertility decline and the net effect of MMR decline on the potential number of maternal lives saved in 2011. The net effect of fertility decline would reflect the proportion of potential number of maternal lives saved through such safe motherhood initiatives as institutional deliveries, skilled birth attendants, and comprehensive emergency obstetric-care services. It can be estimated as follows,

The Net effect of decline in MMR, $D = B - (A \cap B)$

The net effect of fertility decline would reflect the proportion of potential number of maternal lives saved through a reduction in the annual number of live births.

The Net effect of decline in Fertility, $D = B - (A \cap B)$

Now the percent of the potential number of maternal lives saved in 2011 due to safe motherhood programs, decrease in live births and fertility reduction realized through its effect on MMR reduction can be estimated as

$$\text{Effect of safe motherhood} = \left[\frac{D}{(C+D)} \right] \times 100$$

$$\text{Effect of decrease in live births} = \left[\frac{D}{(C+D)} \right] \times 100$$

$$\text{Effect of fertility reduction realized through its effect on MMR reduction} = \left[\frac{(A \cap B)}{(C+D)} \right] \times 100$$

Estimation of the level of MMR implied by fertility decline between 1991 and 2011

The difference between the 2011 level of MMR attributable to fertility decline alone and the observed level of MMR in 2011 reflects the effect of changes in factors that make motherhood safer. The value is estimated in following way:

MMR in 2011 implied by fertility reduction observed during 1991-2011,

$$\hat{M}_{2011} = M_{2011} + \left\{ \left(\frac{B_{2011}}{\hat{B}_{2011}} \right) \times (M_{1991} - M_{2011}) \right\}$$

Maternal Deaths in 2011 implied by fertility reduction observed during 1991-2011,

$$\hat{D}_{2011} = \left(\frac{B_{2011} \times \hat{M}_{2011}}{1000} \right)$$

Decomposition of MMR decline observed between 1991 and 2011

The total maternal mortality ratio decline observed between

Table 1 Birth calculation

Reproductive age group of women	Population at 1991 (in '000)	Average annual growth rate for female at reproductive age, 1991-2001	Estimated population at 2011 with constant annual growth since 1991	Population at 2011	ASFR in 1991	ASFR in 2011	Projected births in 2011 assuming constant fertility	Actual births in 2011
15-19	4681	0.0204744	7049810.92	6352398	80	50	563984.87	317619.9
20-24	5009	0.0216127	7717507.56	7522419	200	140	1543501.51	1053138.66
25-29	4934	0.0199549	7354039.15	7254256	180	110	1323727.04	797968.16
30-34	3301	0.026433	5600668.6	5420659	120	70	672080.23	379446.13
35-39	2782	0.0273169	4804286.26	4859079	80	30	384342.9	145772.37
40-44	2215	0.0240648	3584249.24	3980739	40	10	143369.9	39807.39
45-49	1669	0.0209429	2537256	3016800	20	10	50745.1	30168
Total	24591		45902074	38406350			4681751.6	2763921

1991 and 2011 is decomposed into two components: MMR decline attributable to fertility decline and MMR decline attributable to safe motherhood initiatives.

$$\text{Decline in MMR attributable to fertility reduction} = M_{1991} - \hat{M}_{2011}$$

$$\text{Decline in MMR attributable to safe motherhood initiatives} =$$

$$\hat{M}_{2011} - M_{2011}$$

Similarly, the annual number of actual maternal lives saved between 1991 and 2011 is decomposed into these two components.

$$\text{Decline in actual maternal deaths attributable to fertility reduction} = D_{1991} - \hat{D}_{2011}$$

$$\text{Decline in actual maternal deaths attributable to safe motherhood initiatives} = \hat{D}_{2011} - D_{2011}$$

Results and Discussions

Table 2 shows the potential number of maternal lives saved in 2011 because of fertility and MMR reductions that occurred between 1991 and 2011. The age specific fertility rate (ASFR) is used as an indicator of fertility. Apparently 22097 maternal deaths would have occurred in 2011 in Bangladesh if neither fertility nor the MMR declined during this period. In comparison, about 6025 maternal deaths actually occurred with the declines in fertility and the MMR observed during this period. Reductions in both fertility and the MMR observed during this period thus potentially saved close to 16072 maternal lives in Bangladesh. About 44 percent of the estimated number of maternal lives saved in 2011 is attributable to safe motherhood programs. The remaining 56 percent of the estimated number of maternal lives saved in 2011 can be attributed to fertility reduction: 26 percent attributable to the decrease in the number of live births and 30 percent attributable to changes in the age–parity composition of births.

Table 2 Effect of declines in MMR and fertility on estimated number of maternal deaths

Parameter	Estimated figure
Estimated number of maternal deaths in 2011	
No change in ASFR and no change in MMR	22097.87
No change in ASFR and MMR declined	10206.22
ASFR declined but no change in MMR	13045.7
Both ASFR and MMR declined	6025.35
Potential number of maternal lives saved in 2011 from	
Total effect of decline in MMR	11891.65
Total effect of decline in Fertility decline	9052.16
Total effect of decline in both	16072.52
Overlap between the effect of declines in fertility and in MMR	4871.29
Net effect of decline in MMR	7020.35
Net effect of decline in Fertility	4180.87
Percent distribution of the potential number of maternal lives saved in 2011 (%)	
Effect of safe motherhood	43.68
Effect of safe decrease in live births	26.01
Effect of fertility reduction realized through its effect on MMR reduction	30.3

Decomposition of MMR decline observed between 1991 and 2011

Table 3 shows the estimated decline in the MMR in 2011 attributable to fertility decline observed between 1991 and 2011. The estimates show that the MMR in Bangladesh would have declined

from 472 in 1991 to 218 in 2011 because of fertility reduction alone during this period. This means that 104 points, or about 41 percent of the total decline of 254 points in the MMR in Bangladesh, can be attributed to fertility reduction. The remaining decline of about 59 percent can thus be attributed to safe motherhood programs.

Table 3 MMR in 2011 implied by fertility decline observed between 1991 and 2011

Parameter	Estimated figure
Proportionate decline in actual births from projected births in 2011	0.41
Total decline in MMR between 1991 and 2011	254
MMR in 2011 implied by fertility reduction observed during 1991-2011	367.95
Decline in MMR attributable to fertility reduction	104.04
Decline in MMR attributable to safe motherhood initiatives	149.95
% decline in MMR attributable to fertility reduction	40.96
% decline in MMR attributable to safe motherhood initiatives	59.04

Decomposition of maternal deaths decline observed between 1991 and 2011

Table 4 shows the estimated decline in the maternal deaths in 2011 attributable to fertility decline observed between 1991 and 2011. Number of actual maternal deaths in 1991 was around 1418388 and

it was dropped more than 50 percent in the year 2011. In Bangladesh about 815853 fewer women died in 2011 from childbirth-related complications than in 1991. About 401399 (49 percent) of these actual maternal lives saved in Bangladesh can be attributed to fertility decline observed between 1991 and 2011.

Table 4 Actual maternal deaths in 2011 implied by fertility decline observed between 1991 and 2011

Parameter	Estimated figure
Actual Births in 1991	3005060
Actual Births in 2011	2763920.61
MMR in 1991	472
MMR in 2011	218
Total decline in MMR between 1991 and 2011	254
MMR in 2011 implied by fertility reduction observed during 1991-2011	367.95
Maternal Deaths in 1991	1418388.32
Maternal Deaths in 2011	602534.69
Maternal Deaths in 2011 implied by fertility reduction observed during 1991-2011	1016988.82
Decline in actual maternal deaths between 1991 to 2011	
Total	815853.6
Attributable to Fertility Decline	401399.5
Attributable to Safe Motherhood	414454.13
Percent decline in actual maternal deaths between 1991 to 2011	
Attributable to Fertility Decline	49.2
Attributable to Safe Motherhood	50.8

Conclusions

Maternal mortality ratio (MMR) is defined as the number of maternal deaths per 100,000 live births. There are various causes of maternal deaths, mainly by four causes namely hemorrhage, sepsis, hypertensive disorders, and obstructed labor. Usually, three indicators are used to measure maternal mortality. These are the maternal mortality rate, women's average lifetime risk of death due to pregnancy-related causes, and the annual number of maternal fatalities which can be easily derived mathematically. The calculation of these indicators depend on the proportion of women becoming pregnant in a year, the average number of pregnancies per woman, and the product of population size and crude birth rate. The relationship between fertility decline and MMR is not so easily derived. Since millennium development goal 5 uses MMR as its indicator, and prevailing attributions of declines in maternal mortality entirely to the achievements of safe motherhood, therefore studying such relationship

is crucial.³ In this article we have used a simple method proposed in³ to decompose the overall decline in the MMR in Bangladesh observed between 1991 and 2011 into two components: decline attributable to fertility decline and decline attributable to safe motherhood programs. They have used crude birth rate as fertility indicator. However, this study used age specific fertility rate as fertility indicator. The results show that fertility decline in Bangladesh between 1991 and 2011 has made a substantial contribution to the reduction of the MMR. About 44 percent of the estimated number of maternal lives saved in 2011 is attributable to safe motherhood programs. The estimates show that the MMR in Bangladesh would have declined from 472 in 1991 to 218 in 2011 because of fertility reduction alone during this period. About 41 percent of the total decline of 254 points in the MMR in Bangladesh can be attributed to fertility reduction. About 401399 (or about 49 percent) of actual maternal lives saved can be attributed to fertility decline observed between 1991 and 2011.

Acknowledgement

None.

Conflict of Interest

The authors declare that they have no financial or non-financial competing interests.

References

1. Obaid TA. Fifteen years after the International Conference on Population and Development: what have we achieved and how do we move forward? *Int J Gynaecol Obstet.* 2009;106(2):102–105.
2. United Nations General Assembly. *United Nations Millennium Declaration.* A/RES/55/2. New York: United Nations; 2000.
3. Jain AK. Measuring the effect of fertility decline on the maternal mortality ratio. *Stud Fam Plann.* 2011;2(4):247–260.
4. Goli S, Jaleel ACP. What is the cause of the decline in maternal mortality in India? Evidence from time series and cross sectional analyses. *J Biosoc Sci.* 2014;46(3):351–365.
5. WHO, UNICEF, UNFPA & World Bank. *The World Trends in Maternal Mortality: 1990 to 2010.* World Health Organization, Geneva; 2012.
6. Hogan MC, Foreman KJ, Naghavi M, et al. Maternal mortality for 181 countries, 1980-2008: a systematic analysis of progress towards Millennium Development Goal 5. *Lancet.* 2010;375(9726):1609–1623.
7. BDHS. *Bangladesh Demographic and Health Survey 2011. National Institute of Population Research and Training (NIPORT), Mitra and Associates, and Macro International.* USA: Dhaka, Bangladesh and Calverton, Maryland; 2013.
8. Starrs AM. Safe motherhood initiative: 20 years and counting. *Lancet.* 2006;368(9542):1130–1132.
9. Hill K, Thomas K, AbouZahr C, et al. Estimates of maternal mortality worldwide between 1990 and 2005: an assessment of available data. *Lancet.* 2007;370(9595):1311–1319.
10. United Nations. *Goal 5 Maternal Health Fact Sheet. United Nations Summit High-Level Plenary Meeting of the General Assembly.* New York; 2010.