Bound lengths for type-I progressive hybrid burr type-XII censored data under SS-PALT

Abstract

Our main focus on combining two different approaches, Step-Stress Partially Accelerated Life Test and Type-I Progressive Hybrid censoring criteria in the present article. The fruitfulness of this combination has been investigated by bound lengths for unknown parameters of the Burr Type-XII distribution. Approximate confidence intervals, Bootstrap confidence intervals and One-Sample Bayes prediction bound lengths have been obtained under the above scenario. Particular cases of Type-I Progressive Hybrid censoring (Type-I and Progressive Type-II censoring) has also evaluated under SS-PALT. Optimal stress change time also measured by minimizing the asymptotic variance of ML Estimation. A simulation study based on Metropolis-Hastings algorithm have carried out along with a real data set example.

Keywords: Progressive Type-II, probability, Burr Type-XII distribution, chemical engineering, quality control

Volume 10 Issue 1 - 2021

Gyan Prakash

Department of Community Medicine, Moti Lal Nehru Medical College, India

Correspondence: Gyan Prakash, Department of Community Medicine, Moti Lal Nehru Medical College, Allahabad, U. P., India, Email ggyanji@yahoo.com

Received: June 07, 2020 | **Published:** February 23, 2021

Abbreviations: SS-PALT, step-stress partially accelerated life test; T-IPH, Type-I progressive hybrid; BPBL, bayes prediction bound lengths; BCL, bootstrap confidence lengths; ACL, approximate confidence lengths; T-I, Type-I censoring; PT-II, Progressive Type-II

Introduction

A family of distributions, which includes twelve different cumulative distribution functions, was suggested by Burr.¹ The suggested distribution was named as Burr distribution, having a variety of density shapes. Burr Type-XII distribution have been considered in the present discussion due to its applicability on several applied fields. For example, business, chemical engineering, quality control, duration analysis, failure time modeling and reliability studies etc. The probability density function and cumulative density function along with Hazard rate function and Reliability function of Burr Type-XII distribution are given as

$$f(y; \alpha, \beta) = \beta \alpha y^{\beta-1} (1 + y^\beta)^{-\alpha-1}; \alpha > 0, \beta > 0, y \geq 0, \quad (1)$$

$$F(y; \alpha, \beta) = 1 - (1 + y^\beta)^{-\alpha}; \alpha > 0, \beta > 0, y \geq 0, \quad (2)$$

$$\rho(y) = \beta \alpha \frac{y^{\beta-1}}{1 + y^\beta}; \alpha > 0, \beta > 0, y \geq 0 \quad (3)$$

and

$$R(y) = (1 + y^\beta)^{-\alpha}; \alpha > 0, \beta > 0, y \geq 0. \quad (4)$$

Parameter α and β are denoted by the shape parameters of the Burr Type-XII distribution. The underlying distribution is selected for the present discussion because it covers a wide range of values of skewness and kurtosis and is fit for any general lifetime data model.

A massive amount of references available for concerned distribution, only few important and recent references is included here. Abd-Elfattah et al.² presents some estimation problems under step-stress partially accelerated life tests for the Burr Type-XII distribution on Type-I censoring pattern. Khalaf et al.³ discussed about the finite mixture of Burr Type-XII distribution with its reciprocal and proposed

failure rate for the new model, covers several types of failure rates. Soliman et al.⁴ deals Bayesian inference and prediction problems by using a Gibbs sampling procedure in Burr Type-XII distribution on progressive first failure censoring criteria.

Jang et al.⁵ discussed properties of the estimation on a Bayesian setup by using progressive censoring on underlying distribution. A multi-component stress strength reliability model was proposed by Rao et al.⁶ for Burr Type-XII distribution. Panahi & Sayyareh⁷ presents the statistical inference and prediction on unified hybrid-censored Burr Type-XII data by using Lindley's approximation.

Prakash⁸ discussed different confidence limits on constant stress partially accelerated life test for Burr Type-XII distribution under progressive Type-II censoring. Asl et al.⁹ studied the properties of maximum likelihood and Bayes estimators on progressive Type-II hybrid censored samples for underlying distribution by using expectation -maximization algorithm. Recently, Prakash⁹ studied the properties of Bayes estimators by using constant stress partially accelerated life test, assuming that the lifetimes of test items follow Burr Type-XII distribution on progressive Type-II censored data.

The usefulness of censoring and/or accelerated life test criteria is, minimizes the test expenditures and length of testing. In the present study, we considered both, the Type-I progressive hybrid censoring and step-stress partially accelerated life test. No such study has found for the combination of these two along with Bayes prediction in the literature. So, the main objective of the study is to combine these two for investigating the fruitfulness in terms of bounds lengths for the unknown parameters. One-Sample Bayes prediction bound lengths along with Approximate and Bootstrap confidence lengths are obtained for the study.

T-IPH censoring with their special cases viz., Type-I (T-I) and Progressive Type-II (PT-II) has also been investigated in terms of bound lengths under SS-PALT in the present discussion. Asymptotic variance of ML estimation was minimized for determining an optimal stress change time for refining the quality of the inference. Metropolis-Hastings (M-H) algorithm has been used in the simulation study along with real data examples.

Type-I Progressive hybrid (T-IPH) censoring

Following Prakash,¹¹ let us suppose total n live test units are placed on a life test with $T_1, T_2, \dots, T_n$ corresponding lifetimes respectively. Units under test are identical and independently distributed Eq. (1). It is noted further that, T-IPH censoring is the mixture of Type-I censoring and Progressive Type-II censoring pattern. Hence, in T-IPH censoring, the test stops either at pre-considered number of failure m ($\leq n$) or at pre-considered time t , which one occurred earlier, and both are pre-fixed at the time of life test start.

Now, there arises two circumstances, first, if the trial stops at m^{th} failure then the observed samples are $X_1, X_2, \dots, X_\varepsilon, X_{\varepsilon+1}, \dots, X_m$; $X_m < t$ with assumed censoring pattern $R \equiv (R_1, R_2, \dots, R_m)$ and must be satisfied $n - m = R_1 + R_2 + \dots + R_m$, and is called progressive Type-II censoring. In the second state, if the experiment stops at time t with a number of failures X_j , then they must satisfy the condition $X_j < t < X_{j+1}$ and remaining all the live test units are removed from the test. So, the observed samples are $X_1, X_2, \dots, X_\varepsilon, X_{\varepsilon+1}, \dots, X_j$, if $X_j < t < X_{j+1}$ with $n - R_1 - R_2 - \dots - R_j - j = R^*$ (say). Hence, in T-IPH censoring pattern the observed samples have drawn from following two ways

$$\begin{cases} I: (X_1, X_2, \dots, X_\varepsilon, X_{\varepsilon+1}, \dots, X_m), \text{ if } X_m < t \\ II: (X_1, X_2, \dots, X_\varepsilon, X_{\varepsilon+1}, \dots, X_j), \text{ if } X_j < t < X_{j+1} \end{cases}$$

Numerous literatures on progressive hybrid censoring has available, a little few important studies are discussed here, however, no any such study has found on above considered scenario. Some classical estimation of unknown parameters was discussed by Lin et al.¹² for two-parameter Weibull distribution on Type-II progressive hybrid censored pattern. Lin & Huang¹³ discussed adaptive Type-I progressive hybrid censoring scheme and derived the exact distribution of the maximum likelihood estimator of the mean lifetime of an Exponential distribution. Singh et al.¹⁴ discussed about the properties of Bayes estimation for unknown parameters of the Lindley distribution under Type-II hybrid censored data.

For more informative literature on hybrid censoring one may go with Balakrishnan & Cramer.¹⁵ Elshahhat & Ashour¹⁶ discussed about the Bayesian and non-Bayesian estimation under the generalized Type-II progressive hybrid censoring scheme for Weibull parameters. Some inferences on Burr Type-XII distribution have recently discussed by Kayal et al.¹⁷ under Type-I progressive hybrid censoring.

Step - stress partially accelerated life test

Due to the reliability of the products, there occurred few failures at the end of several life tests. In such lifetime test, the test on normal stress condition more expensive and time intense. In such cases, accelerated life test is used for getting information about the lifetime distribution of item shortly and less expensive by test units kept at higher than usual stress conditions. In step stress accelerated life test, usually the test starts at a low stress (normal) condition, if the units do not fail at a pre-specified time, stress on its elevated and held at specified times. Stress is recurrently increased until the test unit fails or censoring time is reached.¹⁸ A great amount of literature available on SS-PALT, see Prakash¹⁹ for recent studies on SS-PALT.

All live units are verified first at normal stress condition in SS-PALT, if test units do not fail for a pre-assumed time $\hat{a}$ (say), then the

test is swapped at higher stress and continued until test units fail. A reformed random variable model was discussed by DeGroot & Goel²⁰ for the management of such issue, and defined here by the help of stress change time parameter ε and acceleration factor parameter λ as

$$X = \begin{cases} Y & ; 0 < Y \leq \varepsilon \\ \varepsilon + \frac{Y - \varepsilon}{\lambda} & ; Y > \varepsilon \end{cases} \quad (5)$$

Here, the normal stress and higher stress conditions are denoted by “I” and “II” respectively for SS-PALT. If X is assumed as the total lifetime of a test item, then Eq. (5) is rewritten as

$$f(x; \alpha, \beta) = \begin{cases} I: f_1 = \beta \alpha x^{\beta-1} (1+x^\beta)^{-\alpha-1} & ; 0 < x \leq \varepsilon \\ II: f_2 = \beta \alpha \lambda \tilde{x}^{\beta-1} (1+\tilde{x}^\beta)^{-\alpha-1} & ; x > \varepsilon, \tilde{x} = (x - \varepsilon)\lambda + \varepsilon \end{cases} \quad (6)$$

Under T-IPH censoring, the joint probability density (likelihood) function is now written by using Eq. (6) as

$$\begin{cases} I:L \propto \prod_{i=1}^k (f_1(1-F_1)^{R_i}) \times \prod_{i=k+1}^m (f_2(1-F_2)^{R_i}) \\ II:L \propto \prod_{i=1}^l (f_1(1-F_1)^{R_i} (1-F_1(t))^{R^*}) \times \prod_{i=l+1}^j (f_2(1-F_2)^{R_i} (1-F_2(t))^{R^*}) \end{cases} \quad (7)$$

Solving Eq. (7), we get

$$L \propto \beta^d \alpha^d \lambda^{d-\delta} \omega_0 \omega_1 e^{-\alpha \omega_d} ; \quad (8)$$

$$\text{where } \omega_0 = \prod_{i=1}^{\delta} \left(\frac{\xi_{(i)} x_{(i)}}{1 + R_i} \right), \quad W_1 = \sum_{i=1}^k (1 + R_i) \log(1 + x_{(i)}^\beta),$$

$$d = \begin{cases} I: m \\ II: j \end{cases}, \quad \delta = \begin{cases} I: k \\ II: l \end{cases},$$

$$\omega_1 = \prod_{i=1+\delta}^d \left(\frac{\tilde{\xi}_{(i)} \tilde{x}_{(i)}}{1 + R_i} \right), \quad \omega_d = \begin{cases} I: W_1 + W_2 \\ II: W_3 + W_4 \end{cases},$$

$$W_3 = \sum_{i=1}^l (1 + R_i) \log(1 + x_{(i)}^\beta) + R^* \log(1 + t^\beta),$$

$$W_2 = \sum_{i=1+k}^m (1 + R_i) \log(1 + \tilde{x}_{(i)}^\beta),$$

$$W_4 = \sum_{i=l+1}^j (1 + R_i) \log(1 + \tilde{x}_{(i)}^\beta) + R^* \log(1 + \tilde{t}^\beta), \quad \xi_{(i)} = x_{(i)}^{\beta-1}$$

$$\left(\frac{1 + R_i}{1 + x_{(i)}^\beta} \right), \quad \tilde{\xi}_{(i)} = \left(\frac{1 + R_i}{1 + \tilde{x}_{(i)}^\beta} \right) \tilde{x}_{(i)}^{\beta-1} \text{ and } \tilde{t} = (t - \varepsilon)\lambda + \varepsilon.$$

Optimization criterion in SS-PALT

The optimization criterion provided the optimum time duration for the lower stress level. This design is important for improving the precision in estimating the life distribution at design stress and for refining the quality of the inference. The proposed optimization criterion is based on the maximization of the determinant of Fisher's

information matrix which is equivalent to reducing the generalized asymptotic variance of the maximum likelihood estimate of the model parameters. The generalized asymptotic variance is the reciprocal of the determinant of Fisher's information matrix. The Fisher's information matrix is given here as

$$F = - \begin{pmatrix} \frac{\partial^2}{\partial \alpha^2} \text{Log } L & \frac{\partial^2}{\partial \alpha \partial \beta} \text{Log } L & \frac{\partial^2}{\partial \alpha \partial \lambda} \text{Log } L \\ \frac{\partial^2}{\partial \beta \partial \alpha} \text{Log } L & \frac{\partial^2}{\partial \beta^2} \text{Log } L & \frac{\partial^2}{\partial \beta \partial \lambda} \text{Log } L \\ \frac{\partial^2}{\partial \lambda \partial \alpha} \text{Log } L & \frac{\partial^2}{\partial \lambda \partial \beta} \text{Log } L & \frac{\partial^2}{\partial \lambda^2} \text{Log } L \end{pmatrix}.$$

The components of the matrix given in Eq. (9) are obtained as

$$\begin{aligned} \frac{\partial^2}{\partial \alpha^2} \text{Log } L &= -\frac{d}{\alpha^2} \\ \frac{\partial^2}{\partial \beta^2} \text{Log } L &= -\frac{d}{\beta^2} + \sum_{i=1}^{\delta} \left(\frac{(1+x_{(i)}^{\beta} - \beta^2 x_{(i)}^{\beta-1})}{x_{(i)}(1+x_{(i)}^{\beta})^2} \right) + \sum_{i=\delta+1}^d \left(\frac{(1+\tilde{x}_{(i)}^{\beta} - \beta^2 \tilde{x}_{(i)}^{\beta-1})}{\tilde{x}_{(i)}(1+\tilde{x}_{(i)}^{\beta})^2} \right) \\ -\alpha &\left\{ \begin{array}{l} \text{I: } \sum_{i=1}^k \xi_{(i)} + \sum_{i=k+1}^m \tilde{\xi}_{(i)} \\ \text{II: } \sum_{i=1}^l \xi_{(i)} + \sum_{i=l+1}^j \tilde{\xi}_{(i)} + t^* + \tilde{t}^* \end{array} \right. \\ +\alpha\beta &\left\{ \begin{array}{l} \text{I: } \sum_{i=1}^k \xi_{(i)}^2 + \sum_{i=k+1}^m \tilde{\xi}_{(i)}^2 \\ \text{II: } \sum_{i=1}^l \xi_{(i)}^2 + \sum_{i=l+1}^j \tilde{\xi}_{(i)}^2 + t^{2*} + \tilde{t}^{2*} \end{array} \right. \\ \frac{\partial^2}{\partial \lambda^2} \text{Log } L &= -\frac{d-\delta}{\lambda^2} - \prod_{i=\delta+1}^d \left(\beta(x_{(i)} - \varepsilon)^2 \frac{1+\tilde{x}_{(i)}^{\beta} + \beta\tilde{x}_{(i)}^{\beta}}{\tilde{x}_{(i)}^2(1+\tilde{x}_{(i)}^{\beta})^2} \right) \\ +\alpha\beta &\left\{ \begin{array}{l} \text{I: } \sum_{i=k+1}^m \tilde{\xi}_{(i)}^2 (x_{(i)} - \varepsilon)^2 \\ \text{II: } \sum_{i=l+1}^j \tilde{\xi}_{(i)}^2 (x_{(i)} - \varepsilon)^2 - \tilde{t}^{2*} (t - \varepsilon)^2 \end{array} \right. \\ \frac{\partial^2}{\partial \alpha \partial \beta} \text{Log } L &= \frac{\partial^2}{\partial \beta \partial \alpha} \text{Log } L = -\beta \left\{ \begin{array}{l} \text{I: } \sum_{i=1}^k \xi_{(i)} + \sum_{i=k+1}^m \tilde{\xi}_{(i)} \\ \text{II: } \sum_{i=1}^l \xi_{(i)} + \sum_{i=l+1}^j \tilde{\xi}_{(i)} + t^* + \tilde{t}^* \end{array} \right. \\ \frac{\partial^2}{\partial \alpha \partial \lambda} \text{Log } L &= \frac{\partial^2}{\partial \lambda \partial \alpha} \text{Log } L = -\beta \left\{ \begin{array}{l} \text{I: } \sum_{i=k+1}^m (x_{(i)} - \varepsilon) \tilde{\xi}_{(i)} \\ \text{II: } \sum_{i=l+1}^j (x_{(i)} - \varepsilon) \tilde{\xi}_{(i)} + \tilde{t}^* (t - \varepsilon) \end{array} \right. \\ \frac{\partial}{\partial \beta \partial \lambda} \text{Log } L &= \frac{\partial}{\partial \lambda \partial \beta} \text{Log } L = -\beta \sum_{i=\delta+1}^d \left(\frac{1+(1+\beta)\tilde{x}_{(i)}^{\beta}}{\tilde{x}_{(i)}^2(1+\tilde{x}_{(i)}^{\beta})^2} \right) (x_{(i)} - \varepsilon) \\ +\alpha\beta &\left\{ \begin{array}{l} \text{I: } \sum_{i=k+1}^m \tilde{\xi}_{(i)}^2 (x_{(i)} - \varepsilon) \\ \text{II: } \sum_{i=l+1}^j \tilde{\xi}_{(i)}^2 (x_{(i)} - \varepsilon) + \tilde{t}^{2*} (t - \varepsilon) \end{array} \right. \end{aligned}$$

$$\begin{aligned} \text{where } \xi_{(i)}^2 &= \xi_{(i)} \left(\frac{x_{(i)}^{\beta} + 1 - \beta}{(1+x_{(i)}^{\beta})x_{(i)}} \right), \quad \tilde{\xi}_{(i)}^2 = \tilde{\xi}_{(i)} \left(\frac{\tilde{x}_{(i)}^{\beta} + 1 - \beta}{(1+\tilde{x}_{(i)}^{\beta})\tilde{x}_{(i)}} \right), \\ t^* &= \frac{R^*}{1+t^{\beta}} t^{\beta-1}, \quad \tilde{t}^* = \frac{R^*}{1+\tilde{t}^{\beta}} \tilde{t}^{\beta-1}, \quad t^{2*} = t^* \left(\frac{t^{\beta} + 1 - \beta}{t(1+t^{\beta})} \right) \text{ and} \\ \tilde{t}^{2*} &= \tilde{t}^* \left(\frac{\tilde{t}^{\beta} + 1 - \beta}{\tilde{t}(1+\tilde{t}^{\beta})} \right). \end{aligned}$$

The generalized asymptotic variance is the reciprocal of $|F|$. Following Ismail & Aly,²¹ the optimum stress change time ε^* is obtained by maximizing $|F|$ and is obtained by

$$\frac{\partial |F|}{\partial \varepsilon^*} = 0. \quad (10)$$

The solution of Eq. (10) is not in a closed form, numerical solution have been obtained by using Wolfram Mathematica software 10.0.

Approximate confidence lengths

The variance covariance matrix can be approximated by the reciprocal of Fisher's information matrix. All the elements of Fisher's information matrix involved unknown parameters. Hence, an estimate of the variance covariance matrix can be obtained by replacing the parameters by their ML estimate.

Therefore, for obtaining the ML estimators, the first derivatives of the logarithm of the joint probability likelihood function (Eq. (8)) with respect to the corresponding parameters are obtained as

$$\frac{\partial}{\partial \alpha} \text{Log } L = \frac{d}{\alpha} - \omega_d \quad (11)$$

$$\begin{aligned} \frac{\partial}{\partial \beta} \text{Log } L &= \frac{d}{\beta} + \beta \left\{ \sum_{i=1}^{\delta} \frac{\xi_{(i)}}{(1+R_i)x_{(i)}^{\beta}} + \sum_{i=\delta+1}^d \frac{\tilde{\xi}_{(i)}}{(1+R_i)\tilde{x}_{(i)}^{\beta}} \right\} \\ -\alpha\beta &\left\{ \begin{array}{l} \text{I: } \sum_{i=1}^k \xi_{(i)} + \sum_{i=k+1}^m \tilde{\xi}_{(i)} \\ \text{II: } \sum_{i=1}^l \xi_{(i)} + \sum_{i=l+1}^j \tilde{\xi}_{(i)} + t^* + \tilde{t}^*, \end{array} \right. \quad (12) \end{aligned}$$

and

$$\begin{aligned} \frac{\partial}{\partial \lambda} \text{Log } L &= \frac{d-\delta}{\lambda} + \prod_{i=\delta+1}^d \left(\beta(x_{(i)} - \varepsilon) \frac{\tilde{x}_{(i)}^{-1}}{1+\tilde{x}_{(i)}^{\beta}} \right) \\ -\alpha\beta &\left\{ \begin{array}{l} \text{I: } \sum_{i=k+1}^m \tilde{\xi}_{(i)} (x_{(i)} - \varepsilon) \\ \text{II: } \sum_{i=l+1}^j \tilde{\xi}_{(i)} (x_{(i)} - \varepsilon) + \tilde{t}^* (t - \varepsilon) \end{array} \right. \quad (13) \end{aligned}$$

Now, $\hat{\alpha}_{ML}$, $\hat{\beta}_{ML}$ and $\hat{\lambda}_{ML}$ are denoted the ML estimators corresponding to the parameters α, β and λ respectively. The ML estimate corresponding to parameter α is obtained from Eq. (11) as

$$\hat{\alpha}_{ML} = \frac{d}{\omega_d} \quad (14)$$

Consequently, by substituting for $\hat{\alpha}_{MI}$ into Eq. (12) & Eq. (13), the corresponding equations are reduced into two nonlinear equations. The Newton-Raphson method is applied here for simultaneously solving these nonlinear equations for ML estimate $\hat{\beta}_{MI}$ and $\hat{\lambda}_{MI}$, and an iterative procedure is applied to solve these equations numerically by Mathcad statistical package.²²

The applicability of normal approximation of ML Estimation is in small sample size. A log-transformation can be considered for improvements in the performance of the normal approximation. Hence, based on the normal approximation, 100(1 - τ)% approximate confidence lengths are obtained respectively as

$$\alpha_{Acl} = \hat{\alpha}_{MI} \left\{ \exp \left(\frac{Z_{\tau/2} \sqrt{\text{Var}(\hat{\alpha}_{MI})}}{\hat{\alpha}_{MI}} \right) - \exp \left(-\frac{Z_{\tau/2} \sqrt{\text{Var}(\hat{\alpha}_{MI})}}{\hat{\alpha}_{MI}} \right) \right\}, \quad (15)$$

$$\beta_{Acl} = \hat{\beta}_{MI} \left\{ \exp \left(\frac{Z_{\tau/2} \sqrt{\text{Var}(\hat{\beta}_{MI})}}{\hat{\beta}_{MI}} \right) - \exp \left(-\frac{Z_{\tau/2} \sqrt{\text{Var}(\hat{\beta}_{MI})}}{\hat{\beta}_{MI}} \right) \right\} \quad (16)$$

and

$$\lambda_{Acl} = \hat{\lambda}_{MI} \left\{ \exp \left(\frac{Z_{\tau/2} \sqrt{\text{Var}(\hat{\lambda}_{MI})}}{\hat{\lambda}_{MI}} \right) - \exp \left(-\frac{Z_{\tau/2} \sqrt{\text{Var}(\hat{\lambda}_{MI})}}{\hat{\lambda}_{MI}} \right) \right\}. \quad (17)$$

Bootstrap confidence lengths

Another method for obtaining the confidence intervals is considered in this section proposed by Efron & Tibshirani.²³ They have developed a technique that can be routinely applied without any hefty theoretical thought. This method is very useful in such cases when an assumption regarding the normality is invalid. Kreiss & Paparoditis²⁴ stated that, this method is a re-sampling method for estimating biases, risks and confidence intervals in the statistical inference. Following, Kundu et al.,²⁵ the bootstrap confidence limits are obtained for the underlying parameters. The bootstrap samples are obtained as

Step 1: Generate a sample of size d from an underlying model given in Eq. (1) by using inversion method. The estimated distribution has obtained by using generated samples $x_{(1)}, x_{(2)}, \dots, x_{(d)}$.

Step 2: From the estimated distribution obtained in Step 1, generate a bootstrap sample $x_{(1)}^*, x_{(2)}^*, \dots, x_{(d)}^*$. Using T-IPH censoring patterns, the ML estimate $\hat{\alpha}_{MI}$, $\hat{\beta}_{MI}$ and $\hat{\lambda}_{MI}$ say $\hat{\hat{\alpha}}_{MI}$, $\hat{\hat{\beta}}_{MI}$ and $\hat{\hat{\lambda}}_{MI}$ are computed for the estimated distribution.

Step 3: Repeat the above step up $N(=1,000)$ times to obtain N different bootstrap samples. Arrange all these samples ($\hat{\hat{\alpha}}_{MI}$, $\hat{\hat{\beta}}_{MI}$ and $\hat{\hat{\lambda}}_{MI}$) in ascending order to obtain a final sample of the form

$$v_{\alpha}^1 \leq v_{\alpha}^2 \leq \dots \leq v_{\alpha}^N \text{ for } \hat{\hat{\alpha}}_{MI}$$

$$v_{\beta}^1 \leq v_{\beta}^2 \leq \dots \leq v_{\beta}^N \text{ for } \hat{\hat{\beta}}_{MI}$$

and

$$v_{\lambda}^1 \leq v_{\lambda}^2 \leq \dots \leq v_{\lambda}^N \text{ for } \hat{\hat{\lambda}}_{MI}.$$

Step 4: If $H(y) = P(v_{\Theta}^* \leq y)$ be the cumulative density function of v_{Θ}^* ; $\forall \Theta = \alpha, \beta, \lambda$ from final bootstrap samples. Then 100(1 - τ)%

approximate bootstrap confidence limits (BCL) are given for $\Theta(= \alpha, \beta, \lambda)$

$$\left[v_{\Theta}^* \left(\frac{\varepsilon}{2} \right), v_{\Theta}^* \left(\frac{2 - \varepsilon}{2} \right) \right] \quad (18)$$

where $v_{\Theta}^* = H^{-1}(y)$ for given y .

One-sample bayes prediction bound length

In lacks of any authentic advice for selection of prior distribution, it's completely depends on individual interest, belief and objective in Bayesian methodology. Our major objective is to study the fruitfulness of the combination of T-IPH censoring and SS-PALT in terms of bound lengths under Bayesian setup. Prakash⁸ assumed the vague prior for all unknown parameters, so that the prior does not play any significant role in the analyses that follow. Hence, the joint prior probability density function is given as

$$\pi_{(\alpha, \beta, \lambda)} \propto \frac{1}{\alpha \beta \lambda}, \alpha > 0, \beta > 0, \lambda > 0. \quad (19)$$

Based on Bayesian theorem, the joint and marginal posterior densities corresponding to parameters are obtained respectively as

$$\begin{aligned} \pi_{(\alpha, \beta, \lambda)}^* &= \frac{\pi_{(\alpha, \beta, \lambda)} \times L}{\int \int \int \pi_{(\alpha, \beta, \lambda)} \times L \, d\alpha \, d\lambda \, d\beta} \\ &\propto \frac{\beta^{d-1} \alpha^{d-1} \lambda^{d-\delta-1} \omega_0 \omega_1 e^{-\alpha \omega_d}}{\int \beta^{d-1} \omega_0 \int \lambda^{d-\delta-1} \omega_1 \int \alpha^{d-1} e^{-\alpha \omega_d} \, d\alpha \, d\lambda \, d\beta} \\ &\Rightarrow \pi_{(\alpha, \beta, \lambda)}^* \propto \Omega \beta^{d-1} \alpha^{d-1} \lambda^{d-\delta-1} \omega_0 \omega_1 e^{-\alpha \omega_d}; \end{aligned} \quad (20)$$

$$\text{where } \Omega = \left\{ \Gamma(d) \int \beta^{d-1} \omega_0 \int \lambda^{d-\delta-1} \frac{\omega_1}{(\omega_d)^d} \, d\lambda \, d\beta \right\}^{-1}.$$

Corresponding to parameters α , β and λ , marginal posteriors densities are obtained as

$$\pi_{(\alpha)}^* \propto \Omega \alpha^{d-1} \int \beta^{d-1} \omega_0 \int \lambda^{d-\delta-1} e^{-\alpha \omega_d} \, d\lambda \, d\beta, \quad (21)$$

$$\pi_{(\beta)}^* \propto \Omega \Gamma(d) \beta^{d-1} \omega_0 \int \lambda^{d-\delta-1} (\omega_d)^{-d} \, d\lambda \quad (22)$$

and

$$\pi_{(\lambda)}^* \propto \Omega \Gamma(d) \lambda^{d-\delta-1} \int \omega_0 \beta^{d-1} (\omega_d)^{-d} \, d\beta. \quad (23)$$

Prediction of the future observations based on informative sample is an interesting topic in Bayesian analysis and have used in various purposes. Several applications can be found in actuarial studies, rainfall extremes, guarantee data analysis and highest water levels. For example, in guarantee data analysis, setting up warranty period for a product, a manufacturer would use some of the known previous failure times to predict a suitable warranty period of the product. Several applications of Bayes prediction were discussed by Al-Hussaini,²⁶ in different areas of applied statistics.

Now, let us assume $\underline{X}(= x_{(1)}, x_{(2)}, \dots, x_{(d)})$ be the first ordered

observed items from Burr Type-XII distribution under above considered scenario. If Z is assumed as future random variable following underlying model having independent ordered sample $\underline{Z} (= z_{(1)}, z_{(2)}, \dots, z_{(d)})$ under similar scenario. Then the Bayes predictive density function corresponding to a variable Z , is denoted and defined as

$$h(z | \underline{x}) = \int_{\Theta} f(z; \alpha, \beta) \times \pi_{(\Theta)}^* d\Theta; \Theta = \alpha, \beta, \lambda. \quad (24)$$

The Eq. (24) is known as Bayes predictive density function, defined for the parameters α, β and acceleration factor λ separately. No explicit solution of Eq. (24) exist for either parameter. We continue with Eq. (24) numerically for all parameters respectively. The lower and upper Bayes prediction bound limits are denoted by l_1 and l_2 respectively. Thus, the $100(1 - \tau)\%$ BPBL for Z corresponding to the parameter Θ , under the considered censoring scenario is obtained by solving following equation

$$L = \left[P(Z \geq l_2) = \frac{\tau}{2} \right] - \left[P(Z \leq l_1) = \frac{\tau}{2} \right]. \quad (25)$$

Simplifying Eq. (25), we get the non-simplified expressions for the lower and upper Bayes prediction bound limits as

$$\left(1 - \frac{\tau}{2} \right) = \int_{\Theta} (1 - l_1^{-\alpha}) \times \pi_{(\Theta)}^* d\Theta; \Theta = \alpha, \beta, \lambda \quad (26)$$

and

$$\left(\frac{\tau}{2} \right) = \int_{\Theta} (1 - l_2^{-\alpha}) \times \pi_{(\Theta)}^* d\Theta; \Theta = \alpha, \beta, \lambda. \quad (27)$$

Eq. (26) and Eq. (27) has not simplified further for parameters α, β and λ separately. Numerical technique is applied herewith for obtaining the BPBL under the above scenario.

Numerical study based on simulated data

A simulation study by using Metropolis-Hastings algorithm has been carried out for analyzing fruitfulness of considered scenario in this section. Metropolis et al.²⁷ and Hastings²⁸ has deliberated this algorithm which was based on the posterior distribution by use of an arbitrary proposal distribution for simulating samples. Mahmoud et al.²⁹ and Kayal et al., recently explored more about the M-H algorithm. The main objective of the present paper is to assess the effect of combining two different approaches SS-PALT and T-IPH censoring on bound lengths. The numerical findings are computed by using Monte Carlo simulations on 10,000 replications followed by Kayal et al.¹⁷

Thirty test units ($n = 30$) have been fixed for the numerical study in prior. Three different censoring stages 10, 15 and 20 along with four pre-assumed different progressive censoring patterns have been designated for the studying of the behavior of censored sample size and censoring patterns. The pre-assumed progressive censoring schemes are (2, 1, 3, 2, 3, 2, 2, 0, 2, 3), (0, 1, 1, 1, 0, 2, 1, 0, 3, 0, 2, 1, 2, 1, 0), (1, 0, 1, 0, 1, 1, 2, 3, 0, 2, 0, 1, 0, 1, 2) and (0, 1, 0, 1, 0, 1, 0, 0, 2, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0). For perceiving the effect of censoring pattern when other parametric values are fixed a different censoring pattern for $m (= 15)$ has assumed also.

The stress change time $\hat{a}$ is optimized as discussed in the previous section by the method of minimization of the asymptotic variance of ML estimators for α, β and λ . The values for the parameters under study

has assumed for numerical extents as $\alpha = \beta = 0.50$ (0.10) 2.50 = λ . In the present discussion, we also have to find out the optimum values of α, β and λ for which the magnitude of bound lengths maximizes numerically when other parametric values considered to be fixed. Therefore, we assumed all possible combinations of values of these three parameters from 0.50 to 2.50 with an increment of 0.10. The failure time was assumed here as $t = 05, 09$. In the present article, our focus is not only on combining the SS-PALT with T-IPH. We have also studied Type-I censoring & Type-II progressive censoring combined with SS-PALT under similar test situations.

The numerical findings in terms of bound lengths on simulated data are presented in Tables 1-3. The Bayes prediction bound length has been obtained for each parameter for all pre-assumed parametric values as discussed above. From all possible combinations of pre-assumed numerical values of α, β and λ the widest BPBL have noticed for $\alpha = 1.40, \beta = 0.90$ and $\lambda = 1.60$. Hence, the numerical findings are presented here only for three combinations $\alpha = \beta = \lambda = 0.50$, $\alpha = 1.40, \beta = 0.90, \lambda = 1.60$ and $\alpha = \beta = \lambda = 2.50$.

One-sample Bayes prediction bound lengths have been present in Table 03 for selected parametric values as discussed in the above paragraphs. It has been observed that, BPBL getting wider as the level of significance increases. Further, the bound lengths getting narrower when censored sample size m increases. For fixed $m = 15$, two different censoring patterns have been used and it has observed that, the different censoring patterns play a significant role in BPBL. The magnitude of BPBL has observed wider for first censoring pattern when they compared with second one for fixed $m = 15$. The BPBL also has seen wider for small t . It has been already discussed that, the BPBL increases first when values of parameter under consideration increases from $\alpha = \beta = \lambda = 0.50$ and touches maximum at $\alpha = 1.40, \beta = 0.90, \lambda = 1.60$, then the bound lengths decreases for other values of parameter under consideration.

The Table 01 & Table 02 presents the approximate confidence lengths (ACL) and bootstrap confidence lengths (BCL) respectively for selected parametric values. All properties have been seen similar as discussed in above for BPBL. In terms of the bound lengths, the BPBL has seen wider when they compared with BCL and ACL. On the other hand, magnitudes of bootstrap confidence length have seen wider when they compared with ACL. However, for large sample ($m = 20$) the inverse effect has been observed.

In the present discussion, we also appraise two different censoring patterns (Progressive Type-II (PT-II) and Type-I censoring (T-I)) separately under above supposed settings combined with SS-PAT. The numerical findings are obtained for all pre-assumed values; however, avoiding a number of tables, the numerical findings are presented here for selected values only.

Table 04, shows the bound lengths under PT-II censoring on SS-PALT for all parameters under consideration. The numerical findings are reported similar behavior as discussed above in the case of T-IPH censoring pattern. The remarkable point is that, the magnitude of bound lengths was noted narrower for PT-II censoring when they compared with T-IPH censoring pattern (other parametric values assumed to be fixed).

Under T-I censoring scheme on SS-PALT, Table 05 presents the BPBL, BCL & ACL for a total of $n = 30$ units and a fixed censored sample size $m = 15$ with both censoring schemes. The selected time

slot for the analysis are $t = 01, 05, 09, 13$. The numerical findings are reported here only for the selected values as discussed in previous paragraphs. All the properties and behaviors have seen similar discussed in the case of T-IPH and PT-II censoring pattern. Again, it has been observed that, the magnitude of bound lengths under T-IPH

censoring was noted wider as compared with T-I censoring pattern. In terms of bound length, the magnitude of BPBL was observed wider when they compared with ACL or BCL. However, no any clear trends have been observed between BCL and ACL when other parametric values assumed to be fixed.

Table 1 Approximate confidence lengths (ACL) for simulated data

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
α_{Acl}									
0.50	0.50	0.50	10	1.7951	1.7634	1.6895	1.5435	1.4626	1.3823
			15(I)	1.6252	1.6188	1.5533	1.3542	1.3481	1.3023
			15(II)	1.5327	1.5258	1.4431	1.2741	1.2652	1.2078
			20	1.2503	1.2477	1.2247	1.0247	1.0181	0.9522
1.40	0.90	1.60	10	2.5754	2.5302	2.4241	2.2447	2.1323	2.0161
			15(I)	2.3578	2.3481	2.2529	1.9721	1.9623	1.8975
			15(II)	2.2289	2.2196	2.0989	1.8608	1.8386	1.7643
			20	1.8716	1.8634	1.7432	1.5411	1.5289	1.4322
2.50	2.50	2.50	10	1.6616	1.6321	1.5638	1.4285	1.3538	1.2799
			15(I)	1.5038	1.4896	1.4345	1.2533	1.2474	1.2109
			15(II)	1.4186	1.4122	1.3839	1.1792	1.1711	1.1179
			20	1.1578	1.1543	1.1327	0.9477	0.9427	0.8831
β_{Acl}									
0.50	0.50	0.50	10	1.7454	1.7126	1.6417	1.4417	1.3707	1.2962
			15(I)	1.5817	1.5764	1.5116	1.2813	1.2726	1.2334
			15(II)	1.4611	1.4525	1.3747	1.1776	1.1687	1.1157
			20	0.9888	0.9878	0.9689	0.7729	0.7684	0.7187
1.40	0.90	1.60	10	2.8585	2.8064	2.6895	2.4225	2.3006	2.1753
			15(I)	2.6398	2.6301	2.5227	2.1503	2.1409	2.0695
			15(II)	2.4414	2.4292	2.2981	1.9813	1.9571	1.8728
			20	1.7295	1.7229	1.6109	1.3626	1.3523	1.2667
2.50	2.50	2.50	10	1.6161	1.5855	1.5199	1.3394	1.2688	1.1997
			15(I)	1.4629	1.4501	1.3958	1.1852	1.1804	1.1459
			15(II)	1.3528	1.3446	1.3185	1.0999	1.0819	1.0329
			20	0.9152	0.9135	0.8957	0.7146	0.7112	0.6647
λ_{Acl}									
0.50	0.50	0.50	10	1.4666	1.4386	1.3791	1.1736	1.1114	1.0521
			15(I)	1.3369	1.3328	1.2777	1.0523	1.0481	1.0123
			15(II)	1.2023	1.1956	1.1291	0.9412	0.9318	0.8917
			20	0.6789	0.6786	0.6655	0.5057	0.5031	0.4704

Table Continued

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
α_{Acl}									
1.40	0.90	1.60	10	2.7423	2.6927	2.5806	2.2622	2.1482	2.0314
			15(I)	2.5656	2.5563	2.4517	2.0342	2.0255	1.9579
			15(II)	2.3117	2.2998	2.1755	1.8252	1.8025	1.7393
			20	1.3873	1.3823	1.2921	1.0454	1.0376	0.9718
2.50	2.50	2.50	10	1.3581	1.3319	1.2769	1.0865	1.0293	0.9729
			15(I)	1.2364	1.2264	1.1797	0.9733	0.9695	0.9412
			15(II)	1.1143	1.1027	1.0855	0.8714	0.8647	0.8254
			20	0.6282	0.6274	0.6149	0.4674	0.4655	0.4351

Table 2 Bootstrap confidence lengths (BCL) for simulated data

$n = 30$				$t = 05$			$t = 09$					
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$					
α	β	λ	m	99%	95%	90%	99%	95%	90%			
For Parameter α												
0.50	0.50	0.50	10	1.9237	1.8829	1.8102	1.6244	1.5391	1.4546			
			15(I)	1.7414	1.7375	1.6617	1.4328	1.4265	1.3781			
			15(II)	1.6262	1.6118	1.5307	1.3313	1.3217	1.2618			
			20	1.2101	1.2018	1.1855	0.9684	0.9624	0.9443			
			1.40	0.90	1.60	10	2.9493	2.8968	2.7756	2.5335	2.4079	2.2768
						15(I)	2.7132	2.7025	2.5926	2.2394	2.2285	2.1549
15(II)	2.5356	2.5243				2.3874	2.0874	2.0621	1.9789			
			20	1.6957	1.6589	1.6228	1.4762	1.4614	1.3865			
			2.50	2.50	2.50	10	1.7809	1.7485	1.6756	1.5035	1.4247	1.3427
						15(I)	1.6135	1.5987	1.5393	1.3259	1.3198	1.2813
15(II)	1.5053	1.4977				1.4681	1.2323	1.2235	1.1681			
			20	1.1203	1.1173	1.0962	0.8955	0.8921	0.8346			
			For Parameter β									
			0.50	0.50	0.50	10	2.0613	2.0234	1.9394	1.7096	1.6196	1.5306
15(I)	1.8717	1.8652				1.7891	1.5161	1.5095	1.4584			
15(II)	1.7252	1.7157				1.6235	1.3921	1.3808	1.3182			
			20	0.9841	0.9694	0.9474	0.7251	0.7196	0.7006			
			1.40	0.90	1.60	10	3.3771	3.3162	3.1779	2.8627	2.7129	2.5721
						15(I)	3.1227	3.1107	2.9839	2.5432	2.5311	2.4475
15(II)	2.8842	2.8705				2.7152	2.3414	2.3127	2.2194			
			20	1.7195	1.7029	1.5909	1.3526	1.3323	1.2567			

Table Continued

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
For Parameter α									
2.50	2.50	2.50	10	1.9086	1.8743	1.7952	1.5825	1.4992	1.4175
			15(I)	1.7314	1.7146	1.6519	1.4028	1.3965	1.3558
			15(II)	1.5972	1.5883	1.5572	1.2878	1.2783	1.2203
			20	0.9112	0.9035	0.8907	0.7086	0.7012	0.6547
For Parameter λ									
0.50	0.50	0.50	10	1.6272	1.5991	1.5335	1.3033	1.2351	1.1686
			15(I)	1.4875	1.4812	1.4203	1.1704	1.1655	1.1257
			15(II)	1.3342	1.3259	1.2557	1.0451	1.0422	0.9913
			20	0.6598	0.6578	0.6565	0.4997	0.4887	0.4674
1.40	0.90	1.60	10	3.0458	2.9928	2.8691	2.5132	2.3873	2.2568
			15(I)	2.8531	2.8412	2.7215	2.2619	2.2515	2.1758
			15(II)	2.5664	2.5562	2.4187	2.0274	2.0032	1.9253
			20	1.3453	1.3243	1.2791	1.0324	1.0306	0.9688
2.50	2.50	2.50	10	1.5066	1.4804	1.4201	1.2065	1.1437	1.0817
			15(I)	1.3759	1.3631	1.3113	1.0827	1.0781	1.0457
			15(II)	1.2357	1.2306	1.2074	0.9674	0.9607	0.9178
			20	0.6122	0.6094	0.5914	0.4564	0.4555	0.4235

Table 3 Bayes prediction bound lengths for simulated data

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
For Parameter α									
0.50	0.50	0.50	10	2.5541	2.5107	2.4047	2.2368	2.1201	2.0045
			15(I)	2.3126	2.3027	2.2097	1.9541	1.9449	1.8799
			15(II)	2.2023	2.1942	2.0746	1.8597	1.8472	1.7633
			20	1.9728	1.9677	1.9321	1.6553	1.6443	1.5379
1.40	0.90	1.60	10	3.4295	3.3709	3.2289	3.0319	2.8804	2.7234
			15(I)	3.1281	3.1146	2.9889	2.6512	2.6384	2.5504
			15(II)	2.9877	2.9768	2.8141	2.5303	2.5005	2.3995
			20	2.7328	2.7199	2.5452	2.3002	2.2815	2.1372
2.50	2.50	2.50	10	2.3639	2.3235	2.2256	2.0701	1.9621	1.8549
			15(I)	2.1402	2.1193	2.0415	1.8085	1.7999	1.7471
			15(II)	2.0382	2.0307	1.9892	1.7209	1.7098	1.6319
			20	1.8273	1.8209	1.7872	1.5312	1.5227	1.4233
For Parameter β									

Table Continued

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
For Parameter α									
0.50	0.50	0.50	10	3.2855	3.2279	3.0923	2.7762	2.6309	2.4875
			15(I)	2.9889	2.9727	2.8561	2.4545	2.4435	2.3618
			15(II)	2.7765	2.7644	2.6145	2.2749	2.2519	2.1565
			20	2.0752	2.0707	2.0327	1.6599	1.6493	1.5426
1.40	0.90	1.60	10	5.0395	4.9515	4.7437	4.3337	4.1167	3.8924
			15(I)	4.6469	4.6279	4.4403	3.8347	3.8167	3.6895
			15(II)	4.3312	4.3143	4.0794	3.5679	3.5253	3.3831
			20	3.3532	3.3382	3.1223	2.6999	2.6784	2.5089
2.50	2.50	2.50	10	3.0412	2.9875	2.8623	2.5695	2.4349	2.3021
			15(I)	2.7655	2.7395	2.6382	2.2713	2.2611	2.1947
			15(II)	2.5701	2.5587	2.5071	2.1053	2.0911	1.9959
			20	1.9217	1.9159	1.8798	1.5353	1.5271	1.4274
For Parameter λ									
0.50	0.50	0.50	10	3.1472	3.0912	2.9617	2.6123	2.4754	2.3405
			15(I)	2.8703	2.8593	2.7429	2.3237	2.3135	2.2361
			15(II)	2.6323	2.6207	2.4729	2.1251	2.1191	2.0143
			20	1.7976	1.7941	1.7609	1.4039	1.3951	1.3048
1.40	0.90	1.60	10	5.1594	5.0685	4.8561	4.3761	4.1568	3.9303
			15(I)	4.7846	4.7654	4.5719	3.8958	3.8777	3.7485
			15(II)	4.4055	4.3867	4.1482	3.5784	3.5354	3.3928
			20	3.1375	3.1239	2.9221	2.4708	2.4512	2.2961
2.50	2.50	2.50	10	2.9134	2.8611	2.7416	2.4179	2.2921	2.1661
			15(I)	2.6556	2.6321	2.5334	2.1501	2.1406	2.0779
			15(II)	2.4374	2.4258	2.3772	1.9668	1.9532	1.8643
			20	1.6644	1.6598	1.6283	1.2983	1.2916	1.2072

Table 4 Bound lengths for simulated data under Type-II progressive censoring scheme

PT-II	$n = 30$	$t = 05$			$t = 09$		
		$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
Censoring	m	99%	95%	90%	99%	95%	90%
For Parameter α							
BPBL	10	3.1229	3.0691	2.9329	2.7639	2.6047	2.4827
	15(I)	2.8367	2.8225	2.7111	2.4123	2.3998	2.3186
	15(II)	2.7158	2.7053	2.5524	2.3025	2.2658	2.1754
	20	2.4808	2.4587	2.3058	2.0986	2.0808	1.9485

Table Continued

PT-II Censoring	$n = 30$	$t = 05$			$t = 09$		
		$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
	m	99%	95%	90%	99%	95%	90%
For Parameter α							
BCL	10	2.6817	2.6387	2.5275	2.3115	2.1757	2.0772
	15(I)	2.4602	2.4509	2.3513	2.0384	2.0276	1.9595
	15(II)	2.3054	2.2945	2.1625	1.9004	1.8678	1.7935
	20	1.5392	1.4954	1.4683	1.3505	1.3362	1.2669
ACL	10	2.3475	2.3059	2.2083	2.0493	1.9255	1.8405
	15(I)	2.1373	2.1291	2.0429	1.7958	1.7861	1.7258
	15(II)	2.0269	2.0179	1.9031	1.6947	1.6649	1.5987
	20	1.6989	1.6811	1.5776	1.4095	1.3975	1.3084
For Parameter β							
BPBL	10	3.1229	3.0691	2.9329	2.7639	2.6047	2.4827
	15(I)	2.8367	2.8225	2.7111	2.4123	2.3998	2.3186
	15(II)	2.7158	2.7053	2.5524	2.3025	2.2658	2.1754
	20	2.4808	2.4587	2.3058	2.0986	2.0808	1.9485
BCL	10	2.6817	2.6387	2.5275	2.3115	2.1757	2.0772
	15(I)	2.4602	2.4509	2.3513	2.0384	2.0276	1.9595
	15(II)	2.3054	2.2945	2.1625	1.9004	1.8678	1.7935
	20	1.5392	1.4954	1.4683	1.3505	1.3362	1.2669
ACL	10	2.3475	2.3059	2.2083	2.0493	1.9255	1.8405
	15(I)	2.1373	2.1291	2.0429	1.7958	1.7861	1.7258
	15(II)	2.0269	2.0179	1.9031	1.6947	1.6649	1.5987
	20	1.6989	1.6811	1.5776	1.4095	1.3975	1.3084
For Parameter λ							
BPBL	10	3.1229	3.0691	2.9329	2.7639	2.6047	2.4827
	15(I)	2.8367	2.8225	2.7111	2.4123	2.3998	2.3186
	15(II)	2.7158	2.7053	2.5524	2.3025	2.2658	2.1754
	20	2.4808	2.4587	2.3058	2.0986	2.0808	1.9485
BCL	10	2.6817	2.6387	2.5275	2.3115	2.1757	2.0772
	15(I)	2.4602	2.4509	2.3513	2.0384	2.0276	1.9595
	15(II)	2.3054	2.2945	2.1625	1.9004	1.8678	1.7935
	20	1.5392	1.4954	1.4683	1.3505	1.3362	1.2669

Table Continued

PT-II Censoring	$n = 30$	$t = 05$			$t = 09$		
		$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
	m	99%	95%	90%	99%	95%	90%
For Parameter α							
ACL	10	2.3475	2.3059	2.2083	2.0493	1.9255	1.8405
	15(I)	2.1373	2.1291	2.0429	1.7958	1.7861	1.7258
	15(II)	2.0269	2.0179	1.9031	1.6947	1.6649	1.5987
	20	1.6989	1.6811	1.5776	1.4095	1.3975	1.3084

Table 5 Bound lengths for simulated data under Type-I censoring scheme

T-I Censoring	$n = 30$	$m = 15(I)$			$m = 15(II)$		
		$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
	$t \downarrow$	99%	95%	90%	99%	95%	90%
For Parameter α							
BPBL	01	2.5235	2.3571	2.2668	2.2233	2.1845	2.0868
	05	2.6512	2.6384	2.5504	2.5303	2.5005	2.3995
	09	3.1281	3.1146	2.9889	2.9877	2.9768	2.8141
	13	1.9178	1.9007	1.7792	2.2546	2.2241	2.0909
BCL	01	2.1123	1.9672	1.8982	1.9105	1.8794	1.7994
	05	2.2394	2.2285	2.1549	2.0874	2.0621	1.9789
	09	2.7132	2.7025	2.5926	2.5356	2.5243	2.3874
	13	1.2378	1.2239	1.1596	1.3988	1.3485	1.3296
ACL	01	1.9674	1.9516	1.8855	1.6517	1.6272	1.5575
	05	1.9721	1.9623	1.8975	1.8608	1.8386	1.7643
	09	2.3578	2.3481	2.2529	2.2289	2.2196	2.0989
	13	1.4465	1.4334	1.3413	1.7308	1.7023	1.6025
For Parameter β							
BPBL	01	2.5235	2.3571	2.2668	2.2233	2.1845	2.0868
	05	2.6512	2.6384	2.5504	2.5303	2.5005	2.3995
	09	3.1281	3.1146	2.9889	2.9877	2.9768	2.8141
	13	1.9178	1.9007	1.7792	2.2546	2.2241	2.0909
BCL	01	2.1123	1.9672	1.8982	1.9105	1.8794	1.7994
	05	2.2394	2.2285	2.1549	2.0874	2.0621	1.9789
	09	2.7132	2.7025	2.5926	2.5356	2.5243	2.3874
	13	1.2378	1.2239	1.1596	1.3988	1.3485	1.3296
ACL	01	1.9674	1.9516	1.8855	1.6517	1.6272	1.5575
	05	1.9721	1.9623	1.8975	1.8608	1.8386	1.7643
	09	2.3578	2.3481	2.2529	2.2289	2.2196	2.0989
	13	1.4465	1.4334	1.3413	1.7308	1.7023	1.6025

Table Continued

T-I Censoring	$n = 30$	$m = 15(I)$			$m = 15(II)$		
		$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
	$t \downarrow$	99%	95%	90%	99%	95%	90%
For Parameter λ							
BPBL	01	2.5235	2.3571	2.2668	2.2233	2.1845	2.0868
	05	2.6512	2.6384	2.5504	2.5303	2.5005	2.3995
	09	3.1281	3.1146	2.9889	2.9877	2.9768	2.8141
	13	1.9178	1.9007	1.7792	2.2546	2.2241	2.0909
BCL	01	2.1123	1.9672	1.8982	1.9105	1.8794	1.7994
	05	2.2394	2.2285	2.1549	2.0874	2.0621	1.9789
	09	2.7132	2.7025	2.5926	2.5356	2.5243	2.3874
	13	1.2378	1.2239	1.1596	1.3988	1.3485	1.3296
ACL	01	1.9674	1.9516	1.8855	1.6517	1.6272	1.5575
	05	1.9721	1.9623	1.8975	1.8608	1.8386	1.7643
	09	2.3578	2.3481	2.2529	2.2289	2.2196	2.0989
	13	1.4465	1.4334	1.3413	1.7308	1.7023	1.6025

Table 6 Approximate confidence lengths (ACL) for real data

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
α_{Acl}									
0.50	0.50	0.50	10	1.8186	1.7816	1.7103	1.5617	1.4637	1.4033
			15(I)	1.6347	1.6289	1.5631	1.3702	1.3631	1.3157
			15(II)	1.5481	1.5405	1.4512	1.2895	1.2708	1.2142
			20	1.2599	1.2469	1.2294	1.0431	1.0355	0.9678
1.40	0.90	1.60	10	2.6015	2.5589	2.4507	2.2737	2.1387	2.0421
			15(I)	2.3731	2.3639	2.2682	1.9913	1.9822	1.9156
			15(II)	2.2498	2.2398	2.1113	1.8808	1.8487	1.7751
			20	1.8861	1.8674	1.7512	1.5636	1.5504	1.4516
2.50	2.50	2.50	10	1.6814	1.6537	1.5836	1.4511	1.3541	1.3001
			15(I)	1.5124	1.4987	1.4433	1.2685	1.2616	1.2236
			15(II)	1.4331	1.4216	1.3923	1.1938	1.1759	1.1236
			20	1.1666	1.1527	1.1366	0.9655	0.9595	0.8982
β_{Acl}									
0.50	0.50	0.50	10	1.7859	1.7519	1.6786	1.4788	1.3848	1.3295
			15(I)	1.6067	1.6019	1.5362	1.3095	1.2998	1.2586
			15(II)	1.4905	1.4812	1.3968	1.2014	1.1852	1.1326
			20	1.0062	0.9948	0.9812	0.7917	0.7916	0.7397

Table Continued

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
1.40	0.90	1.60	10	2.9119	2.8653	2.7451	2.4772	2.3314	2.2243
			15(I)	2.6837	2.6745	2.5653	2.1941	2.1836	2.1096
			15(II)	2.4884	2.4754	2.3367	2.0221	1.9877	1.9032
			20	1.7601	1.7413	1.6347	1.3973	1.3859	1.2975
2.50	2.50	2.50	10	1.6543	1.6226	1.5546	1.3747	1.2811	1.2313
			15(I)	1.4858	1.4733	1.4183	1.2117	1.2059	1.1695
			15(II)	1.3803	1.3714	1.3396	1.1249	1.0969	1.0483
			20	0.9313	0.9191	0.9067	0.7377	0.7333	0.6847
λ_{Acl}									
0.50	0.50	0.50	10	1.5315	1.5018	1.4389	1.2294	1.1431	1.1021
			15(I)	1.3843	1.3806	1.3236	1.0975	1.0922	1.0538
			15(II)	1.2512	1.2436	1.1694	0.9822	0.9627	0.9224
			20	0.7043	0.6936	0.6857	0.5352	0.5316	0.4963
1.40	0.90	1.60	10	2.8555	2.8034	2.6859	2.3592	2.2192	2.1185
			15(I)	2.6595	2.6505	2.5421	2.1166	2.1067	2.0352
			15(II)	2.4026	2.3897	2.2554	1.8997	1.8664	1.8021
			20	1.4396	1.4242	1.3361	1.0953	1.0863	1.0167
2.50	2.50	2.50	10	1.4189	1.3911	1.3328	1.1319	1.0579	1.0199
			15(I)	1.2801	1.2702	1.2219	1.0155	1.0106	0.9809
			15(II)	1.1598	1.1472	1.1241	0.9097	0.8931	0.8536
			20	0.6517	0.6405	0.6332	0.4954	0.4925	0.4597

Table 7 Bootstrap confidence lengths (BCL) for real data

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
For Parameter α									
0.50	0.50	0.50	10	1.9867	1.9441	1.8682	1.6811	1.5716	1.5053
			15(I)	1.7867	1.7833	1.7056	1.4781	1.4707	1.4196
			15(II)	1.6749	1.6595	1.5709	1.3737	1.3542	1.2939
			20	1.2436	1.2246	1.2136	1.0057	0.9986	0.9787
1.40	0.90	1.60	10	3.0409	2.9863	2.8605	2.6155	2.4647	2.3504
			15(I)	2.7856	2.7752	2.6624	2.3072	2.2951	2.2181
			15(II)	2.6096	2.5974	2.4515	2.1509	2.1152	2.0311
			20	1.7427	1.6945	1.6631	1.5277	1.5116	1.4333
2.50	2.50	2.50	10	1.8399	1.8016	1.7298	1.5567	1.4514	1.3903
			15(I)	1.6552	1.6406	1.5797	1.3682	1.3611	1.3201
			15(II)	1.5506	1.5422	1.5066	1.2712	1.2532	1.1976
			20	1.1513	1.1378	1.1218	0.9308	0.9264	0.8616

Table Continued

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
For Parameter β									
0.50	0.50	0.50	10	2.1378	2.0904	2.0027	1.7701	1.6558	1.5848
			15(I)	1.9223	1.9162	1.8381	1.5651	1.5574	1.5035
			15(II)	1.7782	1.7678	1.6677	1.4375	1.4162	1.3531
			20	1.0121	0.9866	0.9697	0.7563	0.7497	0.7289
			10	3.4836	3.4204	3.2769	2.9564	2.7806	2.6563
			15(I)	3.2093	3.1976	3.0673	2.6217	2.6084	2.5211
1.40	0.90	1.60	15(II)	2.9706	2.9559	2.7909	2.4141	2.3749	2.2802
			20	1.7687	1.7412	1.6317	1.4018	1.3801	1.3011
			10	1.9729	1.9317	1.8544	1.6394	1.5312	1.4684
2.50	2.50	2.50	15(I)	1.7718	1.7613	1.6917	1.4485	1.4411	1.3979
			15(II)	1.6465	1.6367	1.5995	1.3302	1.3107	1.2523
			20	0.9371	0.9188	0.9113	0.7393	0.7308	0.6817
For Parameter λ									
0.50	0.50	0.50	10	2.1378	2.0904	2.0027	1.7701	1.6558	1.5848
			15(I)	1.9223	1.9162	1.8381	1.5651	1.5574	1.5035
			15(II)	1.7782	1.7678	1.6677	1.4375	1.4162	1.3531
			20	1.0121	0.9866	0.9697	0.7563	0.7497	0.7289
			10	3.4836	3.4204	3.2769	2.9564	2.7806	2.6563
			15(I)	3.2093	3.1976	3.0673	2.6217	2.6084	2.5211
1.40	0.90	1.60	15(II)	2.9706	2.9559	2.7909	2.4141	2.3749	2.2802
			20	1.7687	1.7412	1.6317	1.4018	1.3801	1.3011
			10	1.9729	1.9317	1.8544	1.6394	1.5312	1.4684
2.50	2.50	2.50	15(I)	1.7718	1.7613	1.6917	1.4485	1.4411	1.3979
			15(II)	1.6465	1.6367	1.5995	1.3302	1.3107	1.2523
			20	0.9371	0.9188	0.9113	0.7393	0.7308	0.6817

Table 8 Bayes prediction bound lengths for real data

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
For Parameter α									
0.50	0.50	0.50	10	2.9156	2.8656	2.7438	2.5565	2.4021	2.2911
			15(I)	2.6282	2.6175	2.5119	2.2289	2.2175	2.1422
			15(II)	2.5093	2.4995	2.3582	2.1214	2.0975	2.0034
			20	2.2445	2.2283	2.1935	1.8939	1.8804	1.7581

Table Continued

$n = 30$				$t = 05$			$t = 09$		
				$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
α	β	λ	m	99%	95%	90%	99%	95%	90%
1.40	0.90	1.60	10	3.9117	3.8444	3.6816	3.4613	3.2672	3.1019
			15(I)	3.5561	3.5414	3.3986	3.0221	3.0066	2.9052
			15(II)	3.4031	3.3921	3.1997	2.8845	2.8409	2.7273
			20	3.1093	3.0843	2.8912	2.6277	2.6055	2.4401
2.50	2.50	2.50	10	2.6992	2.6526	2.5434	2.3669	2.2223	2.1208
			15(I)	2.4322	2.4088	2.3205	2.0632	2.0525	1.9911
			15(II)	2.3226	2.3134	2.2613	1.9635	1.9412	1.8538
			20	2.0792	2.0613	2.0286	1.7526	1.7421	1.6277
For Parameter β									
0.50	0.50	0.50	10	5.0621	4.9729	4.7631	4.2808	4.0356	3.8356
			15(I)	4.5933	4.5619	4.3899	3.7801	3.7622	3.6353
			15(II)	4.2733	4.2541	4.0183	3.5039	3.4588	3.3134
			20	3.1911	3.1738	3.1211	2.5631	2.5459	2.3805
1.40	0.90	1.60	10	7.7595	7.6236	7.3028	6.6761	6.3207	5.9962
			15(I)	7.1431	7.1145	6.8262	5.9027	5.8741	5.6772
			15(II)	6.6642	6.6376	6.2712	5.4924	5.4171	5.1998
			20	5.1566	5.1231	4.7968	4.1625	4.1285	3.8665
2.50	2.50	2.50	10	4.6863	4.6032	4.4094	3.9629	3.7342	3.5505
			15(I)	4.2497	4.2104	4.0548	3.4983	3.4817	3.3783
			15(II)	3.9558	3.9377	3.8532	3.2413	3.2115	3.0664
			20	2.9551	2.9358	2.8859	2.3714	2.3579	2.2033
For Parameter λ									
0.50	0.50	0.50	10	3.4994	3.4367	3.2919	2.9082	2.7347	2.6056
			15(I)	3.1797	3.1681	3.0392	2.5822	2.5709	2.4828
			15(II)	2.9224	2.9089	2.7398	2.3619	2.3456	2.2307
			20	1.9931	1.9789	1.9478	1.5672	1.5565	1.4551
1.40	0.90	1.60	10	5.7308	5.6294	5.3927	4.8642	4.5993	4.3686
			15(I)	5.3026	5.2819	5.0675	4.3255	4.3046	4.1601
			15(II)	4.8888	4.8673	4.5976	3.9736	3.9162	3.7593
			20	3.4791	3.4535	3.2355	2.7503	2.7277	2.5544
2.50	2.50	2.50	10	3.2401	3.1815	3.0478	2.6926	2.5314	2.4122
			15(I)	2.9416	2.9162	2.8069	2.3897	2.3782	2.3074
			15(II)	2.7063	2.6928	2.6337	2.1864	2.1616	2.0643
			20	1.8454	1.8299	1.8007	1.4501	1.4417	1.3468

Table 9 Bound lengths for real data under Type-II progressive censoring scheme

PT-II Censoring	$n = 30$ m	$t = 05$			$t = 09$		
		$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
		99%	95%	90%	99%	95%	90%
For Parameter α							
BPBL	10	3.5607	3.4939	3.3501	3.1538	2.9559	2.8263
	15(I)	3.2253	3.2125	3.0831	2.7419	2.7341	2.6407
	15(II)	3.0933	3.0824	2.9025	2.6241	2.5748	2.4723
	20	2.8226	2.7895	2.6199	2.3926	2.3749	2.2235
BCL	10	2.7701	2.7199	2.6045	2.3859	2.2273	2.1424
	15(I)	2.5257	2.5169	2.4147	2.1022	2.0881	2.0169
	15(II)	2.3725	2.3609	2.2232	1.9581	1.9136	1.8409
	20	1.5819	1.5277	1.5049	1.3973	1.3818	1.3094
ACL	10	2.3712	2.3319	2.2325	2.0756	1.9313	1.8641
	15(I)	2.1512	2.1435	2.0568	1.8132	1.8034	1.7423
	15(II)	2.0459	2.0362	1.9143	1.7129	1.6724	1.6085
	20	1.7121	1.6847	1.5849	1.4299	1.4117	1.3226
For Parameter β							
BPBL	10	3.5607	3.4939	3.3501	3.1538	2.9559	2.8263
	15(I)	3.2253	3.2125	3.0831	2.7419	2.7341	2.6407
	15(II)	3.0933	3.0824	2.9025	2.6241	2.5748	2.4723
	20	2.8226	2.7895	2.6199	2.3926	2.3749	2.2235
BCL	10	2.7701	2.7199	2.6045	2.3859	2.2273	2.1424
	15(I)	2.5257	2.5169	2.4147	2.1022	2.0881	2.0169
	15(II)	2.3725	2.3609	2.2232	1.9581	1.9136	1.8409
	20	1.5819	1.5277	1.5049	1.3973	1.3818	1.3094
ACL	10	2.3712	2.3319	2.2325	2.0756	1.9313	1.8641
	15(I)	2.1512	2.1435	2.0568	1.8132	1.8034	1.7423
	15(II)	2.0459	2.0362	1.9143	1.7129	1.6724	1.6085
	20	1.7121	1.6847	1.5849	1.4299	1.4117	1.3226
For Parameter λ							
BPBL	10	3.5607	3.4939	3.3501	3.1538	2.9559	2.8263
	15(I)	3.2253	3.2125	3.0831	2.7419	2.7341	2.6407
	15(II)	3.0933	3.0824	2.9025	2.6241	2.5748	2.4723
	20	2.8226	2.7895	2.6199	2.3926	2.3749	2.2235
BCL	10	2.7701	2.7199	2.6045	2.3859	2.2273	2.1424
	15(I)	2.5257	2.5169	2.4147	2.1022	2.0881	2.0169
	15(II)	2.3725	2.3609	2.2232	1.9581	1.9136	1.8409
	20	1.5819	1.5277	1.5049	1.3973	1.3818	1.3094
ACL	10	2.3712	2.3319	2.2325	2.0756	1.9313	1.8641
	15(I)	2.1512	2.1435	2.0568	1.8132	1.8034	1.7423
	15(II)	2.0459	2.0362	1.9143	1.7129	1.6724	1.6085
	20	1.7121	1.6847	1.5849	1.4299	1.4117	1.3226

Table 10 Bound lengths for real data under Type-I censoring scheme

T-I Censoring	$n = 30$	$t = 05$			$t = 09$		
		$\leftarrow \tau \rightarrow$			$\leftarrow \tau \rightarrow$		
	m	99%	95%	90%	99%	95%	90%
For Parameter α							
BPBL	10	2.8666	2.6868	2.5639	2.5244	2.4747	2.3751
	15(I)	3.0221	3.0066	2.9052	2.8845	2.8409	2.7273
	15(II)	3.5561	3.5414	3.3986	3.4031	3.3921	3.1997
	20	2.1748	2.1587	2.0211	2.5656	2.5355	2.3814
BCL	10	2.1687	2.0245	1.9473	1.9639	1.9283	1.8465
	15(I)	2.3072	2.2951	2.2181	2.1509	2.1152	2.0311
	15(II)	2.7856	2.7752	2.6624	2.6096	2.5974	2.4515
	20	1.2701	1.2546	1.1902	1.4379	1.3886	1.3679
ACL	10	1.9866	1.9679	1.8994	1.6643	1.6367	1.5667
	15(I)	1.9913	1.9822	1.9156	1.8808	1.8487	1.7751
	15(II)	2.3731	2.3639	2.2682	2.2498	2.2398	2.1113
	20	1.4527	1.4385	1.3477	1.7445	1.7166	1.6149
For Parameter β							
BPBL	10	2.8666	2.6868	2.5639	2.5244	2.4747	2.3751
	15(I)	3.0221	3.0066	2.9052	2.8845	2.8409	2.7273
	15(II)	3.5561	3.5414	3.3986	3.4031	3.3921	3.1997
	20	2.1748	2.1587	2.0211	2.5656	2.5355	2.3814
BCL	10	2.1687	2.0245	1.9473	1.9639	1.9283	1.8465
	15(I)	2.3072	2.2951	2.2181	2.1509	2.1152	2.0311
	15(II)	2.7856	2.7752	2.6624	2.6096	2.5974	2.4515
	20	1.2701	1.2546	1.1902	1.4379	1.3886	1.3679
ACL	10	1.9866	1.9679	1.8994	1.6643	1.6367	1.5667
	15(I)	1.9913	1.9822	1.9156	1.8808	1.8487	1.7751
	15(II)	2.3731	2.3639	2.2682	2.2498	2.2398	2.1113
	20	1.4527	1.4385	1.3477	1.7445	1.7166	1.6149
For Parameter λ							
BPBL	10	2.8666	2.6868	2.5639	2.5244	2.4747	2.3751
	15(I)	3.0221	3.0066	2.9052	2.8845	2.8409	2.7273
	15(II)	3.5561	3.5414	3.3986	3.4031	3.3921	3.1997
	20	2.1748	2.1587	2.0211	2.5656	2.5355	2.3814
BCL	10	2.1687	2.0245	1.9473	1.9639	1.9283	1.8465
	15(I)	2.3072	2.2951	2.2181	2.1509	2.1152	2.0311
	15(II)	2.7856	2.7752	2.6624	2.6096	2.5974	2.4515
	20	1.2701	1.2546	1.1902	1.4379	1.3886	1.3679
ACL	10	1.9866	1.9679	1.8994	1.6643	1.6367	1.5667
	15(I)	1.9913	1.9822	1.9156	1.8808	1.8487	1.7751
	15(II)	2.3731	2.3639	2.2682	2.2498	2.2398	2.1113
	20	1.4527	1.4385	1.3477	1.7445	1.7166	1.6149

Numerical illustration

In this section, the numerical illustration was based on the dataset provided by Box & Cox³⁰ for survival time of animals observed due to different dosage of poison administered. The observations are listed as

0.18 0.21 0.22 0.22 0.23 0.23 0.23 0.24 0.25 0.29 0.29 0.30 0.30
0.31 0.31 0.31 0.33 0.35 0.36 0.36 0.37 0.38 0.38 0.40 0.40 0.43 0.43
0.44 0.45 0.45 0.45 0.46 0.49 0.56 0.61 0.62 0.63 0.66 0.71 0.71 0.72
0.76 0.82 0.88 0.92 1.02 1.10 1.24.

The Kolmogorov-Smirnov (KS) test (test statistic value = 0.1440 with p-value 0.2724) was afforded a satisfactory fit the underlying distribution of this data set. For the numerical illustration, a set of 30 (= n) randomly selected data from given dataset was assumed. For simplicity in comparison, similar censoring patterns have used for considering censored sample size with all above pre-assumed parametric values. The bound lengths ACL, BCL and BPBL have been presented in Tables 6-10 respectively for the parameters under study. All the properties and trends have seen similar as discussed above in the case of simulated data. The remarkable point is that, the narrower bound lengths have noticed for simulated data set when they compared with the corresponding real dataset under all considered censoring criterion and censoring patterns for all parameters under study.

Summary

No any literature has noticed in studying the properties of the bound length by using SS-PALT and T-IPH censoring. The present article motivated for that. In the present article, we focused not only for combining the SS-PALT with T-IPH censoring. We are also trying to develop the scenario for combining SS-PALT with Type-I censoring and progressive Type-II censoring pattern, respectively and studying the fruitfulness in terms of different bound lengths. The Bayes predictive bound length (BPBL), bootstrap bound length (BCL) and approximate confidence lengths (ACL) has been observed on several combinations of the parametric values. Wider BPBL has been observed in T-IPH censoring over others on both real and simulated datasets. On real and simulated data set, the magnitude of BPBL was seen wider for real data set. The present discussion showed that, the T-IPH censoring maximizes the bound lengths over T-I or PT-II censoring.

Conflicts of interest

There are no conflicts of interest.

Acknowledgement

None.

References

1. Burr WI. Cumulative frequency distribution. *Annals of Mathematical Statistics*. 1942;13:215–232.
2. Abd-Elfattah AM, Hassan AS, Nassr SG. Estimation in step-stress partially accelerated life tests for the Burr Type-XII distribution using Type-I censoring. *Statistical Methodology*. 2008;5(6):502–514.
3. Khalaf EA, Zeinmum J, Heba SM. Finite mixture of Burr Type-XII distribution and its reciprocal: Properties and applications. *Statistical Papers*. 2011;52:835–845.
4. Soliman AA, Abd-Allah AH, Abou-Elheggag NA. et al. Bayesian inference and prediction of Burr Type-XII distribution for progressive first failure censored sampling. *Intelligent Information Management*. 2011;3:175–185.
5. Jang DH, Jung M, Park JH, et al. Bayesian estimation of Burr Type-XII distribution based on general progressive Type-II censoring. *Applied Mathematical Sciences*. 2014;8:3435–3448.
6. Rao GS, Aslam M, Kundu D. Burr-XII distribution parametric estimation and estimation of reliability of multicomponent stress-strength. *Communications in Statistics-Theory & Methods*. 2015;4:4953–4961.
7. Panahi H, Sayyareh A. Estimation and prediction for a unified hybrid-censored Burr Type-XII distribution. *Journal of Statistical Computation & Simulation*. 2015;86(1):55–73.
8. Prakash G. Confidence limits for progressive censored Burr Type-XII data under CP-ALT. *Journal of Statistics Applications & Probability*. 2017;6(2):295–303.
9. Asl MN, Belaghi RA, Bevrani H. On Burr XII distribution analysis under progressive Type-II hybrid censored data. *Methodology & Computing in Applied Probability*. 2017;9(2):665–683.
10. Prakash G. Bayes estimation for C-PALT progressively censored Burr-XII data. *International Journal of Intelligent Technologies & Applied Statistics*. 2018a;11(1): 01–14.
11. Prakash G. Progressively censored Rayleigh data under Bayesian estimation. *The International Journal of Intelligent Technologies & Applied Statistics*. 2015;8(3):257–273.
12. Lin CT, Ng HKT, Chan PS. Statistical inference of Type-II progressively hybrid censored data with Weibull lifetimes. *Communications in Statistics-Theory & Methods*. 2009;38:1710–1729.
13. Lin CT, Huang YL. On progressive hybrid censored Exponential distribution. *Journal of Statistical Computation & Simulation*. 2012;82(5):689–709.
14. Singh B, Gupta PK, Sharma VK. On Type-II hybrid censored Lindley distribution. *Statistics Research Letters*. 2014;3:58–62.
15. Balakrishnan N, Cramer E. The art of progressive censoring: Applications to reliability and quality. Birkhäuser, Boston; 2014.
16. Elshahhat A, Ashour S. Bayesian and non-Bayesian estimation for Weibull parameters based on generalized Type-II progressive hybrid censoring scheme. *Pakistan Journal of Statistics & Operation Research*. 2016;12(2):213–226.
17. Kayal T, Tripathi YM, Rastogi MK, et al. Inference for Burr XII distribution under Type-I progressive hybrid censoring. *Communication in Statistics-Simulation & Computation*. 2017;46(9):7447–7465.
18. Nelson W. Accelerated Life Testing: Statistical Models, Test Plans and Data Analysis. John Wiley & Sons, New York; 1990.
19. Prakash G. Generalized inverted Exponential distribution on Optimum SS-PALT: Some Bayes estimation. *Journal of Statistics Applications & Probability*. 2018b;7(3):457–467.
20. DeGroot MH, Goel PK. Bayesian and optimal design in partially accelerated life testing. *Naval Research Logistics*. 1979;16 (2):223–235.
21. Ismail AA, Aly H M. Optimal planning of failure step-stress partially accelerated life tests under Type-II censoring. *International Journal of Mathematical Analysis*. 2009;3(31):1509–1523.
22. Mathcad. Statistical package. 2001.
23. Efron B, Tibshirani R. Bootstrap methods for standard errors, confidence intervals, and other measures of statistical accuracy. *Statistical Science*. 1986;1(1):54–77.
24. Kreiss JP, Paparoditis E. Bootstrap methods for dependent data: a review. *Journal of Korean Statistical Society*. 2011;40:357–378.
25. Kundu D, Kannan N, Balakrishnan N. *Analysis of Progressively censored*

- competing risks data. In: Balakrishnan N, Rao CR, editors. Handbook of statistics. 2004;23:331–348.
26. Al-Hussaini EK. Predicting observables from a general class of distributions. *Journal of Statistical Planning & Inference*. 1999;79:79–91.
27. Metropolis N, Rosenbluth AW, Rosenbluth MN, et al. Equations of state calculations by fast computing machines. *Journal of Chemical Physics*. 1953;21:1087–1092.
28. Hastings WK. Monte Carlo sampling methods using Markov Chains and their applications. *Biometrika*. 1970;57:97–109.
29. Mahmoud MAW, EL-Sagheer RM, Nagaty H. Inference for constant-stress partially accelerated life test model with progressive Type-II censoring scheme. *Journal of Statistics Applications Proc*. 2017;6(2):373–383.
30. Box GEP, Cox DR. An analysis of transformations (with discussion). *Journal of the Royal Statistical Society-B*. 1964;26:211–252.