

New record of the deep-sea jellyfish *Poralia rufescens* vanhoeffen, 1902 (Class: Scyphozoa) in the Colombian Southern Caribbean

Abstract

Poralia rufescens, Vanhöffen, 1902, is a bathypelagic *Semaestomeae* jellyfish that lives at depths below 600m in the Caribbean; this is the first report of this species and in general of a jellyfish in deep waters of the Colombian Caribbean and spotted in its natural environment thanks to ROV video recordings.

Keywords: *Poralia*, ROV video survey, ultra-deep waters, Colombian Caribbean

Volume 3 Issue 1 - 2019

Cristina Cedeño-Posso,¹ Luisa F Dueñas,^{2,3}
Jorge León,² Vladimir Puentes²

¹Instituto de Investigaciones Marinas y Costeras, INVEMAR, Colombia

²Anadarko Colombia Company-HSE, Colombia

³Departamento de Biología, Facultad de Ciencias, Universidad Nacional de Colombia, Colombia

Correspondence: Vladimir Puentes, Anadarko Colombia Company-HSE, Calle 113 No. 7-80 Piso 11, Bogotá DC, Colombia, Email Vladimir.Puentes@anadarko.com, zanclus0715@gmail.com

Received: December 06, 2018 | **Published:** January 08, 2019

Introduction

Hydrocarbon exploratory drilling activities normally include ROV video surveys and towed camera transects of sites that are preselected for locating wells. During the exploratory activities between 2015 and 2017, jellyfish from the genus *Poralia* were sighted for the first time in deep waters of the Southwestern Colombian Caribbean, below 1500 m. ROV surveys were conducted in 80-100meter-long transects in cross fashion, with headings north, south, east, and west, from a

central point. All video footage and still images were analyzed for the presence of *Poralia*, and, when found in the ROV videos, snapshots were taken using the software GOM player and VLC media player. For every visual confirmation, we registered coordinates, depth, time and date (Table 1).¹

Seven (7) records of the jellyfish from the genus *Poralia* were identified from ROV videos (Figure 1) at a depth between 1565 and 1816m.




Figure 1 Example of images obtained for *Poralia* sp. (A-D) during deep-sea hydrocarbon exploratory activities in the Southwestern Colombian Caribbean. Video-captures of *Poralia* sp. at (A) 1814m, (B) 1816m, (C) 1564m, and (D) 1807m deep.

Table 1 Information on *Poralia* sp. sightings during deep-sea hydrocarbon exploratory activities using ROV surveys

Image	Coordinates		Depth (m)	Time	Date mm/dd/yyyy
	Lat (N)	Long (W)			
ROV_2015_KronosDep_0008	9.165	-76.832	1565	9:48	8/27/2015
ROV_2017_OldPADep_0009	9.207	-76.832	1805	16:27	2/24/2017
ROV_2017_OldPADep_0011	9.207	-76.832	1809	16:19	2/24/2017
ROV_2017_NewPADep_0014	9.209	-76.832	1818	14:58	2/24/2017
ROV_2017_NewPADep_0012	9.209	-76.832	1816	15:11	2/24/2017
ROV_2017_NewPADep_0013	9.209	-76.832	1816	15:12	2/24/2017
ROV_2017_NewPADep_0017	9.209	-76.832	1814	15:36	2/24/2017

Phylum: *Cnidaria*

Class: *Scyphozoa*

Subclass: *Discomedusae*

Order: *Semaeostomeae*

Family: *Ulmaridae*

Subfamily: *Poraliinae*

Genus: *Poralia*

Species: *Poralia rufescens*

Vanhöffen, 1902

Five species of *Scyphozoa* jellyfish are known for the Colombian Caribbean, *Chrysaora* sp., *Aurelia* sp., *Cassiopea* sp., *Lychnorhiza* sp. aff. *lucerna* and *Stomolophus meleagris* most of them from epipelagic waters.^{2,3} This is the first report from a jellyfish in deeper waters from the Colombian Caribbean. *Poralia* jellyfish (*Scyphozoa*: *Semaeostomeae*: *Ulmaridae*) were seen in ROV video surveys (Table 1). *Poralia* belong to the subfamily *Poraliinae* with only one genus and one species: *Poralia rufescens*. In ROV videos *Poralia* is easy to identify by its distinguishing characteristics as are the vertical orientation in the water column, the bell's closed posture, the round ball shape and its reddish brown color typical of deep-sea jellyfish. Besides, it has a slow swim with relatively short tentacles extended out to the side and/or below the bell.^{4,5} Although the identification based on still images is only possible if the diagnostic characters of the species are visible and distinguishable, it was possible to get confirmation of this species thanks to the review of some of its characters (Allen Collins, Smithsonian Institute, Washington DC, personal communication).

The scyphomedusae *Poralia* is a bathypelagic jellyfish that lives below 600m in close association with the benthic boundary layer. *Poralia* has been reported in the Western Atlantic (Bahamas, Dry Tortugas and New England)^{6,7} and in the Gulf of Mexico.⁸ Osborn et al.⁵ analyzed the environmental data where mesopelagic scyphomedusae occur, and their results concluded that *Poralia* was found more often in waters coldest, saltier (34.40ppt) and with lower oxygen concentration (< 0.3ml/l).

Funding details

This research was funded partially by Anadarko Colombia Company and Colciencias' Post-Doctoral Program (Fondo Nacional de Financiamiento para la Ciencia, la Tecnología y la Innovación "Francisco José de Caldas" - COLCIENCIAS. Convocatoria 2017, Número 784)

Acknowledgments

Thanks to Anadarko Colombia Company, a subsidiary of Anadarko Petroleum Company, that provided funding for the exploratory drilling projects, together with Ecopetrol. Other companies involved in the project were Dolphin Drilling that operated the Mobile Operating Drilling Unit, from where the ROV was launched; Subsea 7 (and its division I-TECH services) provided the ROV services. We thank the I-TECH Services technical staff that conducted the ROV surveys, for their willingness, time and patience that made possible the images that led us to this work. Thanks to Allen Collins of the National Systematics Laboratory of NOAA and Smithsonian Institution, for his comments and confirming the ID of these specimens. Contribution No. 1214 of the Instituto de Investigaciones Marinas y Costeras – INVEMAR.

Conflicts of interest

Authors declare that there is no conflicts of interest.

References

1. Dueñas LF, Puentes V. First visual occurrence data for deep-sea cnidarians in the South-western Colombian Caribbean. Anadarko Colombia Company. 2018.
2. Cedeño-Posso C, Lecompte O. Primer registro de medusas del género *Lychnorhiza* (Cnidaria: Scyphozoa: Rhizostomeae: Lychnorhizidae) en el mar Caribe colombiano. *Boletín de Investigaciones Marinas y Costeras*. 2013;42(2):387–393.
3. Oliveira OMP, Araújo EM, Ayon P, et al. Census of Cnidaria (Medusozoa) and Ctenophora from South American marine waters. *Zootaxa*. 2016;4194:1–256.
4. Russel F. On the scyphomedusa *Poralia rufescens* Vanhöffen. *Journal of the Marine Biological Association of the United Kingdom*. 1962;42(2):387–390.
5. Osborn DA, Silver MW, Castro CG, et al. The habitat of mesopelagic scyphomedusae in Monterey Bay, California. *Deep-Sea Research I*. 2007;54(8):1241–1255.
6. Larson RJ, Mills CE, Harbison GR. Western Atlantic midwater hydrozoan and scyphozoan medusae: in situ studies using manned submersibles. *Hydrobiologia*, 1991;216(1):311–317.
7. Larson RJ, Matsumoto GI, Madin LP, et al. Deep-sea benthic and bathypelagic medusae: recent observations from submersibles and a remotely operated vehicle. *Bulletin of Marine Science*. 1992;51(3):277–286.
8. Valentine M, Benfield M. Characterization of epibenthic and demersal megafauna at Mississippi Canyon 252 following the Deepwater Horizon Oil Spill. *Marine Pollution Bulletin*. 2013;779(1–2):196–209.