

Selective Immunoglobulin M Deficiency and Terminal Ileitis due to *Yersinia Enterocolitica* Infection: A Clinically Interesting Coexistence

Letter to Editor

Volume 2 Issue 2 - 2015

Charalampos Spyropoulos*

Department of Surgery, IASO General Hospital, Greece

***Corresponding author:** Charalampos Spyropoulos, Department of Surgery, IASO General Hospital, 4 Foskolou Street, Chalandri, 15232, Athens, Greece, E-mail: charisspyropoulos@gmail.com

Received: April 14, 2015 | Published: April 28, 2015

Abstract

Selective IgM deficiency is a rare type of dysgammaglobulinemia characterized by low levels of serum IgM and repeated multisystemic infections. Yersiniosis is a food-borne disease that has become more prevalent recently due to human transmission via the fecal-oral route and its occurrence in farm animals. No clinically obvious development of Yersiniosis in patients with selective IgM deficiency has been described so far. The present report describes a patient with selective IgM deficiency who presented with repeated microbial infections and terminal ileitis due to *Yersinia* infestation.

Keywords: Terminal ileitis; *Yersinia enterocolitica*; Immunoglobulin M deficiency

Introduction

Selective IgM deficiency is a rare type of dysgammaglobulinemia characterized by low levels of serum IgM and repeated multisystemic infections. Yersiniosis is a food-borne disease that has become more prevalent recently due to human transmission via the fecal-oral route and its occurrence in farm animals. No clinically obvious development of Yersiniosis in patients with selective IgM deficiency has been described so far. The present report describes a patient with selective IgM deficiency who presented with repeated microbial infections and terminal ileitis due to *Yersinia* infestation [1]. A male patient, aged 40, was admitted to our Unit in order to further investigate a recently diagnosed terminal ileitis. His medical history started on nine months ago with 2-3 watery diarrheas daily, without blood or mucus. Furthermore, he had experienced frequent attacks of acute febrile tonsillitis and sinusitis over the last two years. A total colonoscopy with ileoscopy revealed the presence of aphthous ulcers in the mucosa of the terminal ileum. Histology indicated infiltration of the lamina propria by plasmacytes, lymphocytes and neutrophils, eroding the surface epithelium. Mild edema and hyperplasia of goblet cells were also present. Stool cultures and examination for parasites and ova were negative. Liver and renal function tests were normal.

The working diagnosis was terminal ileitis of unknown origin. An examination for possible infection with *Yersinia enterocolitica* revealed 5 IgG and 3 IgA serum YOP antigens positive (Western Blot Analysis - Athens Pasteur Institute) (Table 1). The test is considered positive if at least three IgG or two IgA bands are seen in immunoblotting at a level greater than 10% (IgG) or 5% (IgA) of reference standards [2]. Serum immunoglobulin estimation showed normal IgG and IgA levels but very low levels of IgM. Further studies concerning adaptive and innate immunity and serum auto-antibodies were normal. A head and chest CT scan was performed revealing chronic rhino-sinusitis. Based on these findings, the diagnosis of terminal ileitis due to infection with *Yersinia enterocolitica* was made. The repeated bouts of sinusitis

and tonsillitis were attributed to the patient's immunological defect caused by the coexistent selective immunoglobulin M deficiency [3]. The patient was started on ciprofloxacin per os (500 mg twice daily for three weeks) with total resolution of bowel symptoms. Nine months later, he is in perfect clinical condition. The diagnosis of selective IgM deficiency was based on the low levels of IgM whereas Yersiniosis was evident by the positive results of serum IgG and IgA antibodies against *Yersinia enterocolitica* and *Yersinia pseudotuberculosis* outer membrane proteins (Yop proteins) [4]. Thus, this patient represents the first description of *Yersinia enterocolitica* infection with terminal ileitis coexisting with selective IgM deficiency. We suggest that selective IgM deficiency could be combined with terminal ileitis due to Yersiniosis infection. A search for the presence of *Yersinia* YOP serum antibodies using Western Blot technique should be performed in selected cases.

Table 1: Serum antibodies against *Yersinia enterocolitica* and *Yersinia pseudotuberculosis* (Western Blot analysis).

Antibodies against proteins	Size of proteins (Kd)	IgG antibodies	IgA Antibodies
Yop M (2a)	46	Positive	Positive
Yop H (2b)	44	[-]	[-]
Lor V (V antigen)	38	Positive	Positive
Yop D (4a)	36	Positive	Positive
Yop N (4b)	34	Positive	[-]
Yop P (3o)	30	[-]	[-]
Yop EM (5)	25	Positive	[-]

References

1. Galindo CL, Rosenzweig JA, Kirtley ML, Chopra AK (2011) Pathogenesis of *Y. enterocolitica* and *Y. pseudotuberculosis* in human Yersiniosis. *J Pathog* 2011: 182051.
2. Gupta S, Louis AG (2013) Tolerance and autoimmunity in primary immunodeficiency disease: a comprehensive review. *Clin Rev Allergy Immunol* 45(2): 162-169.
3. Goldstein MF, Goldstein AL, Dunsky EH, Dvorin DJ, Belecanech GA, et al. (2006) Selective IgM immunodeficiency: retrospective analysis of 36 adult patients with review of the literature. *Ann Allergy Asthma Immunol* 97(6): 717-730.
4. Morita M, Saibara T, Nakazawa Y, Miyao M, Okazaki K, et al. (1984) A case of Crohn's disease associated with selective IgM deficiency. *Nihon Shokakibyō Gakkai Zasshi* 81(4): 1070-1075.