

# HCC Burden in Egypt

**Mini Review**

Volume 2 Issue 3 - 2015

**Ehab Abdel-Atti\***Hepatology and Gastroenterology Unit Internal Medicine  
Department, Menoufia University, Egypt**\*Corresponding author:** Ehab Abdel-Atti, Hepatology  
and Gastroenterology Unit Internal Medicine Department,  
Faculty of Medicine, Menoufia University, Shabin Elkom,  
Egypt, Email: ehab\_abdelatty@hotmail.com**Received:** May 13, 2015 | **Published:** July 09, 2015**Abstract**

Hepatocellular carcinoma (HCC) is the commonest primary cancer of the liver. Geographical distribution of HCC varies throughout the world with an incidence rate ranging from 2.1 in Central America to 35.5 in Eastern Asia. The burden of HCC has been increasing in Egypt with a doubling in the incidence rate in the past 10 years. This may be attributed to high prevalence of HCV in Egypt. Egypt has a high incidence of HCC about 21% in cirrhotic Egyptian patients. HCV and HBV infections, diabetes and smoking are the main determinants of HCC development in Egypt. Occult HBV infection may influence the outcome of HCV infection leading to development of HCC. An active surveillance and secondary prevention programs for patients with chronic hepatitis are the most important steps to reduce the risk of HCC.

**Abbreviations:** HCC: Hepatocellular Carcinoma; HCV: Hepatitis C Virus; HBV: Hepatitis B Virus

**Introduction**

Hepatocellular carcinoma (HCC) is one of the most common cancers worldwide. It is rarely detected early. HCC is the commonest primary cancer of the liver. Hepatocellular carcinoma is one of the 130 major causes of morbidity and mortality in the world. It represents the third leading cause of cancer death in males and the fourth in females with more than 600000 deaths per year [1]. Geographical distribution of HCC varies throughout the world with an incidence rate ranging from 2.1 in Central America to 35.5 in Eastern Asia. The burden of HCC has been increasing in Egypt with a doubling in the incidence rate in the past 10 years [2]. There is a geographic correlation between the incidence of HCC and the prevalence of chronic hepatitis B and C, suggesting that these two viral infections are the most important risk factors of HCC worldwide. Co-infection with *Hepatitis B virus* (HBV) and *Hepatitis C virus* (HCV) is associated with a higher risk for developing HCC than either infection alone [3]. Several risk factors for the development of HCV-associated HCC have been reported such as aging, gender and alcohol intake [4].

Egypt has the highest prevalence of HCV worldwide and has rising rates of HCC [5-7]. Our retrospective study about risk factors for hepatocellular carcinoma on cirrhotic Egyptian patients [8] was done on 300 patients with HCC and 50 patients with chronic liver diseases without HCC as controls. Prevalence rate of DM and smoking was significantly higher in HCC cases (59.3% and 69% respectively) than controls (38% and 50% respectively) (P=0.005 and 0.006 respectively). The prevalence of HBsAg and HCV Ab was significantly higher in HCC cases (18% and 70% respectively) than controls (4% and 40% respectively) (P=0.02 and 0.0001 respectively). On multivariate analysis, the risk of HCC development in smokers with HBV or HCV was 4.90 and 8.47 respectively (OR) (P=0.0001). It was higher than in non-smokers with HBV or HCV (OR=2.48 and 4.44 respectively) (P=0.037 and 0.0001 respectively) and in smokers without HBV or HCV (OR=2.56 and 2.77 respectively) (P=0.01). The risk of HCC development in HBV or HCV positive patients

with DM was 3.98 and 9.19 respectively (OR) (P=0.001 and 0.0001 respectively). It was higher than for HBV or HCV positive patients without DM (OR=2.80 and 4.65 respectively) (P=0.031 and 0.0001 respectively) and that for HBV or HCV negative patients with DM (OR=2.56 and 2.23 respectively) (P=0.011 and 0.0001 respectively). A preliminary data from a survey on 1350 Egyptian cirrhotic patients revealed that 289 patients had HCC (21.4%). Also HCV, HBV and smoking are risk factors of HCC development.

Our study about the prevalence of occult HBV (OBI) among HCV positive Egyptian patients documented that out of 100 chronic HCV patients; only 16 patients (16%) had OBI. HCC was significantly more common in OBI/HCV dually infected (31%) than HCV monoinfected patients (7%) (P=0.01). On doing multiple logistic regression analysis, OBI is a risk factor for HCC in Egyptian HCV positive patients (P=0.04) [9]. Extensive collaborative research was carried out during the last decade to explore independent and combined effects of HBV and HCV and other factors in the etiology of HCC. Although HBV is considered worldwide as a major risk factor for liver cirrhosis and HCC, the prevalence of HBV infection in Egypt has been declining over the last two decades, while HCV has increased. Egypt has possibly the highest HCV prevalence worldwide around 14% [10]. In our previous study [8], HCV positive patients are 211 representing 70% of the total number of HCC cases, while HBV positive cases are only 54 representing 18%. There is a higher OR for HCV +ve cases compared with those of HBV representing 24.33 and 17.65 respectively. The rising trend of HCC has been associated with increased prevalence of *Hepatitis C virus* (HCV) infection in Egypt [8]. Moreover, in addition to HCV and HBV infections, DM appears to be associated with HCC [11]. DM increased the risk of HCC development independent of chronic HCV hepatitis and chronic HBV hepatitis as previously described [5,12]. Previous studies suggest that HCV, HBV and DM play an independent role and synergistic effect in liver tumorigenesis [13,14]. HCV positive Egyptian cirrhotic patients with diabetes had a higher risk of HCC than those without diabetes. Seronegative individuals with diabetes also had a higher risk of HCC than those without diabetes [8].

A prominent role of tobacco smoking in the etiology of HCC can be observed. Such a relationship is due to several chemicals in tobacco smoke metabolized and then activated to be carcinogenic in the liver [15]. Smoking increased the risk of HCC development independent of HCV and HBV infections in Egyptian cirrhotic patients [8]. There was a synergism between HBV and smoking as risk factors for HCC in Egyptian cirrhotic patients. Moreover, smokers with HBV +ve have an OR significantly higher than HBV +ve alone or smoking alone (OR= 4.90, 2.84 and 2.56 respectively). Franceschi et al. [16] had demonstrated that tobacco smoking was related to HCC risk and seemed to enhance HCC risk among viral hepatitis patients.

## Conclusion

Egypt has a high incidence of HCC about 21% in cirrhotic Egyptian patients. HCV and HBV infections, diabetes and smoking are the main determinants of HCC development in Egypt. There is a synergistic effect of many risk factors. An active surveillance and secondary prevention programs for patients with chronic hepatitis are the most important steps to reduce the risk of HCC. Occult HBV infection may influence the outcome of HCV infection leading to development of HCC. The persistent HBV infection may have a critical role in the development of HCC in HBsAg-negative patients. So, occult HBV should be considered and evaluated by more sensitive PCR among HCV-infected patients.

## References

- Hassan MM, Spitz MR, Thomas MB, Curley SA, Patt YZ, et al. (2009) The association of family history of liver cancer with hepatocellular carcinoma: A case control study in the United States. *J Hepatol* 50(2): 334-341.
- Gomaa AI, Khan SA, Toledano MB, Waked I, Taylor-Robinson SD (2008) Hepatocellular carcinoma: Epidemiology, risk factors and pathogenesis. *World J Gastroenterol* 14(27): 4300-4308.
- Shi J, Zhu L, Liu S, Xie W-F (2005) A meta-analysis of case-control studies on the combined effect of hepatitis B and C virus infections in causing hepatocellular carcinoma in China. *Br J Cancer* 92(3): 607-612.
- Ohata K, Hamasaki K, Toriyama K, Matsumoto K, Saeki A, et al. (2004) Hepatic Steatosis Is a Risk Factor for Hepatocellular Carcinoma in Patients with Chronic Hepatitis C Virus Infection. *Cancer* 97(12): 3036-3043.
- Ismail AM, Ziada HN, Sheashaa HA, Shehab El-Din AB (2009) Decline of viral hepatitis prevalence among asymptomatic Egyptian blood donors: a glimmer of hope. *Eur J Intern Med* 20(5): 490-493.
- Lehman EM, Wilson ML (2009) Epidemiology of hepatitis viruses among hepatocellular carcinoma cases and healthy people in Egypt: a systematic review and meta-analysis. *Int J Cancer* 124(3): 690-697.
- El-Zayadi A, Badran HM, Barakat EM, Attia ME, Shawky S, et al. (2005) Hepatocellular carcinoma in Egypt: A single center study over a decade. *World J Gastroenterol* 11(33): 5193-5198.
- Baghdady I, El-Kaffrawy N, Abd El-Atti E, Abd El-Bary N, Saber M (2013) Study of the risk factors for hepatocellular carcinoma: effect of their synergism. *Journal of American Science* 9(4): 211-217.
- Baghdady I, Shoeib S, Abdelatti E, Dala A, Eed E, et al. (2011) Occult Hepatitis B Virus infection in Egyptian Hepatitis C Virus positive patients: Prevalence and impact on Hepatocellular Carcinoma development. *MMJ* 24(1): 197-210.
- Anwar WA, Khaled HM, Amra HA, El-Nezami H, Loffredo CA (2008) Changing pattern of hepatocellular carcinoma (HCC) and its risk factors in Egypt: Possibilities for prevention. *Mutat Res* 659(1-2): 176-184.
- Moore MA, Park CB, Tsuda H (1998) Implications of the hyperinsulinaemia diabetes-cancer link for preventive efforts. *Eur J Cancer Prev* 7(2): 89-107.
- Donadon V, Balbi M, Gherseti M, Grazioli S, Perciaccante A, et al. (2009) Antidiabetic therapy and increased risk of hepatocellular carcinoma in chronic liver disease. *World J Gastroenterol* 15(20): 2506-2511.
- Hassan MM, Hwang LY, Hatten CJ, Swaim M, Li D, et al. (2002) Risk factors for hepatocellular carcinoma: synergism of alcohol with viral hepatitis and diabetes mellitus. *Hepatology* 36(5): 1206-1213.
- Mori M, Hara M, Wada I, Hara T, Yamamoto K, et al. (2000) Prospective study of hepatitis B and C viral infections, cigarette smoking, alcohol consumption, and other factors associated with hepatocellular carcinoma risk in Japan. *Am J Epidemiol* 151(2): 131-139.
- Lee YC, Cohet C, Yang Y-Ch, Stayner L, Hashibe M, et al. (2009) Meta-analysis of epidemiologic studies on cigarette smoking and liver cancer. *Int J Epidemiol* 38(6): 1497-1511.
- Franceschi S, Montella M, Polesel J, Vecchia CL, Crispo A, et al. (2006) Hepatitis Viruses, Alcohol, and Tobacco in the Etiology of Hepatocellular Carcinoma in Italy. *Cancer Epidemiol Biomarkers Prev* 15(4): 683-689.