

# Hyponatremic Seizure Associated with Concomitant Use of Standard Dose Trimethoprim-Sulfamethoxazole with a Diuretic

## Proceeding

Trimethoprim-Sulfamethoxazole (TMP-SMX) is an antimicrobial agent used to treat infectious disease pathogens [1]. Electrolyte abnormalities, primarily hyperkalemia and hyponatremia, are rare adverse effects reported with higher doses of trimethoprim use [2]. Paucity of data exists for similar effects in patients using standard doses of TMP-SMX [3], with no previous reported case of trimethoprim induced hyponatremic seizure in the literature. We illustrate a case of severe hyponatremia and hyponatremic seizure in a patient on regular dose of TMP-SMX who was concomitantly using a diuretic.

A 64-year old hypertensive female taking Lisinopril-HCTZ (20-12.5mg) combination pill was started on TMP-SMX (160-800 mg) twice daily for left fourth toe blister. She presented to the emergency room four days later with fatigue, confusion, and a witnessed grandmal seizure. Physical examination, electrocardiogram, urinalysis, chest X-ray, and cranial computed tomography were unremarkable. Serum Na was measured at 119mmol/L (136-145mmol/L) compared to 131mmol/L last week prior to antibiotic use. Other labs included TSH 0.94U/L, glucose 109 mg/dl, K 3.9 (3.5-5.1mmol/L), SOsm 245mOsm/kg (280-303mOsm/Kg), BUN 10mg/dL (7-18mg/dl), SCr 0.9mg/dL (0.4-0.9mg/dL), BUN: SCr 11.1 (10.0-20.0) and creatinine clearance 52.2 mL/min using Cockcroft-Gault equation. Urinary indices showed UOsm 205mOsm/kg (50-800 mOsm/kg), UNa 22mmol/L, UCr 24.5mg/dL and FENa 0.7%. TMP-SMX and Lisinopril-HCTZ were discontinued. The patient received hypertonic saline. Serum Na gradually improved to 131mmol/L over 24 hours and the patient's confusion resolved without further seizures.

Trimethoprim, a heterocyclic weak base, structurally mimics potassium-sparing diuretics, and at higher doses causes natriuresis by blocking sodium reabsorption in the distal nephron [4]. Hyponatremia, while usually asymptomatic, can exert

neurological sequelae. In our case, standard dose TMP-SMX lead to salt wasting and severe hyponatremia which manifested as seizure; its dose-dependent effect was perhaps potentiated by the concomitant use of a diuretic which can induce renal dysfunction and affect sodium and water hemostasis.

## References

1. Khow KS, Yong TY (2014) Hyponatraemia associated with trimethoprim use. *Curr Drug Saf* 9(1): 79-82.
2. Babayev R, Terner S, Chandra S, Radhakrishnan J, Mohan S (2013) Trimethoprim-associated hyponatremia. *Am J Kidney Dis* 62(6): 1188-1192.
3. Mori H, Kuroda Y, Imamura S, Toyoda A, Yoshida I, et al. (2003) Hyponatremia and/or hyperkalemia in patients treated with the standard dose of trimethoprim-sulfamethoxazole. *Intern Med* 42(8): 665-669.
4. Choi MJ, Fernandez PC, Patnaik A, Coupaye-Gerard B, D'Andrea D, et al. (1993) Brief report: trimethoprim-induced hyperkalemia in a patient with AIDS. *N Engl J Med* 328(10): 703-706.

Proceeding

Volume 4 Issue 2 - 2016

**Lynda Hoang\*, Rabaiya Ali, Jose Manriquez and Sulaiman Sultan**

*Internal Medicine Residency, Parkview Medical Center, USA*

**\*Corresponding author:** Lynda Hoang, Internal Medicine Residency, Parkview Medical Center, USA, Email: lynda.hoang@tu.edu

**Received:** January 26, 2016 | **Published:** February 12, 2016