

At The Focal Point-Impacted Foreign Body in Sigmoid Colon Diverticula

Opinion

An 80 years old lady, was scheduled for follow up colonoscopy, due to previous advanced polyps removal and relative sigmoid colon stenosis secondary to severe diverticular disease. She was asymptomatic, without any pain or fever [1,2].

At colonoscopy, an impacted foreign body (Toothpick) was found, impacted in two sigmoid colon ("face to face") diverticulum, one at each end. There was purulent secretion from one of those ends. It was not easy to remove, and mobilization resulted in more purulent exudate. The perforation hazard was considered. It couldn't be cut in its central portion and finally with easy movements with a grasping forceps, it was dislodged from one of its edges, and removed with a snare. The risky meal, assumed to be responsible for the ingestion of the foreign body, was 6 days prior to the colonoscopy [3-10].

Opinion

Volume 5 Issue 3 - 2016

Valle Arturo, Navarrete Claudio and Sáenz Roque*

German School of Clinical Medicine, University Development, Chile

***Corresponding author:** Sáenz Roque, The Latin American GIE Training Center, German School of Clinical Medicine, University Development, Santiago, Chile, Email: rsaenz@alemana.cl

Received: August 27, 2016 | **Published:** September 13, 2016

An abdominal ultrasound failed to show any collection. Patient had no further symptoms, while 7 follow up days, under antibiotic treatment [11] (Figure 1).




Figure 1: Sigmoid colon diverticulum. Impacted foreign body and purulent secretion at the edges.

References

1. Schwartz GF, Polsky HS (1976) Ingested foreign bodies of the gastrointestinal tract. *Am Surg* 42(4): 236-238.
2. Eldridge WW (1961) Foreign bodies in the gastrointestinal tract. *JAMA* 178: 665-667.
3. Fidler M (1972) Foreign-body perforation of a jejunal diverticulum. *Br J Surg* 59(9): 744-745.
4. Ashby BS, Hunter-Craig ID (1967) Foreign-body perforations of the gut. *Br J Surg* 54(5): 382-384.
5. Stack LB, Munter DW (1996) Foreign bodies in the gastrointestinal tract. *Emerg Med Clin North Am* 14(3): 493-521.
6. Selivanov V, Sheldon GF, Cello JP, Crass RA (1984) Management of foreign body ingestion. *Ann Surg* 199(2): 187-191.
7. Wilcher GW (2010) Toothpick perforation of a colon diverticulum: an adjunct autopsy finding. *Med Sci Law* 50(3): 156-158.
8. Chung YS, Chung YW, Moon SY, Yoon SM, Kim MJ, et al. (2008) Toothpick impaction with sigmoid colon pseudodiverticulum formation successfully treated with colonoscopy. *World J Gastroenterol* 14(6): 948-950.

9. Grabysa R, Łowczak R, Kubiak M, Kowalczyk M, Zaborowski P (2010) Toothpick impaction in the sigmoid colon as a cause of chronic abdominal pain. *Endoscopy* 42(Suppl 2): E203.
10. Mohr HH, Dierkes-Globisch A (2001) Endoscopic removal of a perforating toothpick. *Endoscopy* 33(3): 295.
11. Reddy SK, Griffith GS, Goldstein JA, Stollman NH (1999) Toothpick impaction with localized sigmoid perforation: successful colonoscopic management. *Gastrointest Endosc.* 50(5): 708-709.