

A review: micropropagation of guava (*Psidium Spp*)

Abstract

The rapid process for development of identical offspring is micropropagation. In this technique the plants grow from meristematic tissue or somatic cells of superior plants using suitable nutrient media under controlled environmental conditions. Micropropagation in Guava, was successfully done on the variety Allahabad safeda. After that micropropagation progress has been made in the different areas via *in vitro* culture of guava. MSO or MS0 (MS-zero) was mostly used as mineral medium for culturing *Psidium species*. The Woody Plant Medium (WPM) was more suitable for micro propagation of some guava genotypes. Starting material for guava culture (in vitro) is growing shoot tips or nodal segments as explants (1.0-3.0 cm size) collected during in spring. Browning of the explant and culture medium is mostly obstruct in culturing of woody plants, which is remove by avoidance of toxic substances in the medium in guava. Cytokinin (6-Benzylaminopurine) is the most frequently used for guava micropropagation. Use of dual auxins (IBA & NAA) in compare to IBA followed by NAA, resulted higher frequency of rooting, which is important for successful survival of plant. In guava, acclimatization has been experienced by gradually decreasing the relative humidity.

Keywords: guava, micropropagation, BAP, IBA, NAA

Volume 2 Issue 6 - 2018

Krishan Kumar Singh,¹ Shiv Pratap Singh²

¹Department of Horticulture, Chauras Campus, HNB Garhwal University, India

²Department of Botany, Hindu college, India

Correspondence: Krishan Kumar Singh, Department of Horticulture, Chauras Campus, HNB Garhwal University, Srinagar (Garhwal) 246174, Uttarakhand, India, Email forekrishna@gmail.com

Received: June 14, 2018 | **Published:** December 26, 2018

Introduction

Guava (*Psidium guajava* L.) belongs to Myrtaceae family. Place of origin of guava is tropical^{1,2} America. It is utilized as fresh fruit consumption in India along with other Asian sub-continental countries. Genus *Psidium* have about 150 species, most of which are fruit bearing trees. It is native to tropical America, stretching from Mexico to Peru.¹ It is widely adopted and tolerate in frost, drought and saline conditions.³ It has good amount of ascorbic acid, dietary fibres, vitamin-A (about 250 IU/100 g), pectin, calcium, phosphorus, iron and other nutrients. The roots, bark, leaves and green fruits are used as medicine for gastrointestinal problems, diarrhoea and dysentery.⁴ Salad, guava jelly and pudding are made from the of ripe fruits. Besides that, Guava juice and guava pulp wines are also prepared from guava fruit.⁵ It also contains high amount of antioxidants like, lycopene, carotenoids and polyphenols.⁶ Guava has the chemicals which help to reduce the incidence of diseases like arthritis, arteriosclerosis, diabetes, cancer, heart disease, inflammation and brain dysfunction. Antioxidants reported to retard aging, high concentrations of pectin content in guava fruit reduces the cholesterol of human body.⁷⁻⁹ Production of Guava is overlaying the agronomical and horticultural problems including susceptibility to many pathogens, wilting, low fruit growth, short life, higher seed content and stress sensitivity. Improving guava species by traditional methods of breeding are limited because plants generally have long juvenile growth periods, and are heterozygous in nature. While, seed originated guava plantlets do not maintain the genetic purity of the variety due to the segregation and mixing of genetic characters during sexual reproduction. In addition, high internal fungi, bacterial contamination and phenolics compounds exudation tend to limit *in vitro* cultures of the guava plant. Genetic engineering technique reducing the breeding period. Due to this approach micro-propagation and regeneration produce large numbers of rooted plants from unique plants. Clonal propagation can reduce plant-to-plant variation and ensuring the uniform populations of clones.

Micropropagation

Micropropagation refers to *in vitro* plant propagation method. The advantage of micropropagation over conventional propagation technique is to reduce the time needed for achieving large scale propagation of plants which are true to the type and disease free. Murashige¹⁰ and Hammerschlag¹¹ reported three stages i.e. establishment, multiplication and rooting. Hartmann and Kester (1983) added another stage i.e. acclimatization. Micropropagation is now a well-established technology which has made significant contributions to the propagation and improvement of agricultural crops in general and applicable to various crops like *Withania somnifera* L. (Ashwagandha)¹² *Saccharum officinarum* L. (Sugercane).¹³ Greater contribution is predicted from this technology in years to come, both in its own right and as an adjunct to the application of molecular biology.

Explants

Totipotency is the ability of plant tissues/parts (excepting bark) to regeneration *in vitro*. Juvenile explants, comparatively gives better results. New vegetative growths in *Psidium guajava* L. have been reported to be reliable source of explant.^{14,15} The majority of workers have used actively growing shoot tips or nodal segments as explant.¹⁶⁻²⁵ A propagation culture system from germinated seedlings has been established by Yasseen et al,²⁶ and Shah et al.²⁷

Time of collection and size of explants

The explants of guava collected from the base of the main stem when vegetative growth vigorously establish.¹⁴ Parkash and Tiwari,¹⁸ reported that early spring collection of explants show less contamination in compare to late autumn and summer. Some reports by other workers,^{20,21,28,29} on explants collection in spring gives best culture establishment and profuse sprouting. Generally, the size of explants best suited for *in vitro* propagation of guava is 1.0-3.0 cm (Table 1).^{14,16,22,24,30-33}

Table 1 Sterilization of explant

S. No.	Chemical used for sterilization	Reference
1.	Rinse in 70 % ethanol+HgCl ₂ 0.05-0.50 % (containing a few drops of Tween-20) for 2-5 min.	Amin &Jaiswal. ³⁰
2.	70 % ethanol for 1 min+5 % NaOCl for 5 min and treatment of shoots which were plastic wrapped with 0.05 per cent HgCl ₂ for 5 min.	Khattak et al. ³⁷
3.	Wash apical shoots of 5-10 cm for 30-40 min in running tap water + washed with 1 % KCl+1 % NaOCl for 5-7 min.+ washed 3-4 times with sterile distilled water	Parkash & Tiwari ¹⁸
4.	Shoot tip or nodal cuttings (3-5 cm) transferred in a solution 2 g l-l bavistin, 50 mg l-l ascorbic acid, 75 mg l-l citric-acid and 1 ml l-l liauiclean (a.i. benzolkonium chloride); cut 1-2 cm long nodal sections washed thoroughly under running tap water for 15 min.; rinsed with 0.4 % HgCl ₂ for 10 min.	Siddiqui & Farrooq. ³⁵
5.	Sterilization with H ₂ O ₂ (10%), silver nitrate (0.25%) and HgCl ₂ (0.05%) one by one for 5, 6 and 3 min. respectively	Meghwal et al. ²⁰ Bisen& Tiwari, ²¹
6.	Stem segments in 200 mg l-l 8-hydroxy quinaline citrate + 0.1 per cent bavistin; washed in running tap water for 30 min.; stem parts were surface sterilized by dipping them serially in multi disinfectant viz., 70 % ethanol (15 sec.), 10 % (v/v) H ₂ O ₂ (3 min) and 0.05 % HgCl ₂ (3 min)	Singh et al. ¹⁹
7.	Ethyl alcohol (70%) for 30 sec, HgCl ₂ (0.1%) for 5 min and NaOCl(1%) for 6 min	Kumar & Tiwari ³⁹
8.	H ₂ O ₂ (10%, v/v), silver nitrate (0.25%) and HgCl ₂ (0.05%) for 5, 5 and 3 minutes, respectively; 2-3 times rinses with sterile distilled water	Meghwal et al. ²⁹
9.	70% ethanol; 0.05 % HgCl ₂ and a drop of surfactant (Tween 80) was added and agitated at 100 rpm on a rotary shaker for 5 min.; shoot tips were rinsed 3 times with sterile distilled water	Zamir et al. ²²
10.	15% bleach solution for 20 min; culturing in MS medium with 250 mg L-l polyvinyl-pyrrolidone	Xiaomei et al. ²⁴

Phenolics

Guava exude the phenolic compounds into the culture media makes difficulty in regeneration process. *Psidium guajava* L. is a recalcitrant species having high phenolic exudation that kills explants from sources outside the laboratory. Establishment of *in vitro* cultures of woody plants is greatly hampered by the browning of the explant

and culture medium. Browning of explant is due to oxidation of the phenolic compounds, released from the cut ends of the explants. *In vitro* establishment of the guava explants was very difficult due to the exudation of phenolic compounds into cultures, by which the media turned brown or black within 12-24 hrs and most of the explants died within 2 days of inoculation (Table 2).^{15,20,22,24,30,31,32,34-36}

Table 2 Control of phenolics

S. No	Treatment/strategies to remove toxics from media	Reference
1.	Early spring growth which gave less contamination and browning	Kumar, ¹⁵ Singh et al, ¹⁹ Meghal et al. ²⁹
2.	Partial etiolation of stock plants of Allahabad Safeda and aneuploid No. 82 (PusaSrijan) rootstock before explant excision followed by culturing the explants in darkness resulted in early bud sprouting and significant increase in explant survival	Leon-de-Sierralta. ³⁶ Kumar. ¹⁵ Meghwal et al. ²⁹ Bisen ²¹ , Joshee et al. ⁵³
3.	Transferring the explants in fresh medium at short intervals	Amin & Jaiswal, ¹⁴ Singh & Tiwari, ⁵⁸ Siddiquai & Farooq, ³⁵ Kumar & Tiwari ²⁸
4.	Dipping the explants in the antioxidant solution i.e. citric acid (75 mg/l) and ascorbic acid (50 mg/l)	Fitchet, ³⁴ Singh et al. ¹⁹ Kumar, ¹⁵ Meghwal et al. ²⁰ Josheet al. ⁵³
5.	Use of absorbing agents such as activated charcoal, polyvinylpyrrolidone (PVP)	Amin &Jaiswal, ¹⁴ Mohammed et al. ²⁶ Fougat et al. ⁵¹ Siddiqui & Farooq, ³¹ Kumar & Tiwari, ²⁸ Meghwal et al. ²⁰ Concepción et al. ⁴⁹ Xiaomei & Yang, ²⁴
6.	Agitation of explants in antioxidant solution	Meghwal et al. ³² Meghwalet al. ²⁰ Zamir et al. ²²
7.	Air drying of explants in an air flow for 30-45 min prior to inoculation	Fitchet ³⁴
8.	Coating the explants at their cut ends with commercial silicon which is a novel process was established for completely inhibiting phenol-based browning	Youssef et al. (2010)
9.	Regeneration of explants has been successfully achieved mainly in Murashige and Skoog's medium	Amin & Jaiswal, ¹⁴ Fitchet, ³⁴ Loh& Rao; ¹⁶ Khattak et al. ³⁷ Mohammed et al. ²⁶ Parkash & Tiwari ¹⁸ Ramirez & Salazar; ³⁸ Siddiqui & Farrooq ³¹ Meghwal et al. ²⁰ Bisen, ³³ Zamir et al. ²² Shah et al. ²⁷ Xiaomei & Yang, ²⁴ Usman et al. ²⁵

Plant growth regulators

Plant Growth regulator are such organic compounds occurring naturally in the plants as well as synthetic and promote, inhibit or modify any physiological process in small amount. The concentration of plant growth regulator varied from species to species for *in vitro* culture of guava, and type of growth to be initiated i.e. callus formation, shoot proliferation, rooting, etc. In order to support good growth of tissue and organs, it is generally required to add one or more PGR's such as auxin, cytokinin and gibberellins in the medium. Cytokinin levels especially have been shown to be critical for multiplication of many tropical fruit trees. BA has been the most common cytokinin used for guava propagation. In guava, media supplemented with 4.5 mM BA produced 3-6 shoots in 12 weeks¹⁴ while, according to other workers^{16,17} 0.5 to 2.0 mg/l BA produced maximum shoot proliferation. Maximum number of shoots in guava cultivar sardar (L-49) was obtained in media supplemented with 1.0 mg/l BA+0.2 mg/l IBA. Siddiqui and Farooq³² found that 1.0 mg/l BA was effective in stimulating the formation of axillary shoots and similar results were also reported by Fitchet,³⁴ Meghwal et al.³² Khattak et al.³⁷ and Zamir et al.²² who got maximum shoot development in cv. Safeda

with 1mg/l BAP and glutamine 500 mg/l cultured on MS medium. Ramirez and Salazar³⁸ obtained highest percentage of bud sprouting in MS medium supplemented with 4 mg/l BA in nodal segment of guava. A combination of 2 mg/l BA and 0.2 mg/l IBA was found more effective for sprouting and number of shoots per explant at establishment stage. Further increase in cytokinin reduced the explant sprouting in Chinese guava.³⁹ Highest number of shoots proliferated on WPM supplemented with 0.5 to 1.0 mg/l BAP in different cultures of guava.^{19,20} However, Usman et al.²⁵ reported that maximum shoots were induced with 2.0 mg/l of BAP.

Root induction

Media

There is no single medium as well as constituent of medium to suit every stage of *in vitro* cloning. Therefore, to attain full plants, the micro-shoots must be transferred to a rooting medium which is different from the shoot proliferation medium especially in its hormonal composition.⁴⁰ Full strength WPM was used for rooting of guava cultivars Allahabad Safeda, Lucknow-49 and Thailand and Aneuploid No. 82 (Table 3).²⁰

Table 3 Plant growth regulators

S.No.	Treatment of chemicals for root induction	References
1.	Half MS medium supplemented with 1 mm each of IBA and NAA	Amin & Jaiswal, ¹⁴ Yasseen et al. ²⁶ Shah et al. ²⁷
2.	Shoots produced from guava explants produced roots in media containing 9.8 mM TBA	Mohammed et al. ²⁶
3.	Shootlets cultured on half MS 0.2 mg/l IBA and incubated for one week in dark resulted in early and better rooting of guava	Parkash & Tiwari ¹⁸
4.	Shoot explants were easily rooted in vitro using Rugini olive medium (OM) with 0.5 or 1.0 mg NAA or IBA/l	Papadatau et al. ¹⁷
5.	Best rooting (66.66%) was recorded with IBA+NAA (0.2+0.2 mg/ l) in Chinese guava	Kumar ¹⁵
6.	Achieved best rooting of micro shoots on half strength modified Murashige and Skoog's (MMS) medium supplemented with 4.90 mM indole-3-butyric acid along with 100 mg l-l activated charcoal	Singh et al. ¹⁹
7.	Excised proliferating shoot segments of 1.5 to 2.0 cm for in vitro rooting on WPM containing 200 mg/l activated charcoal and dual auxins (IBA and NAA) @ 0.2 mg/l each which resulted in higher frequency of rooting	Meghwal et al. ²⁹
8.	Obtained maximum (54) plants rooted with average number of roots (3.8) per plantlet in MS medium supplemented with 2.5 mg/L IAA + 2.5 mg/L IBA	Zamir et al. ²²
9.	Achieved maximum rooting (65%) when shoots were dipped in 4.9 mM Indole-3-butyric acid (IBA) solution for 1 min and then rooted	Xiaomei & Yang ²⁴

Rooting

Among the various auxins viz., IBA, IAA, NAA etc. used for *in vitro* rooting, IBA is the most commonly used auxin. Khattak et al.³⁷ reported that the initiation of adventitious roots at 1/2MS+1 mg/l IBA from *in vitro* derived guava shoots, whereas 2.5 mg/l each of IAA and IBA for root development was noticed by Raziuddin et al.⁴¹ The best rooting of *in vitro* desired microcutting of guava cv. Hisar Safeda with 1/2MS+2 mg/l IBA+200 mg/l activated charcoal.⁴² Biswas et al.⁴³ reported healthy root system on half strength WPM media containing 0.1 mg/l NAA. Shah et al (2008) observed NAA supplementation at 0.5 mg/l in 1/2MS + IBA media was found most responsive for cv. Safeda. Singh et al.²⁸ reported the *in vitro* clonal propagation of guava cultivars Allahabad Safeda, Lucknow -49 and Thailand and concluded that the rooting was noted to be the best in WPM + 0.5 or 1.0 mg/l IBA and NAA + 200 mg/l activated charcoal.

Hardening of rooted plantlets

The success of any micropropagation research depends on the success of plantlet transferring technique, where shoot or plantlets that have been growing heterotrophically under an aseptic environment of test tube (having very high humidity) become autotrophic and grows under condition of moderate to low humidity. For acclimatization rooted plantlets of guava were taken out of culture tubes, washed thoroughly to remove any remaining medium and planted in small plastic pots filled with garden soil and compost (1:1). During first 7-10 days, the potted plantlets were covered with glass beakers to provide high humidity.³⁰ Loh and Rao¹⁶ transferred the rooted plantlets to small plastic pots (containing vermiculite) and covered initially with thin plastic film to maintain high humidity and then small holes were made in the plastic to acclimatize it gradually. The plantlets were kept outdoor under 80 per cent shade for about a week, after which

they were transplanted in pots with soil. More than 90 per cent of the plantlets survived after transplantation to soil. According to Khattak et al.³⁷ and Zamir et al.²² the rooted plantlets were transferred to potting medium (sand, clay, compost at 1:1:1 by volume) with a layer of sand in greenhouse and initially watered with half Knop's solution. Parkash and Tiwari¹⁸ transferred rooted plantlets for hardening treatments 30 days after rooting and 86 per cent survival was recorded in the pots containing a mixture of sand, soil and FYM (1:1:1). Same technique was followed by Kumar¹⁵ for Chinese guava. Whereas, Singh et al.¹⁹ washed the rooted plantlets and removed the adhering agar in sterile water and transplanted in container filled with peat + perlite (1:1) and moisture with one-fourth strength MMS (Modified Murashige and Skoog) macro-salts solution (pH 5.8). The hardened plantlets (30-45 days) were then transferred to small plastic pots filled with soil: farmyard manure, sand (1:1:1) and shifted to the glasshouse. The plants were regularly misted with one tenth MMS macro-salts + 0.1 per cent bavistin with pH adjusted to 5.7 and exposed to light of high intensity ($62 \text{ m mol m}^{-2} \text{ s}^{-1}$) with temperature maintained at $26 \pm 1^\circ\text{C}$. The plantlets (after 4 weeks) were then shifted to the shade house in small plastic pots. A regular irrigation to these plantlets were practiced at 15 days' interval and later on shifted to natural environment. Xiaomei and Yang²⁴ planted rooted shoots (with 3-5 fully expanded leaves) in 15cm diameter plastic pots containing a mixture of sterile sand and garden soil (1:3), covered with polyethylene bags for 21 days to prevent excessive water loss. The pots were watered once a week. Full strength MS macronutrient solution 10 mL was applied every other week. Plantlets were kept at 25°C in artificial light (16 h photo period and irradiance of $50 \text{ imol mm}^{-2} \text{ s}^{-1}$) provided by white florescent tubes for 6 weeks and were then transferred to the temperature controlled ($25.6/18.3^\circ\text{C}$, day/night) greenhouse to grow under natural light.

Genetic fidelity

The commercial multiplication of a large number of diverse plants species represents one of the major success stories of utilizing tissue culture technology profitably. However, a major problem often encountered with the use of tissue culture techniques such as SE is the occurrence of soma clonal variation, which is often heritable as it represents induced genetic changes.^{44,45} Thus, genetic fidelity testing is an important prerequisite for *in vitro* regeneration protocols of many crop species, particularly if the resultant plants are to be transplanted to the field. Several strategies have been employed to assess the genetic fidelity of regenerated plants, each with their own advantages and disadvantages. Molecular markers facilitate the screening of SE regenerated plants with high precision, and since these markers are unaffected by environmental factors (that can alter phenotypes), they produce reliable and reproducible results. However, for an effective analysis of the genetic stability of *in vitro* regenerated plantlets, a combination of markers that amplify different regions of the genome should be used.^{46,47} Kamle et al.⁴⁸ have reported various DNA based molecular markers for assessment of genetic fidelity in guava. Liu and Yang⁴⁷ have reported ISSR markers for assessment of genetic fidelity in guava.

Conclusion

In vitro clonal propagation protocols for different guava spp. and varieties have been developed in all over the world. For extending the guava production in tropical and subtropical areas a rapid and efficient method for clonal propagation of elite mature genotype is necessary. During past few decades, transpiring biological techniques for micropropagation and tissue culture of predominant guava cultivars

have been discussed by several researchers are effective. However, there are several problems related with *in vitro* cultures of these explants including browning or blackening of culture medium due to leaching of phenolics, microbial contamination, and *in vitro* tissue recalcitrance which require to be given more attention. Understanding of the biological processes that permit the manipulation of *in vitro* morphogenesis and investigations on various physiological, biochemical and molecular aspects of plant hormones will highly improve our understanding and gives details that will necessary for addressing the problems of *in vitro* recalcitrance or *in vitro* plant growth and development.⁴⁹⁻⁵⁸

Acknowledgments

None.

Conflicts of interest

Authors declare that there is no conflict of interest.

References

1. Singh KK, Mahato S. Influence of indol-3-butyric acid (IBA) and various time on rooting of guava (*Psidium guajava* L.) air layering. *Journal of Plant Development Sciences*. 2016;8(4):193-196.
2. Krishan Kumar Singh. A review: Macro-propagation of guava (*Psidium guajava*). *Journal of Pharmacognosy and Phytochemistry*. 2018;7(2):2629-2634.
3. Samson JA. Tropical fruits 2nd ed. Trop Agr Ser. New York: Longman Scientific and Technical; 1986.
4. Rathore DS. Effect of season on the growth and chemical composition of guava (*Psidium guajava* L.) fruits. *Journal of Horticulture Science*. 1976;51:41-47.
5. Bardiya MC, Kundu BS, P Tauro. Studies on fruit wines Ist. Guava Wine. *Haryana Journal of Horticultural Sciences*. 1974;3(3-4):140-146.
6. Jiménez-Escrig A, Rincón M, Pulido R, et al. Guava Fruit (*Psidium guajava* L.) as a new source of antioxidant dietary fiber. *J Agric Food Chem*. 2001;49(11):5489-5493.
7. Feskanich D, Ziegler RG, Michaud DS, et al. Prospective study of fruit and vegetable consumption and risk of lung cancer among men and women. *J Natl Cancer Inst*. 2000;92:1812-1823.
8. Gordon MH. Dietary antioxidants in disease prevention. *Nat Prod Rep*. 1996;13(4):265-273.
9. Halliwell B. Antioxidants in human health and disease. *Anu Nutr Rev*. 1996;16:33-50.
10. Murashige T. Plant propagation through tissue culture. *Ann Rev Plant Physiol*. 1974;25:135-166.
11. Hammerschlag FA. Factors affecting establishment and growth of peach shoot *in vitro*. *Hort Sci*. 1982;17:85-86.
12. Bhuria P, Yogendra Singh S, Tiwari M, et al. Multiple shoot regeneration from nodal explant of Ashwagandha (*Withania somnifera*). *The Indian Research Journal of Genetics and Bio Technology*. 2014;6(3):521-525.
13. Srivastava B, Yogendra Singh, Tiwari S, et al. "Studies on Micropropagation of Sugarcane from leaf roll, eye bud and meristem tip". *The Indian Research Journal of Genetics and Bio Technology*. 2014;6(1):319-323.
14. Amin MN, VS Jaiswal. Clonal propagation of guava through *in vitro* shoot proliferation on nodal experiments of mature trees. *Plant Cell Tissue Organ and Culture*. 1987;9:235-244.
15. Kumar R. *In vitro* cloning of Chinese guava (*Psidium friedrichsthalianum*

- (Berg.) Nierdz.) Pantnagar: PhD Thesis Submitted to G. B. Pant University of Agriculture and Technology; 2001. 120 p.
16. Loh CS, Rao AN. Clonal propagation of guava (*Psidium guajava* L.) from seedlings and grafted plants and adventitious shoot formation in vitro. *Scientia Horticulturae*. 1989;39(1):31–39.
 17. Papadatau P, Pontikis CA, Ephtimiation E, et al. Rapid multiplication of guava seedlings by in vitro shoot tip culture. *Scientia Horticulturae*. 1990;45:99–105.
 18. Parkash H, Tiwari JP. Micropropagation of guava (*Psidium guajava* L.). *J Applied Horticulture*. 1996;2(1–2):98–101.
 19. Singh SK, Singh SP, Sharma HC. In vitro clonal propagation of guava (*P. guajava* L.) from field grown mature plants. *Physiol Mol Biol Plants*. 2001;7:33–38.
 20. Meghwal PR, Singh SK, Sharma HC. Micropropagation of aneuploid guava. *Indian Journal of Horticulture*. 2003;60(1):29–33.
 21. Bisen BP. *In vitro cloning of Brazilian Guava (Psidium guineense Sw.)*. Pantnagar: Ph.D. Thesis submitted to G.B. Pant University of Agriculture and Technology; 2004. 114 p.
 22. Zamir R, Ali N, Shah ST, et al. In vitro regeneration of guava (*Psidium guajava* L.) from shoot tips of mature trees. *Pakistan Journal of Botany*. 2007;39(7):2395–2398.
 23. Rai MK, VS Jaiswal, U Jaiswal. Encapsulation of shoot tips of guava (*Psidium guajava* L.) for short-term storage and germplasm exchange. *Scientia Horticulturae*. 2008;118(1):33–38.
 24. Xiaomei L, Guochen Y. Clonal propagation of guava (*Psidium guajava* L.) on nodal explants of mature elite cultivar. *International Journal of Plant Biology*. 2011;2:121–125.
 25. Usman M, Butt M, Fatima B. Enhanced in vitro multiple shoot induction in elite Pakistani guava cultivars for efficient clonal plant multiplication. *African Journal of Biotechnology*. 2012;11(44):10182–19187.
 26. Mohamed-Yasseen Y, Barringer SA, Schnell RJ. In vitro shoot proliferation and propagation of guava (*Psidium guajava* L.) from germinated seedlings. *Plant Cell Rep*. 1995;14(8):525–528.
 27. Shah ST, Zamir R, Ahmad J, et al. *In vitro* regeneration of plantlets from seedling explants of guava (*Psidium guajava* L.) cv. Safeda. *Pak J Bot*. 2008;40:1195–1200.
 28. Kumar R, Tiwari JP. Effect of chemicals, light intensity and its duration on in vitro establishment of Chinese guava (*Psidium friedrichsthalianum*). *Progressive Horticulture*. 2001;33(1):1–6.
 29. Singh SK, Meghwal PR, Sharma HC, et al. Direct shoot organogenesis on hypocotyls explants from in vitro germinated seedlings of *Psidium guajava* L. cv. Allahabad Safeda. *Scientia Horticulturae*. 2002;95:213–221.
 30. Amin MN, VS Jaiswal. Micropropagation as an aid to rapid cloning of guava cultivar. *Scientia Horticulture*. 1988;36(1–2):89–95.
 31. Siddiqui ZM, Farooq SA. Role of anti-oxidants in the elimination of phenolic compounds from the in vitro cultures of *Psidium guajava* L. (guava). *Advances in Plant Sciences*. 1996;9(2):155–158.
 32. Meghwal PR, Sharma HC, Goswami AM, et al. Effect of stock plant etiolation on in vitro phenol exudation during culture establishment (*P. guajava* L.). *Indian Journal of Horticulture*. 2001;58(4):328–331.
 33. Bisen BP, JP Tiwari. Evaluation of surface sterilizing agent on in vitro culture establishment of Brazilian guava (*Psidium guineense SW*). *Progressive Horticulture*. 2006;38(1):55–57.
 34. Fitchet-Purnell M. Dimple guava established in tissue culture. *Inlightingsbulletin- Navorsingsinstituut- vir - citrus-en subtropics vrugte*. 1990;212–215.
 35. Siddiqui ZM, Farooq SA. Tissue culture studies on the nodal explant of guava. *Indian Journal of Horticulture*. 1997;54(4):276–279.
 36. Leon-de-Sierrolta, Arehas S, de-Mareno L, et al. Effect of exposure to sunlight of stock plants on the in vitro culture initiation of guava (*Psidium guajava* L.). *Revista-de-la-faculad-de-agronomiauniversidad-del-zulia*. 1997;14(1):531–539.
 37. Khattak MS, Malik MN, Khan MA. Guava propagation via in vitro technique. *Sarhad Journal of Agriculture*. 2002;18(2):199–202.
 38. Ramirez M, Salazar E. In vitro establishment of nodal segments of guava (*Psidium guajava* L.). *Revista de la Facultad de Agronomia Universidad del Zulia*; 1997;14(5):497–506.
 39. Kumar R, Kanawjia A, Tiwari JP. In vitro cloning of Chinese guava (*P. friedrichsthalianum* (Berg) Nierdz.). Kanpur: International Seminar on rec. *Trend Hi-Tech Horticulture and PHT*; 2004;4–6:74.
 40. Bhojwani SS, MK Razdan. *Plant Tissue Culture: Theory and Practice*. New York: Elsevier Science Publishing Company, 1992. p. 502.
 41. Raziuddin Shah, SS Farhad, Sardar Ali. Micropropagation of guava through shoot culture. *Sarhad J Agri*. 2004;20(4):523–527.
 42. Kumar S. Studies on *in vitro* propagation in guava (*Psidium guajava* L.). Hisar: PhD Thesis submitted to CCS Haryana Agricultural University; 2003.
 43. Biswas BK, Joshee N, Yadav A, et al. Development and application of biotechnology in guava: a nutraceutical fruit. In: Desjardins Y, editors. *ISHS Acta Hort. 744: 1st international symposium on human health effects of fruits and vegetables*, 2007. p. 267–276.
 44. Larkin PJ, Scowcroft WR. Somaclonal variation: a novel source of variability from cell cultures for plant improvement. *Theoretical and Applied Genetics*. 1981;60:197–214.
 45. Svabova L, Lebeda A. In vitro selection for improved plant resistance to toxin-producing pathogens. *Journal of Phytopathology*. 2005;153:52–64.
 46. Alizadeh M, SK Singh. Molecular assessment of clonal fidelity in micropropagated grape (*Vitis* spp.) rootstock cultivars using RAPD and ISSR markers. *Iranian Journal of Biotechnology*. 2009;7(1):37–44.
 47. Liu X, Yang G. Assessment of clonal fidelity of micro-propagated guava (*Psidium guajava*) plants by ISSR Markers. *Australian Journal of Crop Science*. 2012;6(2):291–295.
 48. Kamle M, Kumar P, Bajpai A, et al. Assessment of genetic fidelity of somatic embryogenesis regenerated guava (*Psidium guajava* L.) plants using DNA-based markers. *New Zealand Journal of Crop and Horticultural Science*. 2013.
 49. Concepcion O, Napoles L, AT Perez. The effects of three antioxidants on the in vitro culture of guava (*Psidium guajava* L.) shoot tips. Relationship between explant origin and phenolic compound content. *Cultivos Tropicales*. 2005;26:33–39.
 50. Fitchet M. Tissue culture of guava. *Information Bulletin. Citrus and Subtropical Fruit. Res Instt South Africa*. 1989;201:4–5.
 51. Fougat RS, SB Pandya, T Ahmad, et al. In vitro studies in pomegranate (*Punica granatum* L.). *Journal of Applied Horticulture*. 1997;3(1–2):23–29.
 52. Hartmann HT, Kester DE. *Plant propagation: Principles and practices, 4th ed*. New Delhi: Prentice Hall of India Pvt. Ltd; 1983.
 53. Joshee NM, Mutua, AK Yadav. In vitro shoot bud induction and plantlet regeneration in guava as influenced by genotype. *Acta Horticulture*. 2004;632:279–285.
 54. Khattak MS, Malik MN, Khan MA. Effect of surface sterilization agents on in vitro culture of guava. *Sarhad Journal of Agriculture*. 1990;6:151–154.

55. Mangal M, Sharma D, Sharma M. In vitro plantlet regeneration in guava from nodal segments. *Phytomorphology*. 2008;58:103–8.
56. Murashige T, Skoog F. A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiol Plant*. 1962;15:473–497.
57. Singh KK. Rooting response of Guava (*Psidium guajava* L.) through cutting under Garhwal Himalayan region. *Journal of Plant Development Sciences*. 2016;8(3):167–169.
58. Singh R, Tiwari JP. In vitro clonal propagation of jackfruit (*Artocarpus heterophyllus* Lamk.). *Indian Journal of Horticulture*. 1998;55:213–217.