

Diversity, abundance and habitat association of avifauna in Menagesha Amba Mariam and Gara Medhanialem forest, in Oromia Region, Ethiopia

Abstract

The study was conducted from August 2018 to March 2019 by considering the wet and dry seasons. The aim of this study is to investigate diversity, abundance and habitat association of bird species in Menagesha Amba Mariam and Gara Medhanialem forest. Stratified random sampling technique was employed. Point transect techniques was applied in forest and woodland habitats and line transects technique was used in the farmland habitat. Shannon diversity index and chi-square test were employed for data analysis. A total of 112 bird species that belong to 16 orders and 45 families were recorded. Three are endemic to Ethiopia and Twelve species endemic to both Ethiopia and Eritrea. The highest species diversity ($H'=3.60$) was recorded from the forest habitat and the lowest ($H'=2.95$) in the farm land. The association of bird species with habitat was statistically significantly different in wet season ($\chi^2=1702.9$, $df=180$, $p<0.001$) and in dry season ($\chi^2=1497.5$, $df=172$, $p<0.001$). Availability of food and nesting sites were the major factors determining the diversity of birds. Government officials and local community participation is needed for conservation of Menagesha Forest resources.

Keywords: bird species, habitat type, Menagesha forest and species similarity

Volume 6 Issue 1 - 2021

Tamenut Desalegn,¹ Shimelis Aynalem,² Nega Tassie³

¹Bahir Dar University College of Agriculture and Environmental Sciences, School of Fisheries and Wildlife Management, Ethiopia

²Dean of Bahir Dar University College of Agriculture and Environmental Sciences, Ethiopia

³Dean of Biology Department at Bahir Dar University, Ethiopia

Correspondence: Shimelis Aynalem, Dean of Bahir Dar University College of Agriculture and Environmental Sciences, Ethiopia, Tel 0918008194, Email shimelis.aynalem@gmail.com

Received: November 02, 2020 | **Published:** January 29 2021

Introduction

Ethiopia is one of the top 25 biodiversity rich countries in the world and hosts two of the African biodiversity hotspots, namely the eastern Afromontane and the horn of Africa hotspots.¹ The two hotspots comprise highland massive surrounded by arid lowland and contains various wildlife and their habitats ranging from 116 meters (b.s.l) in the Afar Depression to 4620 meters (a.s.l) at Ras Dajen² It is a significant regional center for biological diversity due to its wide ranges of elevation and wider variations of topography.³ The major geographic features are massive highland mountains and plateau divided by the Great Rift Valley and surrounded by lowlands along the periphery. The Great Rift Valley runs from Ethiopia extending in to Tanzania and Great Lakes region of Africa, in the south creating a vast depression.² Those natural factors described above powerfully influenced Ethiopia's extraordinary range of terrestrial and aquatic ecosystems, and contributed to be endowed with a high diversity and rate of endemism.⁴ Consequently, the subsistence of such diverse ecosystems has gifted Ethiopia to be known for its avifauna diversity.

Worldwide, there are over 10, 000 species of birds⁵ that belonging to 204 families.⁶ More than 50 percent of the extant avian species belong to the order Passeriformes. There are over 2,341 bird species in Africa.⁷ Ethiopia has 867 species of birds, where 19 are endemic, 38 are globally threatened, 1 is introduced and 13 species restricted to the geographical region of Ethiopian and Eritrean highlands and thus, shared only by Ethiopia and Eritrea.⁸

Birds perform several ecological roles for ecosystem functioning and produce great benefits for human population in some circumstances.⁹ In the last decades, many studies have assessed an array of ecosystem functions mediated by birds, such as pollination,¹⁰ control of insect populations¹¹ in different ecosystems. About half of the plants are clearly adapted to bird pollination as well as indicates environmental health and they have been considered as environmental indicator species.¹² They are also called bio indicators of climate

change and water quality and also potential tourist attractions.¹³ The abundance of birds as bio indicator species showed strong and significant positive correlations with both tree abundance and cover in each vegetation type.¹⁴

In Ethiopia, there are many designated protected areas (PAs) including National Parks, Wildlife Reserves, Controlled Hunting areas, Priority Forest area, Biosphere Reserves and Community Conservation Areas.¹⁵ Protected areas have a vital role for conservation and protection of biodiversity. Before 1970 Ethiopia has two protected areas but now it has 67 protected areas to conserve natural ecosystems and wildlife heritage.¹⁶ There are 24 National Parks, 3 Biosphere Reserve, 2 Wildlife Sanctuaries, 6 Wildlife Reserve, 5 Community Conservation Areas, 21 Controlled Hunting Areas and 6 Open Controlled Hunting areas.¹⁷

Menagesha Amba Mariam and Gara Medhanialem forest is one of the conservations protected area in Oromia Region Ethiopia is rich in flora and fauna.¹⁸ This forest also one of the important bird area in Ethiopia. However, there is little scientific investigation on the avifauna of state forest. Therefore, this study helps to fill the gap on the diversity, abundance and habitat association of avifauna at Menagesha Amba Mariam and Gara Medhanialem forest.

Materials and methods

Description of the study area

This study was carried out at Menagesha Amba Mariam and Gara Medhanialem Forest. The forest is found in the central part of Ethiopia 30km far to the west of Addis Ababa, the capital city of Ethiopia neighbored by Sebeta and Holeta towns. It is located between 9°01' -9°02'N latitude and 38°34' -38°35'E longitude in the Oromia regional state, Ethiopia, between an altitudinal range of 2200-2910m. The total area of the State Forest is 207 ha (Figure 1) of which 31% is under cultivation for growing both annual and perennial crops,

28% woodland while 41% is forest. The study area flanked by Mt. Wochacha to the south, Kolbo Kebele to the north and Sademo town to the west and Gefersa town to the east.




Figure 1 Location Map of the study area.

Flora and fauna

There are 32 plant species, of which African olive (*Olea europaea*) cover (17.7%) followed by African juniper (*Juniperus procera*) (15.8%) Blue bitter-berry (*Olinia rochetiana*) (11.3%) and Tree heath (*Erica arborea*) (9.73%).18

The study area is the habitat of numerous wild animals including, Olive baboon (*Papio Anubis*), Black and White Colobuses (*Colobus guereza*), Spotted hyena (*Crocota crocuta*) and Grivet monkey (*Chlorocebus aethiops*).

Climate

There is no available metrological data in Menagesha Amba Mariam and Gara Medhaniale Forest. But there are two stations near the forest, Sebeta and Addis Ababa (Ayertena) where weather data was obtained. Approximately the climate of the area was the mean of the two stations. The rainfall and temperature condition of the area was defined based on the data collected from 2007-2017 by the National Metrological Agency (NMA). According to the data from NMA, the result of the analysis showed that the mean monthly maximum and minimum temperature of the study area is 23.5°C and 8.6°C respectively. The hottest month is March with maximum temperature of 26.13°C, followed by May 25.19°C and the coldest month is December with a temperature record of 8.5°C (Figure 2).




Figure 2 Mean monthly maximum and minimum temperature and rain fall of Menagesha Amba-Mariam and Gara-Medhaniale Forest (2007-2017).

The mean annual rainfall is 1050mm and the mean annual

temperature is 23.8°C. The rain mainly falls from June to September, with the highest concentration in July and August. June to December is comprises the wet season and December to March the dry season.

Materials used

Materials used during the field survey include 10×50 field binoculars; Geospatial Positioning System (GPS), sound recorder, digital camera and bird guide books.^{19,20}

Sampling design and data collection

A preliminary survey was carried out in July, 2018 to gather basic information about the study site. The physical features of the study area were assessed using ground survey. The actual study was carried out from August, 2018 to March, 2019 by considering both the wet and the dry seasons. The wet season data covers from August to November and the dry season was from December to March. In this study Stratified random sampling design was employed, since the study area was not uniform in terms of habitat types. The area was stratified into three habitats types (Forest, Farmland and Woodland).

A total of 25 sampling plots were purposively placed covering the whole study area 207 hectare. Different types of sampling plots were assigned for each habitat. The sampling plots were selected based on the area proportion of each representative habitat. Each study plot had consisted of 9-point counts. Each plot measures 3.24 hectare to cover 39% of the study area. The radius of each point was 30m and there was 200m distance between plots to avoid double counting of individual birds.²¹

In forest area and wood land point transect technique was used while in the farmland the researcher used line transect count techniques. There were ten plots in the forest, seven plots in woodland and eight plots in the farm land (Table 1). There were 16 lines transect to record the abundance of birds in the farmland.

Table 1 Habitat classification and allocation of plots based on area coverage

Habitat type	Forest	Woodland	Farmland
Area coverage in hectare	84.5ha	58.5ha	64ha
Proportion of sample	41%	28%	31%
Number of plots	10	7	8
Number of point counts for each plot	9	9	-
Total number of point count	9×10=90	7×9=63	-
Total number of lines transect	-	-	8×2= 16

In each site, bird observation was carried out twice daily 6:30 AM to 10:00 AM in the morning and 4:00 PM and 6:00 PM in the afternoon. Birds were counted as seen, heard and when flight on the sky. Field data collection was carried out for three days per week depending on the weather conditions and time of the day when bird species were active. Bird identifications and counting of individuals was conducted by direct observations. Observations were made by standing in the middle of the sampling point and observing 360° round quietly and gently. Upon reaching a point, 2-5 minutes was stayed for the birds to settle in case of any disturbances.

Data analysis

The data was analyzed by using simple descriptive statistics; bird abundance and diversity was calculated using Shannon-Wiener Diversity Index which provides an account for both abundance

and evenness²² and the data was analyzed using SPSS version 20.0 software.

Shannon-Wiener Diversity Index (H') was calculated as:

$$\text{Diversity index } H' = \sum (pi \ln pi) \dots\dots\dots (1)$$

Where,

H' = the Shannon-Wiener Index of Diversity

Pi = the proportion of the species relative to the total number of species

ln= Indicates natural logarithm

Sorensen similarity index (S) used to measure species similarity of different habitat types. It is equal to 1 in case of complete species similarity between three habitats and 0 if species of two habitat types are dissimilar.²³

Sorensen similarity index (S) is computed as:

$$S = 2C / (A + B + D) \dots\dots\dots (2)$$

Where:

A = Number of species that occur in a site A

B = Number of species that occur in a site B

C = Number of common species that occur in a site A, B and D

D=Number of species that occur in a site D

Species evenness is often assessed by Shannon equitability index (E) which is calculated by:

$$E = H' / H_{\max} \dots\dots\dots (3)$$

Where; H'max = ln(S) = natural logarithm of the total number of species (S) in each habitat.²⁴ E values ranges from 0 to 1, in which 1 indicates complete evenness.

Richness index (RI) of each species was determined using the formula:

$$RI = S - 1 / \ln pi \dots\dots\dots (4)$$

Where, S = Number of species in each habitat, ln = Natural logarithm, pi= Number of individual species in each habitat.

Relative abundance of avian species was determined by using encounter rates that give crude ordinal scales of abundance (abundant, common, frequent, uncommon and rare).²⁵

$$\text{Encounter rate} = \frac{\text{Total number of individual birds recorded}}{\text{Period of observation per 100 field hour}} * 100$$

Encounter rate was calculated for each species by dividing the number of birds recorded by the number of hours spent searching, in order to get a figure of birds per hour for each species (Table 2).

Table 2 Encounter rates used to give a crude ordinal scale of abundance

Abundance category	Abundance score	Ordinal scale
<0.1	1	Rare
0.1-2.0	2	Uncommon
2.1-10.0	3	Frequent
10.1-40.0	4	Common
>40.0	5	Abundant

Relative abundance= Total number of individuals of one species (n)/ Total no. of individuals of all species (N) ×100 (5)

Chi-square test (χ^2) was employed to see the association of birds among the three habitat types (Forest, Woodland and Farmland) and Chi-square (χ^2) goodness of fit was used for test abundance of bird species in the study area.

Results

Abundance and composition

A total of 112 avian species that belongs to 16 and 45 family's orders were recorded during the present study period (Table 2). Majority of 67% them are under order Passeriformes. The highest number of species was recorded in families: Accipitridae, Columbidae, Ploceidae and Fringillidae. The lowest number of species was recorded from seventeen Families, 1 species each. Yellow-fronted Parrot (*Poicephalus flavifrons*), Abyssinian Catbird (*Parophasma galinieri*) and Abyssinian Woodpecker (*Dendropicus abyssinicus*) are endemic to Ethiopia. Twelve species are endemic to both Ethiopia and Eritrea. Five species near endemic to Ethiopia and one species is Intra-Africa migrant were recorded, but the remaining 91 species were residents (Table 3).

Table 3 Bird species recorded during the wet and the dry seasons at Amba-Mariam and Gara-Medhaniale forest

Order	Family	Common name	Scientific name
Accipitriformes	Accipitridae	African Fish-Eagle	<i>Haliaeetus vocifer</i> **
		African Hawk-Eagle	<i>Haliaeetus spilogaster</i> +
		Augur Buzzard	<i>Buteo augur</i> **
		Bateleur	<i>Terathopius ecaudatus</i> **
		Black-chested Snake-Eagle	<i>Circaetus pectoralis</i> **
		Hooded Vulture	<i>Necrosyrtes monachus</i> *
		Tawny Eagle	<i>Aquila rapax</i> *
		Verreaux's Eagle	<i>Aquila Verreauxii</i> *
		White-headed Vulture	<i>Trigonoceps occipitalis</i> *
		Yellow-billed Kite	<i>Milvus aegyptius</i> +
Anseriformes	Anatidae	Egyptian Goose	<i>Alopochen aegyptiaca</i> **

Table continued...

Order	Family	Common name	Scientific name	
Apodiformes	Apodidae	Nyanz Swift	<i>Apus niansae</i> +	
Bucerotiformes	Bucerotidae	African Grey Hornbill	<i>Tockus nasutus</i> +	
		Red-billed Hornbill	<i>Tockus erythrorhynchus</i> **	
Coliiformes	Collidae	Speckled Mousebird	<i>Colius striatus</i> **	
Coraciiformes	Meropidae	Blue-breasted Bee-eater♦	<i>Meros lafresnayii</i> +	
		Little Bee-eater	<i>Merops pusillus</i> *	
Columbiformes	Columbidae	African Olive Pigeon	<i>Columba arquatrix</i> *	
		Blue-spotted Wood Dove	<i>Turtur afer</i> **	
		Dusky Turtle Dove	<i>Streptopelia lugens</i> **	
		Laughing Dove	<i>streptopelia senegalensis</i> **	
		Red-eyed Dove	<i>Streptopelia semitorquata</i> +	
		Lemon Dove	<i>Aplopelia larvata</i> *	
		Speckled Pigeon	<i>Columba guinea</i> **	
		Tambourine Dove	<i>Turtur tympanistris</i> *	
		White-collared Pigeon	<i>Columba albitorques</i> **	
		Falconiformes	Falconidae	Lanner Falcon
Galliformes	Phasianidae	Erckel's Francolin♦	<i>Pternistis erckelii</i> **	
Gruiformes	Rallidae	Rouget's Rail	<i>Rougetius rougetii</i> **	
Musophagiformes	Musophagidae	White-cheeked Turaco♦	<i>Tauraco leucotis</i> **	
Passeriformes	Alaudidae	Erlanger's Lark	<i>Calandrella erlangeri</i> **	
	Buphagidae	Red-billed Oxpecker	<i>Buphagus erythrorhynchus</i> **	
	Cisticolidae	Buff-bellied Warbler	<i>Phyllolais pulchella</i> *	
		Grey-backed Camaroptera	<i>Camaroptera brachyura</i> **	
		Red-faced Cisticola	<i>Cisticola erythrops</i> *	
		Tawny-flanked Prinia	<i>Prinia subflava</i> +	
		Corvidae	Cape Rook	<i>Corvus capensis</i> +
			Fan-tailed Raven	<i>Corvus rhipidurus</i> **
	Pied Crow		<i>Corvus albus</i> +	
	Thick-billed Raven		<i>Corvus crassirostris</i> +	
	Estrildidae	African Firefinch	<i>Lagonosticta rubricata</i> **	
		Bronze Mannikin	<i>Spermestes cucullata</i> *	
		Red-cheeked Cordon-bleu	<i>Uraeginthus bengalus</i> **	
		Red-billed Firefinch	<i>Lagonosticta senegalensis</i> **	
		Yellow-billed Waxbill	<i>Coccozygia quartina</i> **	
	Fringillidae	African Citril	<i>Serinus citrinelloides</i> **	
		Brown-rumped Seedeater	<i>Serinus tristriatus</i> **	
		Streaky Seedeater	<i>Serinus striolatus</i> **	
		Stripe-breasted Seedeater	<i>Serinus reichardi</i> **	
		Yellow-crowned Canary	<i>Serinus Serinus flavivertex</i> +	
Yellow-fronted Canary		<i>Serinus mozambicus</i> +		
Hirundinidae	Rock Martin	<i>Ptyonoprogne fuligula</i> *		
	Lesser Striped Swallow	<i>Cecropsis abyssinica</i> **		
Laniidae	Common Fiscal	<i>Lanius collaris</i> **		

Table continued...

Order	Family	Common name	Scientific name
	Locustellidae	Cinnamon Bracken Warbler	<i>cinnamomeus</i> +
	Macrosphenidae	Northern Crombec	<i>Sylvietta brachyura</i> +
	Malaconotidae	Ethiopian Boubou	<i>Laniarius aethiopicus</i> **
		Back-crowned Tchagra	<i>Tchagra senegalus</i> *
	Monarchidae	Abyssinian Slaty Flycatcher ^x	<i>Melaenornis chocolatinus</i> +
		African Dusky Flycatcher	<i>Muscicapa adusta</i> +
		African Paradise Flycatcher	<i>Trochocercus cyanomelas</i> **
	Motacillidae	White Wagtail♣	<i>Motacilla alba</i> *
		Mocking Cliff Chat	<i>Thamnolaea</i>
	Muscicapidae		<i>cinnamomeiventris</i> **
		Moorland Chat ^x	<i>Cercomela sordida</i> **
		Ruppell's Black-Chat	<i>Myrmecocichla melaena</i> +
		Ruppell's Robin-Chat ^x	<i>Cossypha semirufa</i> **
	Nectariniidae	Beautiful Sunbird	<i>Cinnyris pulchellus</i> *
		Malachite Sunbird	<i>Nectarinia famosa</i> **
		Scarlet-chested Sunbird	<i>Chalcomitra senegalensis</i> **
		Shining sunbird	<i>Cinnyris habessinicus</i> **
		Tacazze Sunbird	<i>Nectarinia tacazze</i> **
		Variable Sunbird	<i>Cinnyris venustus</i> **
	Oriolidae	Abyssinian Oriole	<i>Oriolus monacha</i> **
		Black-headed Oriole ^x	<i>Oriolus larvatus</i> +
	Paridae	White-backed Black Tit ^x	<i>Parus leuconotus</i> **
	Passeridae	Swainson's Sparrow♠	<i>Passer swainsonii</i> **
	Phylloscopidae	Brown Woodland Warbler	<i>Phylloscopus umbrovirens</i> +
	Ploceidae	Baglafecht Weaver	<i>Ploceus baglafecht</i> **
		Fan-tailed Widowbird	<i>Euplectes axillaris</i> **
		Little Weaver	<i>Ploceus luteolus</i> +
		Red-collared Widowbird	<i>Euplectes ardens</i> *
		Spectacled Weave	<i>Ploceus ocularis</i> +
		Village Weaver	<i>Ploceus cucullatus</i> +
		Vitelliane Masked Weaver	<i>Ploceus vitellinus</i> +
		Yellow Bishop	<i>Euplectes capensis</i> **
		Yellow-crowned Bishop	<i>Euplectes afer</i> *
		Yellow-mantled Widowbird	<i>Euplectes macroura</i> *
	Pycnonotidae	Common Bulbul	<i>Pycnonotus barbatus</i> **
	Sturnidae	Greater Blue-eared Starling	<i>Lamprotornis chalybaeus</i> +
		Red-winged Starling	<i>Onychognathus morio</i> **
	Sylviidae	Abyssinian Catbird♣	<i>Parophasma galineri</i> **
	Timaliidae	White-rumped Babbler♠	<i>Turdoides leucopygia</i> **
	Turdidae	African Thrush	<i>Turdus pelios</i> +
		Mountain Thrush	<i>Turdus abyssinicus</i> **
		Groundscraper Thrush	<i>Psophocichla litsitsirupa</i> **
		White-winged Cliff Chat ^x	<i>Thamnolaea semirufa</i> +

Table continued...

Order	Family	Common name	Scientific name
	Viduidae	Pin-tailed Whydah	<i>Vidua macroura</i> +
		Village Indigobird	<i>Vidua chalybeata</i> +
	Zosteropidae	Abyssinian White-eye	<i>zosterops abyssinicus</i> **
		Yellow White-eye	<i>Zosterops senegalensis</i> **
Pelecaniformes	Threskiornithidae	Hadada Ibis	<i>Bostrychia hagedash</i> +
		Wattled Ibis \times	<i>Bostrychia carunculata</i> **
Piciformes	Scopidae	Hamerkop	<i>Scopus umbretta</i> +
	Indicatoridae	Lesser Honeyguide	<i>Indicator minor</i> *
	Picidae	Abyssinian Woodpecker \clubsuit	<i>Dendropicus abyssinicus</i> **
		Cardinal Woodpecker	<i>Dendropicos fuscescns</i> *
Grey-headed Woodpecker		<i>Dendropicos spodocephalus</i> +	
Psittaciformes	Psittaculidae	Nubian Woodpecker	<i>Campethera nubica</i> *
		Black-winged Lovebird \times	<i>Agapornis taranta</i> **
		Yellow-fronted Parrot \clubsuit	<i>Poicephalus flavifrons</i> **
Trogoniformes	Trogonidae	Narina Trogon	<i>Apaloderma narina</i> *

*=Dry; +=wet; **both Wet and Dry; \clubsuit Endemic; \times Endemic to Ethiopia and Eretria; \diamond Near Endemic; \spadesuit Inter Africa migrant; unmarked species are resident birds)

Forest habitat had high number of individuals in both seasons followed by woodland and least number of individuals recorded at the farmland habitat (Figure 3).




Figure 3 Abundance of birds in the three habitat types.

The Chi-square test for abundance of birds during the study period in the two seasons showed that there was statistically significant difference between observed and expected counts of each species. The abundance of bird chi-square test during the wet season ($\chi^2=1504.8$, $df=90$, $p<0.001$) and the dry season ($\chi^2=1842.7$, $df=86$, $p<0.001$).

During the wet and dry seasons, 90 and 81 bird species were recorded, respectively. Among them 59 bird species were common for both seasons, while 22 and 31 species were exclusively recorded during the wet and the dry seasons, respectively (Table 2).

The dominant bird species recorded in the study area were Brown-rumped Seadeater ($n=198$), Swainson's Sparrow ($n=127$), Ruppell's Robin-Chat ($n=122$) and Abyssinian Oriole (98).The number of bird species in the forest habitat was higher compared to the Woodland and Farmland (Figure 4).

Species diversity

The highest species diversity ($H'=3.60$) of birds was recorded from the forest habitat during the wet season, followed by woodland habitat ($H'=3.49$), whereas the least diversity ($H'=3.37$) was recorded from the farmland habitat during the wet season. During the dry season, the highest species diversity was recorded in the forest habitat

($H'=3.36$) and the least was in the farmland ($H'=2.95$). During the wet season, the highest and the lowest species evenness was recorded in woodland (0.95) and farmland (0.86) habitats, respectively. The woodland habitat had high species evenness but the least evenness was registered from the farmland habitat in both seasons.




Figure 4 Number of bird species recorded by habitats and seasons.

The highest richness index was recorded ($RI=7.61$) from the forest habitat and the least richness ($RI=4.79$) was recorded from the farmland habitat (Table 4). The highest number of species was recorded in the forest habitat in both seasons. A total of 50 and 44 species were recorded in the forest habitat in wet and dry seasons, respectively. But the smallest number of bird species was recorded from the farmland habitat 29 during dry season.

Relative abundance of birds in wet and dry seasons

During the study period 34, 18 and 22 species were uncommon; 15, 21 and 17 species were frequent and 1 species was recorded common in each habitat during the wet season. Likewise, during the dry season 26, 12 and 14 species were uncommon; 17, 22 and 14 species were frequent; 1 species was common at each habitat (Table 5).

There was 40% avian species similarity between forest and woodland habitats during the wet season. But 8.2% avian species similarity was observed between farmland and forest habitat during the dry season. Similarity decrease as we go from farmland with woodland, farmland with woodland and farmland with forest,

respectively in both seasons (Table 6 & 7). The three habitat types had 9.2% and 5.6% avian species similarity during the wet and dry seasons, respectively.

(forest, woodland and farmland) showed that there was statistically significantly different, in wet season ($\chi^2=1702.9$, $df=180$, $p<0.001$) and in dry season ($\chi^2=1497.5$, $df=172$, $p<0.001$) across in the studied habitats throughout the study period.

The chi-square test for habitat association of bird species during the study period in the two seasons with the three habitat types

Table 4 Bird species abundance, diversity, richness and evenness during the wet and dry season

Habitat	Season	No. of species	Abundance (no. of individuals)	D	RI	H'	H'/Hmax
Forest	Wet	50	625	0.959	7.61	3.6	0.92
	Dry	44	387	0.943	7.22	3.36	0.89
Woodland	Wet	40	445	0.958	6.39	3.49	0.95
	Dry	35	384	0.946	5.71	3.24	0.91
Farmland	Wet	40	433	0.947	6.42	3.37	0.91
	Dry	29	342	0.921	4.79	2.95	0.86

H', Shannon-weiner Index; $D, 1 - \sum p_i^2$; RI=Richness Index; H'/Hmax=Evenness; Hmax=Ln(S); S, The total number of species

Table 5 Number of bird species between different habitat relative abundance categories

Habitat	Season	Uncommon	Frequent	Common
Forest	Wet	34	15	2
	Dry	26	17	1
Woodland	Wet	18	21	1
	Dry	12	22	3
Farmland	Wet	22	17	1
	Dry	14	14	2

Table 6 Sorensen similarity index (S) of bird species between different habitats during the wet season

Habitats	Common species	Similarity index
Farmland with Forest	9	0.2
Farmland with Woodland	10	0.25
Forest with Woodland	19	0.4
Forest with woodland with Farmland	4	0.092

Table 7 Sorensen similarity index (S) of bird species in different habitats during the dry season

Habitat	Common species	Similarity index
Farmland with Forest	3	0.082
Farmland with Woodland	5	0.156
Forest with Woodland	13	0.329
Forest with woodland with Farmland	2	0.056

Discussion

A total of 112 avian species that belong to 45 families were recorded during the present study period with large proportion of species recorded under order Passeriformes. In line with this study, Kalkidan & Afework²⁶ and Girma et al.²⁷ found that Passeriformes was the most dominant order. Likewise, the recent report by Seyoum et al.²⁸ also pointed out that order Passeriformes was numerically the dominant order represented with 22 species which accounts for 44% of the identified species. Ploceidae and Colombidia were the most dominant families in the present study. This might be due to habitat

quality, flower composition, high productivity of fruits, availability different nesting vegetation's.²⁹ Likewise, Gloria³⁰ reported that birds select habitats that fit their requirements for successful reproduction and survival although some generalist species may utilize several habitats.

The diversity of birds varied within the three habitat types, which could be due to vegetation type, foliage height, nesting availability and floristic composition of the area. Similarly, Waterhouse et al.³¹ showed that the food and cover requirement of bird species is determined mainly by the vegetation structure and composition. Soderstrom & Part³² in Sweden reported that birds often prefer to

utilize multiple habitats and depend on the quality and productivity of habitats in terms of food availability, shelter and breeding areas in order to maintain viable population. In addition to these Akobu et al.³³ reported that the distribution of nectarivorous birds positively associated with habitat of complex vegetation in the Assop forest that had higher plant diversity.

In this study the diversity, abundance and habitat association of birds are determined in terms of water availability, feeding requirement, floristic composition and nesting habitats. Similarly, Rodriguez-Estrella³⁴ reported that the pattern of bird distribution is strongly related to environmental factors (topography, food and habitats) and human interventions. Meseret & Solomon³⁵ also supported the present study that wild species prefer to habitats where their ecological needs are fulfilled.

The highest numbers of bird species were observed in the forest habitat during both the wet and dry seasons. Kalkidan & Afework²⁶ in Entoto Natural Park and Escarpment also reported that the highest numbers of avian was observed in the forest habitats. This is possibly due to the diversity of vegetation that provided for different bird species. In addition to that Hailemariam et al.³⁶ suggested that species composition and abundance of birds were mainly determined by vegetation structures that used as breeding sites, food source and shelter. In recent study Dagnaw & Mesele³⁷ in Zengo Forest also reported that, the highest diversity of bird species was observed at the forest habitat. Furthermore, Eshetu et al.³⁸ had also reported that the highest abundance of birds was recorded in dense and ticket forest habitat types. But the lowest abundance was recorded in farmland habitat types. These might be related with the fact that forest habitats are much conducive for birds in the availability of food and roosting sites.

In contrary to this study, Zerihun et al.³⁹ stated that Wondo Genet Forest, natural forest which is dominated by few tree species is not suitable to different bird species, because it does not fulfill the feeding preferences of most birds together with risk of predation and it is preferred by few forest specialist bird species alone as they obtained sufficient cover and food necessities. The second highest bird individuals were recorded at the woodland habitat in present study area. In line with this study, Gloria³⁰ reported that bird species diversity was higher in forest followed by woodland and shrub land than in the settlement and farmland. This might be differences in resource availability between habitats such as breeding sites, nesting material, cover, food and water restrict some species to certain habitat type while allowing others to be widely distributed.

The small number of bird species was recorded from the farmland habitat. This might be due to habitat modification for agricultural activities, road construction, removing of nesting grounds, spraying of herbicides on agricultural areas, transmitted diseases and computation of birds with domestic animals. Aggeliki et al.⁴⁰ reported that farmland biodiversity continues to decline mainly because of agricultural intensification and land abandonment.

Furthermore, Anteneh et al.⁴¹ reported that permanent removal of forest components in farmland destroys species foraging and nesting resources and unnaturally enhances exposure to natural enemies such as predators and also destabilizes the balance of processes such as interspecific competition and co-existence. In contrary to this Henderson et al.⁴² found out that land use change is a factor in both declining and increasing of bird populations. For instance, the introduction of set-aside has resulted in a relatively large area of farmland being converted in to a habitat that is preferred by a number

of farmland bird species.

In the case of species similarity, communities showed the highest similarity during the wet season than dry. The highest species similarity was observed between forest and woodland in wet season. This might be related to the availability of insects, flower composition, high canopy cover and presence of fruit. In addition to this they occur in similar locality could be occupied by similar bird species and generalist species interchanging between the two habitat types. In agreement with the findings of Kamal et al.⁴³ avian species diversity and richness and similarity were positively correlated with the similarity in vegetation type and foliage height.

In present study high species evenness was observed at the woodland habitat. It might be related to habitat homogeneity. Species evenness might be closely linked to habitat complexity or heterogeneity; greater heterogeneity might explain the relatively low species evenness.⁴⁴ The lowest evenness was recorded from the farmland. This is supported by Wilsey & Stirling⁴⁵ that the increase competition particularly can result in dominant species having greater proportional abundances leading to decreased evenness.

Large number of forest bird species were found to be ranked under uncommon based on their relative abundance scores during the wet season. This might be related to vegetation complexity, inconspicuousness of small birds, roosting and feeding. Similarly, Shimelis & Afework⁴⁶ stated that the presence of high number of uncommon bird species in Zegie Peninsula could be related to cutting of trees, clearing vegetation for coffee plantation and firewood production to sell to the nearest town; those activities could reduce the individual bird species. Likewise, Kalkidan & Afework²⁶ reported that the relative abundance of birds in the forest habitat showed large number of bird species grouped as uncommon. This might be due to the vegetation complexity and inconspicuousness of small birds. In recent study Seyoum et al.²⁸ also reported that the majority of 74% of birds were registered under uncommon species. This might be due to majority of the species had low population sizes as a result they were grouped under uncommon species. Only five bird species Swainson's Sparrow, Little Weaver, Brown-rumped Seadeater, Ruppell's Robin-Chat, Abyssinian Oriole are grouped under common species because they had relatively a greater number of individuals.

In the study period rare species was not observed, this might be the area less comfortable for the specialist bird species and the absence of rarity was the area support birds and related to the habitat condition.

Conclusion

The present study documented base line information about avifauna in Menagesha Amba Mariam and Gara Medhaniale forest. The occurrence of 112 species of birds confirms the importance of Menagesha forest for biodiversity protection. The study area, in spite of small in size it supports endemic, migratory and near endemic species where they get better protection and good access for foraging and nesting opportunities. Habitat type determines bird species diversity, abundance, evenness, similarity and richness. Birds are associated with different habitat types where basic requirements like nesting grounds, roosting sites water and food resources are available.

In both dry and wet seasons, the highest species diversity richness and individual abundance was recorded from the forest habitats, and the highest evenness recorded at the woodland habitat. This indicates that forest and woodland habitats are suitable for birds by providing cover nesting materials and feeding requirements. Bird diversity, abundance and evenness were high in wet season than dry seasons.

The small number of bird species diversity, individual abundance and evenness were recorded from the farmland habitat. In the farmland habitat there are different human activities practiced that affect bird's diversity and abundance, such as farming, livestock grazing, cutting of trees for charcoal, road construction, erosion and need for additional farming area.

Recommendations

- i. Local communities have not enough awareness about the importance of birds, awareness creation campaigns should be practiced at least to the local communities living within and around the state forest.
- ii. Even though the study area is relatively protected by regional government, some illegal activities are being practiced by the local communities so the communities should get train to ensure conservation of the area.

Acknowledgments

The author would like to thank everyone who helped him to write his thesis successfully and also thanks Dr. Shimelis Aynalem and Dr. Nega Tassie for their professional guidance, intellectual stimulation, which made my study successful. Besides, I would like to thank Dr. Mezgebu Ashagrie for his support in statistical analysis and Belayneh Abebe for providing me a bird guide book for the field data collection. In addition to these thanks the local communities support me during data collection and sharing ideas. Finally, Special thanks go to his parents for their financial support.

Conflicts of interest

The author declares that there is no conflict of interest.

Funding

None.

References

1. Conservation International. Biodiversity hotspots. 2011.
2. Husen A, Mishra VK, Semwal K, et al. Biodiversity Status in Ethiopia and Challenges. In: Bharati KP, Chauhan A, Kumar P, editors. *Environmental Pollution and Biodiversity*. New Delhi, India: Discovery Publishing House Pvt Ltd; 2012;1:31–79.
3. Kelbessa E, Demissew S. Diversity of vascular plant taxa of the flora of Ethiopia and Eritrea. *Ethiopian Journal of Science*. 2014;13(1S):37–45.
4. Tesfaye A, Gashaw M, Tesfaye G, et al. Ecosystems of Ethiopia. National Biodiversity Strategy and Action Plan (NBSAP) Project, IBCR, Addis Ababa, Ethiopia. 2003:103.
5. Bird Life International. Handbook of the Birds of the World Bird species distribution maps of the world. 2016.
6. George WC. *Bird migration and climate change*. 2nd ed. Island Press, Washington, DC. 2010.
7. Bird Life International. State of Africa's bird: Outlook for our changing environment. Nairobi, Kenya: BirdLife International Africa Partnership. 2013.
8. Lepage D. Avibase Bird Checklists of the World Ethiopia. Birdlife International. 2019.
9. Jose CMF, Deborah F. An appraisal of bird-mediated ecological functions in a changing world. *Tropical Conservation Science*. 2017;10:1–12.
10. Anderson SH, Kelly D, Ladley JJ, et al. Cascading effects of bird functional extinction reduces pollination and plant density. *Science*. 2011;331(6020):1068–1071.
11. Van Bael SA, Bichier P, Barber NA, et al. Birds as predators in tropical agro-forestry systems. *Ecology*. 2008;89(4):928–934.
12. Rajashekara S, Venkatesha GM. Community composition of aquatic birds in lakes of Bangalore, India. *J Environmen Biol*. 2011;32(1):77–83.
13. EWNHS. Important Bird Areas of Ethiopia. A first inventory. Ethiopian Wildlife and Natural History Society. Addis Ababa. 1996:300.
14. Asefa A, Mengesha G, Yosef Mamo Y. Application of birds as ecological bioindicators for monitoring habitat change: A case study from Abijata-Shalla lakes national park, Ethiopia. *International Journal Art Science*. 2016;4(2):112–121.
15. Ethiopian Wildlife Conservation Authority (EWCA). First enhancing our Heritage Assessment at Simien Mountains National Park (SMNP), WHS. 2014.
16. Murray M, Admasu B. Development of a Marketing Strategy for Wildlife Tourism in Ethiopia. *Natural Resources*. 2013:65.
17. Vreugdenhil D, Vreugdenhil AD, Tamirat Tilahun, et al. Gap Analysis of the Protected Areas System of Ethiopia, with technical contributions from Nagelkerke L, Gedeon K, Spawls S, Yalden D, Lakew Berhanu and Siegf L. World Institute for Conservation and Environment, USA. 2012.
18. Tilahun A, Teshome Soromessa T, Kelbessa E. Structure and Regeneration Status of Menagesha Amba Mariam Forest in Central Highlands of Shewa, Ethiopia. *Agriculture, Forestry and Fisheries*. 2015;4(4):184–194.
19. Redman N, Stevenson T, Fanshawe J. Birds of the Horn of Africa Ethiopia, Eritrea, Djibouti, Somalia, Socotra. Princeton University Press; Revised ed. 2009:512.
20. Aynalem S. Birds of Lake Tana Area Ethiopia. A photographic Field Guide. 2013:235.
21. Sutherland WJ. *Ecological Census Techniques*. 2nd ed. Cambridge: Cambridge University press; 2016:320–326.
22. Shannon CE, Wiener N. *The Mathematical Theory of Communication*. The University of Illinois, Urbana; 1999:117.
23. Krebs CJ. *Ecological Methodology*. 2nd ed. California: Addison-Welsey Educational Publishers Inc; 1999:620.
24. Tramer EJ. Bird species diversity: components of Shannon's formula. *Ecology*. 1969;50:929–931.
25. Bibby C, Jones M, Marsden S. Expedition Field Technique. Bird surveying manual. Royal Geographical Society of London. 1998.
26. Esayas K, Bekele A. Species Composition, Relative Abundance and Distribution of the avian fauna of Entoto Natural Park and Escarpment, Addis Ababa. *Ethiopian Journal of Science*. 2011;34(2):113–122.
27. Mengesha G, Mamo Y, Bekele A. A comparison of terrestrial bird community structure in the undisturbed and disturbed areas of the Abijata Shalla lakes national park, Ethiopia. *International Journal of Biodiversity and Conservation*. 2011;3(9):389–404
28. Kiros S, Afework B, Legese K. A preliminary study on bird diversity and abundance from Wabe fragmented forests around Gubre subcity and Wolkite town, Southwestern Ethiopia. *International Journal Avian and Wildlife Biology*. 2018;3(5):333–340.
29. Tassie N, Bekele A. Diversity and habitat association of birds of Dembia plain wetlands, Lake Tana, Ethiopia. *SINET, Ethiopian Journal of Science*. 2008;31(1):1–10.
30. Gloria B. Diversity, Distribution and Abundance of Avifauna in respect to habitat types. M.Sc. Thesis, Kilakala and Bigwa, Morogoro, Tanzania. 2013:5–33.

31. Waterhouse FL, Mather MH, Seip D. Distribution and abundance of birds relative to elevation and biogeoclimatic zones in coastal old-growth forests in southern British Columbia B.C. *Journal of Ecosystems and Management*. 2002;2:1488–4674.
32. Soderstrom B, Part T. Influence of landscape scale on farmland birds in semi-natural pastures. *Conservation Biology*. 2000;14:522–533.
33. Akobu SA, Ihuma JO, Manu S. Survey of abundance and diversity of avian species in Assop Forest Reserve and surrounding Farmlands in Jos, Nigeria. *International journal*. 2012;2(3):506–514.
34. Rodriguez-Estrella R. Land use changes affect distributional patterns of desert birds in the Baja California peninsula, Mexico. *Diversity and Distribution*. 2007;13:877–889.
35. Chanea M, Yirga S. Diversity of Medium and Large-sized Mammals in Borena-Sayint National Park, South Wollo, Ethiopia. *International Journal of Sciences: Basic and Applied Research*. 2014;15(1):95–106.
36. Areaya H, Yonas M, Hadush HT. Community composition and abundance of residential birds in selected church forests, Tigray Region, Northern Ethiopia. *Academic journal*. 2013;8(22):1038–1047.
37. Belay D, Yihune M. Diversity, Distribution and Relative Abundance of Avian Fauna in and Around Zengo Forest, East Gojjam, Ethiopia. *International Journal of Ecology and Environmental Sciences*. 2017;43(4):287–293.
38. Moges E, Masersha G, Tebaber Chanie T, et al. Species diversity, habitat association and abundance of avifauna and large mammals in Gonde Teklehimant and Aresema monasteries in North Gondar, Ethiopia. *International Journal of Biological Conservation*. 2018;10(4):185–191.
39. Girma Z, Mamo Y, Mengesha G, et al. Diversity, relative abundance and distribution of avian fauna in and around Wondo Genet forest, South-central Ethiopia. *Research Journal of Forestry*. 2017;11(1):1–12.
40. Aggeliki D, Maria Luisa P, Philippe P, et al. Low-intensity agriculture increases farmland bird abundances in France. *Journal of Applied Ecology*. 2010;47(6):1348–1356.
41. Aynalem S, Mitchell P, Addisu Asefa. Impact Forest Structure Simplification on Bird Species Richness in the Harena Forest of the Bale Mountains National Park (BMNP), South Eastern Ethiopia. *An International Quarterly Scientific Journal*. 2013;12:321–324.
42. Henderson IG, Vickery JA, Fuller RJ. Summer bird abundance and distribution on set-aside fields on intensive arable farms in southern England. *Ecography*. 2000;23:50–59.
43. Kamal K, Dinesh B, Aashish T. Avian diversity and its association with vegetation structure in different elevational zones of Nainital district (Western Himalayan) of Uttarakhand. *International Journal of Biodiversity and Conservation*. 2012;4(11):364–376.
44. Willig MR, Kaufman DM, Stevens RD. Latitudinal gradients of biodiversity: pattern, process, scale and synthesis. *Annual Review of Ecology, Evolution, and Systematics*. 2003;34:273–309.
45. Wilsey B, Stirling G. Species richness and evenness respond in a different manner to propagule density in developing prairie microcosm communities. *Plant Ecology*. 2007;190:259–273.
46. Aynalem S, Bekele A. Species Composition, Relative Abundance and Habitat Association of the bird fauna of the Montane Forest of Zegie Peninsula and nearby Islands, Lake Tana, Ethiopia. *Ethiopian Journal of Science*. 2009;32(1):45–56.