Resveratrol normalizes hyperammonemia induced proinflammatory and pro-apoptotic conditions in rat brain

Abstract

Chronic liver failure (CLF) led hyperammonemia (HA) is known to develop a metabolic brain disorder known as hepatic encephalopathy (HE). TNF- α is now considered responsible for mounting neurological complications associated with HA/HE. However, the mechanism that connects inflammation and neuroexcitotoxicity is not yet clear. Resveratrol (RSV) is a natural antioxidant and known to mediate its therapeutic actions mainly by scavenging Reactive oxygen species (ROS). RSV is predicted to modulate many cellular targets as well; however, there is little information on its neuroprotective roles. This article describes the effect of RSV treatment on the expression profiles of TNF- α , NFkB and apoptotic factors; Bcl₂ & Bax in cerebral cortex and cerebellum of the CLF rats (induced by administration of 100mg thioacetamide i.p. for 10days) confirming moderate grade HA. A significant increase in mRNA level of TNF- α ($p < 0.001$) in both of the brain regions (cerebral cortex and cerebellum) of HA rats was observed. This was consistent with a similar enhancement in the level of NFkB, a transcription factor for TNF- α synthesis, and thus, suggests induction of TNF- α led inflammation in the brain during persistent HA. A significant decline in Bcl₂ level ($p < 0.001$) with a significantly enhanced level of Bax further suggested a neurodegenerative condition in those brain regions of the HA rats. Moreover, post treatment of those HA rats, with 10mg/Kg b.w. RSV for a week, was found to bring all the factors towards their normal level in both the brain regions. The finding suggest that RSV is able to ameliorate moderate HA induced TNF- α led inflammatory cascade and proapoptotic neurodegenerative condition in the rat brain primarily by modulating expression of the inflammatory and apoptotic factors.

Keywords: hyperammonemia, neuro inflammation, moderate he, tnf- α , apoptosis, cortex, cerebellum

Volume 4 Issue 2 - 2016

Archita Khanna, Surendra K Trigun

Department of Zoology, Banaras Hindu University, India

Correspondence: Surendra K Trigun, Department of Zoology, Institute of Science, Banaras Hindu University, Varanasi-221005, India, Tel 91-542-6702523, Email sktrigun@bhu.ac.in

Received: July 31, 2016 | **Published:** August 26, 2016

Introduction

Ammonia neurotoxicity is considered responsible for development of a serious nervous system disorder called hepatic encephalopathy (HE). This situation arises mainly due to liver dysfunction led hyperammonemia (HA). HE is characterized by the neuropsychiatric manifestations related to motor dysfunction, memory impairment, and deranged sleep-awake cycle.^{1,2} Since HE affects the quality of life of the patients, it is important to develop effective therapeutic strategy against HE. This necessitates understanding the neuro-chemistry of HA pathogenesis.

Since ammonia crosses the blood brain barrier easily, prolonged HA condition, a common situation during chronic liver failure (CLF), is likely to maintain increased ammonia concentration in brain.^{3,4} The information, mainly derived from the cell culture system and from animal models with acute HE, suggest that the increased brain ammonia level mainly drives astrocytes to undergo significant morpho-pathological changes and also over activates glutamate-NMDA receptor (N-Methyl-D-Aspartate Receptor) pathway in the post-synaptic neurons.^{1,5-7} Moreover, NMDAR inhibition could not be translated into prevention of ammonia neurotoxicity, mainly because threshold glutamate-NMDAR activation is necessary for normal neurological functions including higher order brain functions and normal synaptic activity.

During recent past, "TNF theory" (Tumor necrosis factor) of HE pathogenesis could draw much attention due to a close association between the level of TNF- α and ammonia in the brain of CLF patients

with HE.⁸ Later studies have also emphasized significant involvement of TNF- α and other cytokines in HE to the extent that these cytokines could be used as markers for encephalopathy grading.^{4,9-13}

It is now evident that the microglia and astroglia cells in brain synthesize TNF- α to regulate higher order brain functions and astrocyte induced synaptic strengthening.¹⁴⁻¹⁷ However, when produced in higher amounts during neuropathology, they are known to potentiate glutamate induced cytotoxicity by inhibiting glutamate transport to the glial cells from the synaptic cleft.^{12,16-19} Even development of MHE has been described to be dependent more on enhanced inflammatory markers than the severity of CLF or HA in the CLF patients.²⁰ Moreover, some findings from in vitro studies suggest that resveratrol treatment could prevent ammonia toxicity in astroglial cells by normalizing ROS/RNS level.^{12,21,22} Thus, hinting towards a significant role of resveratrol in ameliorating ammonia toxicity. However, the mechanism by which resveratrol does so is unclear.

Resveratrol (3, 3', 4'-trihydroxy-Trans stilbene), a natural polyphenol, is found in a number of dietary sources such as grapes, berries, and red wine.²³⁻²⁵ It acts as an effective cardioprotector, anti-aging, and anti-carcinogenic agent in vivo.^{12,26-28} Some reports also suggest that this natural polyphenol exerts neuroprotection during certain neurodegenerative challenges^{12,26-28} mainly due to its ROS scavenging properties and/or by activating some antioxidant enzymes. Some recent observations, mainly derived from in vitro studies, suggest that ammonia induced impaired glutamatergic communication in astroglial cells could be recovered with resveratrol treatment.²⁹ Since astroglia cells are the main source of inflammatory

cytokines, it is argued that resveratrol could also prevent inflammatory mechanisms implicated in impaired glutamate transport during ammonia neurotoxicity. And if it does so, then this polyphenol must be able to down regulate apoptotic mechanism which otherwise becomes upregulated during such challenges.³⁰⁻³²

Thus, to investigate whether resveratrol is able to modulate inflammation-apoptosis pathway in the HA brain, the expression profiles of TNF- α (a proinflammatory marker), NF κ B (a transcriptional regulator of inflammatory cytokines) and intrinsic apoptotic factors (Bcl₂/Bax) were examined in the cortex and cerebellum (the two susceptible brain regions) of the CLF rats with moderate grade HA.

Material and methods

Chemicals

All chemicals used were of analytical grade purchased from E-Merck and Sisco research Laboratory, Mumbai (India) except N,N'-methylenebisacrylamide, Acrylamide, N,N,N',N'-tetramethylethylenediamine (TEMED), phenylmethylsulfonyl fluoride (PMSF), Ponceau S and bromophenol blue, which were purchased from Sigma-Aldrich, USA. Primary antibodies used were supplied from the following companies: rabbit polyclonal Bcl₂ and Bax from Cell Signaling Technology and rabbit monoclonal β -actin from Sigma Aldrich. Goat anti-Rabbit horseradish peroxidase (HRP) conjugated secondary antibodies were supplied from Genei. ECL western blotting detection kit was procured from Thermo Scientific. The trans-Resveratrol was supplied from the Cayman chemicals company.

The following molecular biology kits and reagents were used: TRI reagent and DEPC (Sigma Aldrich, Revert Aid first strand cDNA synthesis kit (Thermo Scientific, Lithuania), TURBO DNA-free™ (Ambion RNA by Life Technologies™, USA) Maxima SYBR Green/ ROX qPCR master mix 2X (Thermo Scientific). The primers used in the present study were synthesized by Eurofins Genomics India Pvt. Ltd. The rat gene specific primers used were: Bax (Forward 5'- GATCATGAAGACAGGGGCCTT-3'; Reverse 5'-CCTGGATGAAACCCTGTAGCA-3'); Bcl₂ (Forward 5'-ACACCAGAATCAAGTGTTCGT-3'; Reverse 5'-TGCATTCTTGGATGAAGGGGT-3'); TNF- α (Forward 5'TCATCAGTTCCATGGCCAGAC-3'; Reverse 5'-GGTTTGCTACGACGTGGGCTA-3'); NF κ B (Forward 5'- GCGTTTCCGTTACAAGTGCAGG-3'; Reverse 5'-CCCCAGGAATACTGCCTGCAGAG-3'). The primers were designed using the Fast PCR 6.1 software.

Animals

The adult male albino rats weighing 130-160g, used in this study, were fed the recommended diet and maintained on 12:12h light: dark cycle, in an animal house, as per recommendations from the institutional animal ethical committee (IAEC) of Banaras Hindu University, for the care and use of the laboratory animals.

Development of chronic liver failure (CLF)/HA and treatment schedule

The CLF/HA model of neuroinflammation in adult male albino rats was developed by the administration of thioacetamide (TAA) as standardized in our lab.³³ The rats were divided into three groups with six rats in each. Group A: control (C), administered with 0.9% NaCl i.p, once daily for 10days; Group B: Hyperammonemia (CLF/HA), administered with 100mg/Kgbw TAA (prepared in 0.9 % NaCl) i.p

once daily for 10days; Group C: CLF/HA + Resveratrol (CLF/HA + RSV) administered with 10mg/Kgbw Resveratrol i.p once daily, dissolved in 1% DMSO, starting from 8th day onwards till 14th day. The Resveratrol was administered 4h after the TAA treatment. The selected dose of Resveratrol was able to recover TAA induced neurobehavioural deficit in the rats. After 24h of the last dose given, all the rats were sacrificed on 15th day. To confirm the hyperammonemic condition, ammonia assay was performed in the pooled serum of all the six rats from control and HA groups. The cortex and cerebellum was dissected out and stored at -80°C for further studies.

Preparation of cortex and cerebellum extracts

Cerebral cortex and cerebellum extracts were prepared in 0.02 M Tris-Cl (pH 7.4) containing protease inhibitors. Extracts were centrifuged at 35,000 g for 45min at 4°C. The supernatant collected was used for the ammonia assay. Protein content was determined by the method of Lowry et al.,³⁴

Mitochondria-free tissue extract was prepared in an extraction medium consisting of 400 mM sucrose, 0.2mM benzamidine, 1mM EDTA, 0.1mM phenylmethylsulfonyl fluoride (PMSF) (pH 7.4), and 0.02% heparin. Initially, the extracts were centrifuged at 12,000 $\times$ g for 15min and finally at 19,000 $\times$ g for 40min at 4°C to obtain cytosolic fractions. The protein content in the extract was estimated by Lowry method as mentioned earlier.

Ammonia assay: The ammonia assay was performed using a kit supplied by Sigma-Aldrich, USA. The serum as well as cortex and cerebellum extract were deproteinized in 1/5volume of ice-cold 100g/L Trichloroacetic acid and kept on ice for 15min. After centrifugation at 15,000Xg for 15min at 4°C, the supernatants were neutralized with 2M KHCO₃, centrifuged and used for ammonia assay. The quantification of ammonia involved measuring the rate of conversion of α -ketoglutarate to glutamate catalyzed by glutamate dehydrogenase in the presence of ammonia. The decrease in O.D. at 340nm was recorded.

The 1mL reaction mixture, contained 50 μ L samples, 3.4mM α -ketoglutarate and 0.23mM reduced NADPH in water. The mixture was incubated for 5min at 18-35°C and reaction was started by the addition of suitably diluted GDH. Initial and final (at 5min) O.D. at 340nm was used to calculate concentration of ammonia in mM.

Western blotting: Briefly, 100 μ g of cytosolic protein was loaded in each lane and subjected to 10% denaturing polyacrylamide gel electrophoresis at constant voltage of 100volts for 2hours followed by transferring the protein bands on nitrocellulose membrane over night at 50mA at 4°C. Efficiency of Protein transfer was checked by Ponceau S staining. The membrane was then placed in a blocking solution (5% non fat dried milk in 1X PBS) for 2h. The membrane was processed for immunodetection of Bcl₂ antibody (1:1000) and Bax antibody (1:500). HRP-conjugated secondary antibody was used for final detection of the respective proteins using ECL western blotting detection kit. For loading control, monoclonal anti- β -actin peroxidase antibody (1:10,000) was used. The normalized densitometric values of each protein vs β -actin, was recorded using gel densitometry software Alpha Imager 2200.

RNA isolation and cDNA synthesis: Total RNA from cortex and cerebellar extracts was isolated using TRI reagent following the manufacture's protocol. Briefly, the homogenate prepared in 1mL TRI reagent was centrifuged at 12,000g for 10min at 4°C and supernatant was collected. Chloroform was added (0.2mL) to the supernatant and vortexed. After 3min, the tube was centrifuged at 12,000 Xg for

10min, at 4°C. The centrifugation separated the mixture into 3 phases, out of which the colorless upper aqueous phase containing RNA was collected and 0.5mL isopropanol was added to it. After 5min, the tube was again centrifuged at 12,000X g for 10min, at 4°C. The RNA precipitated out in the form of a pellet at bottom of the tube and was washed in 1 mL 75% ethanol. After air drying the RNA pellet, it was dissolved in 60µL DEPC treated water by repeated pipetting. The dissolved RNA pellet was subjected to DNase treatment (DNA free-Ambion) to remove any sort of DNA contamination. The total RNA yield was determined and quality assessed by agarose formaldehyde RNA gel electrophoresis.

The RNA samples with an A_{260}/A_{280} ratio of 1.8-2.0 were used in the cDNA synthesis reaction. RevertAid first strand cDNA synthesis kit was used for cDNA synthesis. The reaction mixture consisted of 4µg RNA, 1 µL random hexamer primer, made up to 12µL with DEPC treated water. The mixture was gently mixed, centrifuged briefly and the following components were added; 4µL of 5X reaction buffer, 1µL of RiboLock™ Ribonuclease Inhibitor (20U/µL), 2µL of 10mM dNTP mix, 1µL of reverse transcriptase to make a final volume of 20µL. The components were mixed gently and incubated for 5min at 25°C followed by cDNA strand synthesis for 60min at 42°C. The reaction was terminated by heating at 72°C for 5min. The cDNA was stored at -80°C.

Real time (qPCR): The real time qPCR was done with an ABI Prism 7500 Sequence Detection System (PE Applied Biosystems, CA, USA). The PCR reaction mixture of (20µL) consisted of 1µL of sample cDNA (diluted 1:10), 1µL of 10 pM of forward and reverse primers and 10µL of 2X Thermo Scientific SYBR Green/ROX qPCR Master Mix and brought to final volume with RNase free water.

PCR condition was: 95°C for 30sec, followed by 40 cycles at 95°C for 5 sec, and 60°C for 20sec. Real-time PCR data was analyzed using the $\Delta\Delta C_t$ method, normalizing the C_t values of the indicated gene to the C_t values of GAPDH relative to a control sample.

Statistical analysis: The statistical analysis of experimental data was done by applying Student's t test, expressed as mean $\pm$ SD and $P < 0.05$ was taken as the level of significance between the control and experimental sets.

Results

Development of chronic liver failure/hyperammonemia

The HA condition due to TAA induced CLF was evident from a ~2 x increase ($P < 0.01$) in serum ammonia level of CLF/HA rats (Figure 1a). Also, as depicted in Figure 1b&c, as compared to the control group rats, the level of ammonia was significantly increased ($P < 0.05$) in the cortex and in the cerebellum of CLF/HA rats. These findings confirm the induction of chronic liver failure led HA condition, both in the blood and in the susceptible brain regions.

Effect of resveratrol on inflammatory factors: TNF- α and NF- κ B

Development of neuroinflammation is considered an important marker for HA in the brain. Neuroinflammation was monitored by measuring the transcript level of neuroinflammatory markers TNF- α and NF- κ B. According to Figure 2a&b, as compared to the control group rats, a significant increase ($P < 0.01$) in the TNF- α transcript in cortex and cerebellum of CLF/ HA rats was observed. A similar pattern was observed for NF- κ B transcript (Figure 3a&3b), which showed significantly enhanced levels ($P < 0.001$) in both the brain

regions of the HA rats. However, the enhanced levels of both the neuroinflammatory markers, TNF- α and NF- κ B, were restored back to their normal levels in the HA rats treated with RSV.

Figure 1 Effect of TAA treatment on ammonia level in serum (a), cortex (b) and cerebellum (c). Values are represented as mean $\pm$ SD where $n = 4$; * $P < 0.05$; ** $P < 0.01$; (control versus CLF), CLF: chronic liver failure.

Figure 2 Expression of TNF- α in cerebral cortex (a) and cerebellum (b) of control, HA and HA+RSV treated rats. Cycle threshold value of TNF- α mRNA level was normalized with the cycle threshold value of Gapdh for each sample to give a relative quantity of the mRNA expression by $2^{-\Delta\Delta C_t}$. Values are represented as mean $\pm$ SD from 4 repeats; Gapdh was taken as the internal control. ** $P < 0.01$; (control versus HA) # $P < 0.05$; ## $P < 0.01$; ### $P < 0.001$; (HA versus HA+RSV), HA: Hyperammonemia, HA+RSV: Hyperammonemia + Resveratrol.

Figure 3 Expression of NF- κ B gene in cerebral cortex (a) and cerebellum (b) of control, HA and HA+RSV rats. Cycle threshold value of NF- κ B RNA level was normalized with the cycle threshold value of Gapdh for each sample to give a relative quantity of the mRNA expression by $2^{-\Delta\Delta C_t}$. Values are represented as mean $\pm$ SD from 4 repeats; Gapdh was taken as the internal control. *** $P < 0.001$; (control versus HA) ### $P < 0.001$; #### $P < 0.001$; (HA versus HA+RSV), HA: Hyperammonemia, HA+RSV: Hyperammonemia + Resveratrol.

Effect of resveratrol on Bcl₂ and Bax expression

Looking at the ratio of Bcl₂/Bax is the most effective way to determine the susceptibility of cells for undergoing apoptosis. According to Figure 4a&c, it is evident that the expression level of Bcl₂ is significantly declined in both, the cortex ($P < 0.001$) and the cerebellum ($P < 0.01$), as compared to that observed in the control

group rats. The expression level of Bax (Figure 5a&c) was also found to be significantly enhanced in the cortex ($P<0.001$) and in the cerebellum ($P<0.01$) of the HA rats, as compared to the control group of rats; resulting in a significant decline in the Bcl₂/Bax ratio presented in Figure 6a for the cortex ($P<0.001$) and in Figure 6b for the cerebellum ($P<0.001$). However, on treatment with RSV, the Bcl₂/Bax ratio was found to be restored to normal levels in both the brain regions of HA rats (Figure 6a&b).

Figure 4

Figure 4 Expression profile of Bcl₂ in cerebral cortex (a) and cerebellum (c) of the HA and RSV treated HA rats. Western blot analysis of Bcl₂ was performed as described in the text of methods. The level of β-actin was taken as the loading control. In panels (b) and (d) normalized densitometric values of Bcl₂/β-actin for cortex and cerebellum respectively, have been presented as mean ± SD from three western blot repeats. ** $P<0.01$; *** $P<0.001$; (control versus HA rats), ### $P<0.01$; #### $P<0.001$ (HA versus HA+ RSV rats). HA: Hyperammonemia, HA+RSV: Hyperammonemia + Resveratrol.

Figure 5 Expression profile of Bax in cerebral cortex (a) and cerebellum (c) of the HA and RSV treated HA rats. Western blot analysis of Bax was performed as described in the text of methods. The level of β-actin was taken as the loading control. In panels (b) and (d) normalized densitometric values of Bax/β-actin for cortex and cerebellum respectively, have been presented as mean ± SD from three western blot repeats. □□ $P<0.01$; *** $P<0.001$; (control versus HA rats), #### $P<0.001$ (HA versus HA+ RSV rats). HA: Hyperammonemia, HA+RSV: Hyperammonemia + Resveratrol.

These findings were further confirmed by conducting the qPCR analysis of Bcl₂ and Bax. According to Figure 7a&b, the transcript level of Bcl₂ was significantly decreased ($P<0.01$) in both the cortex and the cerebellum of the HA rats, as compared to the control group counterparts. This could be normalized in both the brain regions of the HA rats with the resveratrol treatment. Similarly, the transcript level of

Bax, as shown in Figure 8a&b, was significantly increased ($P<0.001$) in both the cortex and the cerebellum of the HA rats, however, due to the RSV treatment, its level recovered to the normal values.

Figure 6 Ratio of Bcl₂/Bax in cerebral cortex (a) and cerebellum (b) presented as mean ± SD from three western blot repeats. *** $P<0.001$; (control versus HA) #### $P<0.001$; (HA versus HA+ RSV rats). HA: Hyperammonemia, HA+RSV: Hyperammonemia + Resveratrol.

Figure 7 Expression of Bcl₂ gene in cerebral cortex (a) and cerebellum (b) of control, HA and HA+RSV rats. Cycle threshold value of Bcl₂ RNA level was normalized with the cycle threshold value of Gapdh for each sample to give a relative quantity of the mRNA expression by 2^{-ddCt}. Values are represented as mean ± SD from 4 repeats; Gapdh was taken as the internal control. ** $P<0.01$; (control versus HA) #### $P<0.001$; (HA versus HA+ RSV rats). HA: Hyperammonemia, HA+RSV: Hyperammonemia + Resveratrol.

Figure 8 Expression of Bax gene in cerebral cortex (a) and cerebellum (b) of control, HA and HA+RSV rats. Cycle threshold value of Bax RNA level was normalized with the cycle threshold value of Gapdh for each sample to give a relative quantity of the mRNA expression by 2^{-ddCt}. Values are represented as mean ± SD from 4 repeats; Gapdh was taken as the internal control. *** $P<0.001$; (control versus HA) #### $P<0.001$; (HA versus HA+ RSV rats). HA: Hyperammonemia, HA+RSV: Hyperammonemia + Resveratrol.

Resveratrol is known to exhibit anti-pathogenic activity against a wide range of diseases mainly by acting as a potent ROS scavenger. Moreover, inflammation and ROS generation are now evident to go hand to hand in a number of pathological manifestations, wherein, TNF-α has been found to play a regulatory role.^{35,36} In case of brain disorders also, neuroinflammation and excitotoxicity have been found to be responsible for triggering neuro-excitotoxic disorders/diseases like; traumatic brain injury, Ischemia, Alzheimers disease, Parkinson's disease, etc.^{17,37-47} Recent experimental evidences also suggest that natural antioxidants may modulate the expression of inflammatory cytokines while protecting diseased cells from oxidative insult. For example, fisetin, a natural product has been demonstrated to decrease

TNF- α mRNA in HCC cells in vivo.⁴⁸ In this article, we examined this hypothesis for RSV, a natural polyphenol, against an inflammation driven excitotoxic brain disorder developed due to persistent HA in CLF rats.

The enhanced levels of ammonia (Figure 1), in blood and in the two brain regions selected (cortex and cerebellum), represent a moderate grade HA condition exposing brain cells with increased ammonia concentration. And as described earlier,^{33,49,50} these rats were considered to be representative of a true CLF led HA model of neuroinflammation/excitotoxicity. Using these HA rats, we examined the expression profiles of TNF- α , NF κ B and apoptotic factors in the cortex and cerebellum. This is because, these two brain regions have been demonstrated to undergo significant neuro-chemical and neuroanatomical changes during ammonia toxicity and HE pathogenesis.^{33,50,51}

The “TNF theory” of HA/HE pathogenesis⁸ emphasizes significant roles of TNF- α and other cytokines in HE up to the extent that these cytokines could be used as markers for encephalopathy grading.^{4,9–13} Obviously, TNF- α could be considered a suitable target for therapeutic intervention, however, with limited information on its responsiveness towards RSV in neuroinflammation and neuroexcitotoxicity. Modulation of the expression of this cytokine by natural product is an upcoming strategy.⁴⁸ In this respect, results in Figure 2 clearly demonstrate a cause and effect relationship between HA & TNF- α mRNA level in cerebral cortex and cerebellum of the HA rats and thereby support the TNF- α theory of HA pathogenesis. However, it was interesting to record recovery of TNF- α expression reaching to normal levels due to the in vivo treatment with RSV. Various strategies have been employed to decrease enhanced TNF- α level by blocking its synthesis and enhancing its degradation as therapeutic mechanisms in excitotoxic neurological disorders.¹⁷ However, examples are limited with regards to RSV and the modulative expression of this cytokine. In this respect, results in Figure 2 are the first of their kind to demonstrate Resveratrol dependent decline in TNF- α mRNA particularly in the cortex and cerebellum of the rats with HA induced neuroinflammation.

The argument gets support from the results in Figure 3 describing HA dependent similar enhancement in NF κ B mRNA level. This is because, NF κ B is known to be a constitutive type transcription factor that is translocated to the nucleus for enhancing expression of the inflammatory cytokines including TNF- α amongst the most prominent ones.^{12,52,53} Moreover, concordant to the RSV dependent decline in TNF- α expression (Figure 3), it was interesting to observe a similar recovery in the level of NF κ B mRNA in both the brain regions of HA rats treated with RSV. Thus, RSV dependent recovery in the expression of TNF- α and NF κ B together, in both the brain regions of HA rats, advocates that RSV could modulate TNF- α level by altering its transcription at gene level. The argument gets support from a recent report describing modulation of TNF- α mRNA by another natural product, fisetin, in case of HCC pathogenesis.⁴⁸

In case of resveratrol, as a neuroprotectant, another issue remaining unsolved is how it prevents possible neurodegeneration under the excitotoxic conditions involving TNF- α as the main culprit. This aspect is equally important because TNF- α is described to inhibit glutamate transport to the glial cells from the synaptic cleft and thereby allowing over activation of glutamate-AMPA and NMDA receptors while decreasing GABA receptor activity of the post synaptic neurons.¹⁷ Such a situation causes enhanced Ca²⁺ entry into the neurons and thereby inducing neuronal death.¹⁷ Such neuronal deaths then must

implicate apoptotic mechanisms to execute safe removal of the dying neurons. The most plausible determinant of intrinsic pathway of apoptosis is the mitochondria damage induced interplay of Bcl₂/Bax ratio under neuropathological conditions.^{30–32} RSV is known to prevent such neurodegeneration by eliminating the ROS insult.^{12,54,55} However, ROS generation and mitochondrial damage also go hand to hand by inducing neurodegeneration in acute HA condition.^{3,50} Thus, it is likely that brain cells, particularly, post synaptic neurons, start undergoing apoptosis by involving classical Bcl₂-Bax pathways.^{30–32}

Moreover, in the present context, the findings in Figure 4,5& 6 clearly demonstrate a significant decline in Bcl₂/Bax ratio attributed mainly to the declining Bcl₂ (an antiapoptotic factor) level with the enhanced Bax (pro-apoptotic factor) level in cerebral cortex and cerebellum of the HA rats and thus suggesting induction of apoptosis in both these brain regions of the HA rats. However, when these HA rats were treated with RSV, the Bcl₂/Bax ratio was found to recover back to normal levels. This further suggests that resveratrol is able to modulate Bcl₂ and Bax expression and thereby, allow the recovery of brain cells towards an anti-apoptotic condition from a HA induced pro-apoptotic microenvironment. This argument gets support from a similar finding on recovery of Bcl₂/Bax ratio in the cerebellum of CLF rats with even mildest HA condition due to the in vivo treatment with Bacopa extract.⁵¹ Indeed, Resveratrol could also diminish memory impairment in the vascular dementia model by attenuating increased expression of Bax and declining Bcl₂ expression.³²

Conclusion

In conclusion, these findings not only provide evidence for a significant neuroprotective role of Resveratrol during moderate grade HA but also demonstrate that it does so by modulating enhanced expressions of TNF α and NF κ B and by reciprocally regulating expression of Bax and Bcl₂ in the cortex and cerebellum of the HA rats. Unravelling such a mechanistic aspect of in vivo neuroprotective action of resveratrol, under a neuro-excitotoxic/neuro-inflammation condition, is likely to open up the research for further examination of this mechanism in other brain disorders arising out of neuroinflammation and excitotoxicity.

Acknowledgements

This work was financially supported by an ICMR project (54/38/CFPGER/2011/NCD-II) to SKT. A. Khanna acknowledges the Lady Tata Memorial Trust- JRF/SRF’ awarded to her during the tenure of this work. The authors are thankful to the DST FIST, CAS COSIST, DST Purse and DBT-BHU ISLS programmes in the Department & Institute of Science for providing instrumentation facilities.

Conflicts of interest

Author declares there are no conflicts of interest.

Funding

None.

References

1. Blei AT, Cordoba J. The Practice Parameters Committee of the American College of Gastroenterology Hepatic Encephalopathy. *Am J of Gastroenterol.* 2001;96(7):1968–1976.
2. Bajaj SJ, Wade JB, Sanyal AJ. Spectrum of neurocognitive impairment in cirrhosis: implication for the assessment of hepatic encephalopathy. *Hepatology.* 2009;50(6):2014–2021.

3. Felipe V. Hyperammonemia. In: A Iaztha, N Banik, SK Ray, Editors. Brain and spinal cord trauma. 3rd (edn), Handbook of Neurochemistry and molecular neurobiology. Berlin: Springer; 2009,p.43–69.
4. Felipe V, Urios A, Montesinos E, et al. Contribution of hyperammonemia and inflammatory factors to cognitive impairment in minimal hepatic encephalopathy *Metabolic Brain Disorder*. 2012;27(1):51–58.
5. Hermenegildo C, Montoliu C, Llansola M, et al. Chronic hyperammonemia impairs the glutamate-nitric oxide-cyclic GMP pathway in cerebellar neurons in culture and in the rat in vivo. *Eur J Neurosci*. 1998;10(10):3201–3209.
6. Hermenegildo C, Monfort P, Felipe V. Activation of N-methyl-D-aspartate receptors in rat brain in vivo following acute ammonia intoxication: characterization by in vivo brain microdialysis. *Hepatology*. 2000;31(3):709–715.
7. Poh Z, Chang PEJ. A Current Review of the Diagnostic and Treatment Strategies of Hepatic Encephalopathy. *International Journal of Hepatology*. 2012;2012:1–10.
8. Odeh M, Sabo E, Srujo I, et al. Relationship between tumor necrosis factor- α and ammonia in patients with hepatic encephalopathy due to chronic liver failure. *Ann Med*. 2005;37(8):603–612.
9. Odeh M. Pathogenesis of hepatic encephalopathy: the tumor necrosis factor- α theory. *Eur J Clin Invest*. 2007;37(4):291–304.
10. Butterworth RF. Neuroinflammation in acute liver failure: Mechanisms and novel therapeutic targets. *Neurochem Int*. 2011;59(6):830–836.
11. Butterworth RF. Hepatic encephalopathy: a central neuroinflammatory disorder? *Hepatology*. 2011;53(4):1372–1376.
12. Bobermin LD, Quincozes-Santos A, Guerra MC, et al. Resveratrol prevents ammonia toxicity in astroglial cells. *PLoS One*. 2012;7(12):e52164.
13. Chastre A, Blanger M, Beauchesne E, et al. Inflammatory cascade driven by tumor necrosis factor- α play a major role in the progression of acute liver failure and its neurological complications. *PLoS One*. 2012;7(11):e49670.
14. Beattie EC, Stellwagen D, Morishita W, et al. Control of synaptic strength by glial TNF α . *Science*. 2002;295(5563):2282–2285.
15. Beste C, Baune BT, Falkenstein M, et al. Variations in the TNF- α gene (TNF- α -308G $\rightarrow$ A) affect attention and action selection mechanisms in a dissociated fashion. *Journal of Neurophysiology*. 2010;104(5):2523–2531.
16. Santello M, Bezzi P, Volterra A. TNF α controls glutamatergic gliotransmission in the hippocampal dentate gyrus. *Neuron*. 2011;69(5):988–1001.
17. Olmos G, Llad J. Tumor Necrosis Factor Alpha: A link between neuroinflammation and excitotoxicity Mediators of *Inflammation*. 2014;2014:1–12.
18. Tanabe K, Matsushima-Nishiwaki R, Yamaguchi S, et al. Mechanisms of tumor necrosis factor- α -induced interleukin-6 synthesis in glioma cells. *J Neuroinflammation*. 2010;7:16.
19. Tanabe K, Kozawa O, Iida H. Midazolam suppresses interleukin-1 β -induced interleukin-6 release from rat glial cells. *Journal of Neuroinflammation*. 2011;8:68.
20. Shawcross DL, Wright G, Olde Damink SW, et al. Role of ammonia and inflammation in minimal hepatic encephalopathy. *Metab Brain Dis*. 2007;22(1):125–138.
21. Quincozes-Santos A, Andreazza AC, Nardin P, et al. Resveratrol attenuates oxidative-induced DNA damage in C6 Glioma cells. *Neurotoxicology*. 2007;28(4):886–891.
22. Quincozes-Santos A, Andreazza AC, Goncalves CA, et al. Actions of redox-active compound resveratrol under hydrogen peroxide insult in C6 astroglial cells. *Toxicology In Vitro*. 2010;24(3):916–920.
23. Pervaiz S. Resveratrol: from grapevines to mammalian biology. *FASEB J*. 2003;17(14):1975–1985.
24. Kulkarni SS, Cant C. The molecular targets of resveratrol. *Biochim Biophys Acta*. 2015;1852(6):1114–1123.
25. Park EJ, Pezzuto JM. The pharmacology of resveratrol in animals and humans. *Biochim Biophys Acta*. 2015;1852(6):1071–1113.
26. Delmas D, Jannin B, Latruffe N. Resveratrol: preventing properties against vascular alterations and ageing. *Mol Nutr Food Res*. 2005;49(5):377–395.
27. Baur JA, Sinclair DA. Therapeutic potential of resveratrol: the in vivo evidence. *Nat Rev Drug Discov*. 2006;5(6):493–506.
28. Vang O Ahmad N, Baile CA, Baur JA, et al. What is new for an old molecule? Systematic review and recommendations on the use of resveratrol. *PLoS One*. 2011;6(6):e19881.
29. Bobermin LD, Hansel G, Scherer EB, et al. Ammonia impairs glutamatergic communication in astroglial cells: protective role of resveratrol. *Toxicol in Vitro*. 2015;29(8):2022–2029.
30. Clementi ME, Pezzotti M, Orsini F, et al. Alzheimer’s amyloid beta-peptide (1–42) induces cell death in human neuroblastoma via bax/bcl-2 ratio increase: an intriguing role for methionine 35. *Biochem Biophys Res Commun*. 2006;342(1):206–213.
31. Sun ZK, Yang HQ, Wang ZQ, et al. Erythropoietin prevents PC12 cells from beta-amyloid-induced apoptosis via PI3K/Akt pathway. *Translational Neurodegeneration*. 2012;1:1–7.
32. Sun ZK, Ma XR, Jia YJ, et al. Effects of resveratrol on apoptosis in a rat model of vascular dementia. *Exp Ther Med*. 2014;7(4):843–848.
33. Singh S, Trigun SK. Activation of neuronal nitric oxide synthase in cerebellum of chronic hepatic encephalopathy rats is associated with up-regulation of NADPH-producing pathway. *Cerebellum*. 2010;9(3):384–397.
34. Lowry OH, Rosebrough NJ, Farr AL, et al. Protein measurement with the Folin phenol reagent. *J Biol Chem*. 1951;193(1):265–275.
35. Afonso V, Santos G, Collin P, et al. Tumor necrosis factor- α downregulates human Cu/Zn superoxide dismutase 1 promoter via JNK/AP-1 signaling pathway. *Free Radic Biol Med*. 2006;41(5):709–721.
36. Singh KB, Maurya BK, Trigun SK. Activation of oxidative stress and inflammatory factors could account for histopathological progression of aflatoxin-B1 induced hepatocarcinogenesis in rat. *Molecular Cellular Biochemistry*. 2015;401(1–2):185–196.
37. Goodman JC, Robertson CS, Grossman RG, et al. Elevation of tumor necrosis factor in head injury. *J Neuroimmunol*. 1990;30(2–3):213–217.
38. Fillit H, Ding W, Buee L, et al. Elevated circulating tumor necrosis factor levels in Alzheimer’s disease. *Neurosci Lett*. 1991;129(2):318–320.
39. Sharief MK, Hentges R. Association between tumor necrosis factor- α and disease progression in patients with multiple sclerosis. *N Engl J Med*. 1991;325(7):467–472.
40. Boka G, Anglade P, Wallach D, et al. Immunocytochemical analysis of tumor necrosis factor and its receptors in Parkinson’s disease. *Neurosci Lett*. 1994;172(1–2):151–154.
41. Mogi M, Harada M, Riederer P, et al. Tumor necrosis factor- α (TNF- α) increases both in the brain and in the cerebrospinal fluid from parkinsonian patients. *Neurosci Lett*. 1994;165(1–2):208–210.
42. Feuerstein GZ, Liu T, Barone FC. Cytokines, inflammation, and brain injury: role of tumor necrosis factor- α . *Cerebrovasc Brain Metab Rev*. 1994;6(4):341–360.

43. Liu T, Clark RK, McDonnell PC, et al. Tumor necrosis factor- α expression in ischemic neurons. *Stroke*. 1994;25(7):1481–1488.
44. Rieckmann P, Albrecht M, Kitze B, et al. Tumor necrosis factor- α messenger RNA expression in patients with relapsing-remitting multiple sclerosis is associated with disease activity. *Ann Neurol*. 1995;37(1):82–88.
45. Poloni M, Facchetti D, Mai R, et al. Circulating levels of tumour necrosis factor- α and its soluble receptors are increased in the blood of patients with amyotrophic lateral sclerosis. *Neurosci Lett*. 2000;287(3):211–214.
46. Alvarez A, Cacabelos R, Sanpedro C, et al. Serum TNF- α levels are increased and correlate negatively with free IGF-I in Alzheimer disease. *Neurobiol Aging*. 2007;28(4):533–536.
47. Babu GN, Kumar A, Chandra R, et al. Elevated inflammatory markers in a group of amyotrophic lateral sclerosis patients from northern India. *Neurochem Res*. 2008;33(6):1145–1149.
48. Maurya K, Trigun SK. Fisetin Modulates Antioxidant Enzymes and Inflammatory Factors to Inhibit Aflatoxin-B1 Induced Hepatocellular Carcinoma in Rats. *Oxidative Medicine and Cellular Longevity*. 2016;2016(2016):1–9.
49. Felipo V, Butterworth R. Neurobiology of ammonia. *Prog Neurobiol*. 2002;67(4):259–279.
50. Singh S, Trigun SK. Low grade cirrhosis induces cognitive impairment and motor dysfunction in rats: Could be a model for minimal hepatic encephalopathy. *Neurosci Lett*. 2014;559:136–140.
51. Mondal P, Trigun SK. Bacopa monnieri Extract (CDRI-08) Modulates the NMDA Receptor Subunits and nNOS-Apoptosis Axis in Cerebellum of Hepatic Encephalopathy Rats. *Evid Based Complement Alternat Med*. 2015;2015:535013.
52. Baltimore D. NF- κ B is 25. *Nature Immunology*. 2011;12:683–685.
53. Gerlo S, Kooijman R, Beck IM, et al. Cyclic AMP: a selective modulator of NF- κ B action. *Cell Mol Life Sci*. 2011;68(23):3823–3841.
54. Shen HM, Pervaiz S. TNF receptor superfamily-induced cell death: redox-dependent execution. *FASEB J*. 2006;20(10):1589–1598.
55. Hamby ME, Gragnoli AR, Hewett SJ, et al. TGF β 1 and TNF α potentiate nitric oxide production in astrocyte cultures by recruiting distinct subpopulations of cells to express NOS-2. *Neurochem Int*. 2008;52(6):962–971.