The use of herbal medicine in a rural community in Vitoria da Conquista, Bahia, Brazil: an indication for pharmacological studies

Abstract

The present study evaluated ethno medicinal knowledge of people in the rural community of São Sebastião, Vitória da Conquista, Bahia, Brazil. Using semi-structured interviews, 84 randomly chosen individuals informed about the use of traditional medicinal plants in their community, thereby identifying their therapeutic potential, which were categorized according to the ICD-10 (International Statistical Classification of Diseases and Related Health Problems). The collected data were analyzed with using the Informant consensus factor (ICF), the Relative importance (RI) and the scores were calculated by using Fidelity level (FL). Forty-six species traditionally used in treatment of ailments were categorized as diseases: digestive system, respiratory problems, general symptoms/signs, nervous system and circulatory system. *Lippia alba*, *Cymbopogon citratus* and *Foeniculum vulgare* had a RI >1, indicating important value to the community. Highest values of FL were: *Spondias purpurea*, *Bauhinia forficata*, *Punica granatum*, *Matricaria chamomilla*, *Stryphnodendron adstringens*, *Foeniculum vulgare*, *Mentha piperita*, *Amburana cearensis* and *Dysphania ambrosioides*. *S. purpurea*, to treat hypertension, and *Amburana cearensis* (indigestion) deserve in depth pharmacological and toxicological studies so that they might be included in the National Program of Medicinal Plants and Phytotherapies of the Brazilian Sistema Único de Saúde and safely prescribed by regional health teams, lowering costs for local health system.

Keywords: ethnobotanical study, northeast brazil, traditional medicine, medicinal plants, *spondias purpurea*

Volume 7 Issue 1 - 2017

Gabriele Marisco,¹ Thalana SS Silva,¹
 Renata Assun,^{1,2} Martin Brende,^{1,2} Cristina
 Pungartnik²

¹Departamento Ciências Naturais, Universidade Estadual do Sudoeste da Bahia, Brazil

²Departamento Ciências Biológicas, Universidade Estadual de Santa Cruz, Brazil

Correspondence: Cristina Pungartnik, Departamento Ciências Biológicas, Universidade Estadual de Santa Cruz, Laboratório de Biologia dos Fungos, Rodovia Ilhéus-Itabuna, Km-16, Salobrinho, Ilhéus, Bahia, Brazil, Tel 557336805438, Email cpungartnik@yahoo.com.br

Received: February 20, 2017 | **Published:** May 17, 2017

Abbreviations: ICF, informant consensus factor; RI, relative importance; FL, fidelity level; UESB, universidade estadual do sudoeste da bahia, HUESBVC, herbarium at uesb, vitoria da conquista

Introduction

Brazil is a continental country with six different biomes, Amazon, Atlantic Forest, Cerrado, Caatinga, Pantanal and Pampa,¹ of which 3 are represented in the State of Bahia (Atlantic Forest, Caatinga, Cerrado).^{2,3} Traditional knowledge of medicinal plants used by indigenous populations of rural areas of Bahia has not received enough attention so far, although in northeastern Brazil there are ethnobotanical studies recently published.⁴⁻⁸ Vitoria da Conquista, Bahia was formerly (18th Century) inhabited by the Indian tribes Mongoiós (subgroup Camacãs), Ymborés (or Aimorés) and to a lesser extent by the Pataxós. Their villages were scattered over a wide range, known as the Sertão da Ressaca. Today, the original vegetation cover in Vitória da Conquista does not exceed 10%, consisting of Caatinga and Atlantic Forest biomes.⁹ This vegetation type is of fundamental importance for maintaining the quality of regional watersheds, such as Rio Mole, Rio Pardo and Rio Caculé (rivers) that benefit a large population of southwestern Bahia.¹⁰ The rural communities, mainly extinguished Brazilian natives (indigenous communities) own diverse knowledge about plants and caring for the environment. Thus, the need of research on communities and how they deal with plants so that their descendants acquire their traditional knowledge.¹¹

Use of medicinal plants has grown considerably in the second half of the 20th century, from many parts of the world. In the developing countries 65-80% of the population depends exclusively on the medicinal plants for the basic healthcare,¹² including Northeastern

Brazil.⁵ Brazil was considered the seventh largest market, and from 2006 to 2010 grew by 14% and it is also estimated that by 2015 Brazil will be the sixth largest market in this sector.¹³ Important policies were implemented in the last decade to encourage research and development related to the use of medicinal plants and herbal remedies which may be provided with quality, safety and efficacy to the population, prioritizing the country's biodiversity.¹⁴ Considering the actual worldwide popular use of medicinal plants, the *Sistema Único de Saúde* (SUS, Brazilian Health System), via the Brazilian Ministry of Health, has introduced the National Policy on Integrative and Complementary Practices and the National Policy on Medicinal Plants in 2006^{13,15} which lists some indications for the use of medicinal plants, such as *Aloe vera* L. (against psoriasis and burns), *Salix alba* L. (against backache) and *Rhamnus purshiana* DC. (to treat constipation). The increased prescription of medicinal plants via physicians of SUS has not only raised their consume in health care but has also helped to educate about possible dangers of their indiscriminate use. As prescription of local herbal medicines by qualified professionals and accurate information have increased and this will help to prevent medical application of unsafe and ineffective plants.^{16,17}

Different tools to quantify ethno botanical data are applied in complementing studies of plant usage in folk medicine. Thus, collection of quantitative data could support the conservation of certain plant species and knowledge of their popular medical use, mainly by informing about species with several therapeutic indications.^{18,19} Some of these tools have been successful in identifying the pharmacological potential of medicinal plants, thus enabling their effective application in herbal medicine.²⁰⁻²² The present study evaluated the ethno medicinal knowledge of the people in the community of São Sebastião, Vitória da Conquista, Bahia, using ethno botanical tools in order to

provide baseline information for future ethno pharmacological and phytochemical studies.

Materials and methods

Study area

The study area is located in the state of Bahia and the survey was conducted in the village of São Sebastião, which belongs to the city of Vitória da Conquista (14° 47' S and 40° 39' W). São Sebastião is a small rural community of 2,730 inhabitants (1,325 men and 1,405 women, average monthly income of US\$ 280), to whom the use of medicinal plants may not only be a tradition but also an economic necessity; indeed, this area still hosts many Brazilian native plants, which are worth thorough study and protection, as elsewhere in Brazil.^{23,24} Figure 1 shows the detailed location map of the study area.

Figure 1 Location of the study area, village of São Sebastião, Vitória da Conquista, Bahia, Brazil.

Ethnobotanical survey

Data were collected during 2012 to 2014. Participants signed the Term of Free and Clarified Assent (demanded by the National Health Council through the Ethics and Research Committee, Resolution 196/96 of the CNS/MS). The study was approved by the Committee of Ethics in Research with Human Beings (CEP) of the State University of Southwestern Bahia (Universidade Estadual do Sudoeste da Bahia-UESB) (N° 0134.0.454.000.11). Data of the semi-structured questionnaires informed on the respondents profile and knowledge about plants used in medicinal therapy, how the knowledge was acquired, plants frequently used by that community and their common therapeutic indications. The cited regional plant species were collected, identified and integrated into the collection of the herbarium at UESB, Vitoria da Conquista (HUESBVC).

Data analysis

The citations for therapeutic purposes were classified according to ICD-10 (International Statistical Classification of Diseases and Related Health Problems) established by World Health Organization.²⁵ The Informant Consensus Factor (ICF) was calculated for each ailment category to identify the agreements of the informants on the reported cures for the group of diseases.²⁶ The Relative Importance Value (RI), a measure of the diversity of medicinal applications, was calculated for each medicinal plant cited by informants.²⁷ The highest possible RI value (2.0) indicates the species with the highest diversity of medicinal use. Scores were calculated Fidelity Level (FL).²⁸

Results and discussion

Socio-demographic information

Eighty-four (84) community residents were interviewed (83.3% women aged between 18 and 80 years, half declaring themselves “housewives”) representing 84 families of whom 96% used medicinal plants to treat different diseases. In regard to the level of schooling, 10.8% were illiterate, 63.1% of respondents did not complete elementary school, 11.8% did not complete secondary school and 14.3% completed secondary school.

Acquisition, parts used and mode of preparation of medicinal plants

Table 1 reports 46 medicinal plants belonging to 30 families. This survey showed that some 80% of the species with medicinal properties were cultivated in orchards or gardens in backyards of the houses; 15% are considered wild and extracted from the surrounding environments and 5% of the plants were bought at popular markets or drugstores, demonstrating the importance of cultivation as the main source of medicinal plants. Several authors in similar ethnopharmacological studies have also reported that the majority of respondents acquired plants grown in their own backyard cultivars.^{20,24} The leaves (78.2%) were the most frequently used plant part for the treatment of diseases, followed by root (8.2%) and seed (6.4%) (Figure 2), which corroborated similar results found in other ethnobotanical studies.^{29,30} In many parts of Brazil the use of teas is a common practice. The main preparation methods of the plants cited were by decoction (69.3%), boiling plant material in water, and infusion (21.2%), which is worrying since infusion is used to prepare all tender parts of medicinal plants such as leaves, buttons and flowers, rich in volatile components, delicate aromas and active principles, degraded by the action of water and heat.³¹

Ailments treated by medicinal plants

Respondents were asked to associate a medicinal plant with possible health care indications, according to WHO.^{24,32} The highest number of citations (106) was for digestive system, while the highest ICF value was obtained for nervous problems (0.85) and digestive system (0.84). Values for nine common diseases categories are given in Table 2. Most cited were those categories related to the digestive system (32.8%), followed by respiratory diseases (17%), general symptoms/signs of inflammation (14.5%), nervous system (13.3%) and circulatory system diseases (12.6%). Similar results were already found in other studies^{20,24,33,34} indicating that the residents of São Sebastião use medicinal plants as the first line of treatment for most common diseases.

Use and efficacy of the medicinal plants

Species with high value of RI (>1) are used to treat a variety of diseases and are therefore considered important to a community.²⁷ *Lippia alba* with RI = 2 was the most versatile species in relation to its use and was indicated for cures of up to 6 body systems, followed by *Cymbopogon citratus* and *Foeniculum vulgare* (Table 3). All plants with RI >1 through this methodology assume that a plant is important when it is most versatile; they are measures of folk knowledge and interpretations of their use must be carefully made.⁴ The data indicated that some plants have more diversified medicinal uses or applications than others.³⁵ Among all plants, *Lippia alba* with RI=2 was the most versatile species in relation to its use and has often been reported in ethnobotanical studies³⁶ and pharmacological use, for example, vasorelaxant effect,³⁷ action as a tranquilizer³⁸ and effective

at preventing gastric ulceration.³⁹ However, when the objective is the search for new plant-derived drugs, the concordance of the answers as to the medicinal use of a particular plant is extremely important. Plants indicated for many different applications would have less credibility

compared to those with a higher fidelity of use.³³ Plants with relatively high level of concordance, i.e., with multiple informants agreeing on their therapeutic value, suggest a real efficacy in disease treatment.²⁸

Table 1 List of traditional medicinal plants investigated in this study with their related information

Voucher	Scientific name	Family	Popular name brazil	Part plant indicated	preparation
6453	<i>Dysphania ambrosioides</i> (L) Mosyakin & Clemants	Amaranthaceae	Mastruz	Root, leaves	Juice, infusion, decoction, maceration (cutaneous)
8010	<i>Allium sativum</i> L	Amaryllidaceae	Alho	Bulb	Decoction, juice, infusion
6452	<i>Spondias purpurea</i> L	Anacardiaceae	Seriguela	Leaves	Decoction
6377	<i>Foeniculum vulgare</i> Mill.	Apiaceae	Erva doce	Leaves, seed	Infusion, decoction
8008	<i>Lepidium virginicum</i> L.	Brassicaceae	Agrião	Leaves	Syrup
8007	<i>Ananas comosus</i> (L) Merr.	Bromeliaceae	Abacaxi	Fruit	Decoction
8018	<i>Matricaria chamomilla</i> L.	Compositae	Camomila	Flower	Infusion
7994	<i>Bidens pilosa</i> L.	Compositae	Carrapicho de agulha	Root	Infusion, decoction
8012	<i>Solidago microglossa</i> DC.?	Compositae	Arnica	Leaves	Infusion
7999	<i>Baccharis trimera</i> (Less.) DC.	Compositae	Carqueja	Leaves	Decoction
8019	<i>Costus</i> sp.	Costaceae	Cana água	Leaves	Decoction
8040	<i>Cucumis sativus</i> L.	Cucurbitaceae	Pepino	Fruit	Juice
7989	<i>Sechium edule</i> (Jacq.) Sw.	Cucurbitaceae	Chuchu	Leaves	Decoction
8030	<i>Eleocharis</i> sp.	Cyperaceae	Junça	Root	Decoction
8037	<i>Aleurites moluccanus</i> (L) Willd	Euphorbiaceae	Nogueira	Seed	Maceration
8045	-	Lamiaceae	Tioioiô	Leaves	Decoction
6459	<i>Plectranthus barbatus</i> Andrews	Lamiaceae	Boldo	Leaves	Infusion, decoction, juice
8002	<i>Mentha spicata</i> L.	Lamiaceae	Hortelãzinho	Leaves	Syrup, infusion, decoction
8028	<i>Plectranthus amboinicus</i> (Lour.) Spreng.	Lamiaceae	Hortelã grosso	Stem, leaves, root	Syrup, decoction
8005	<i>Ocimum basilicum</i> L.	Lamiaceae	Alfavaca	Leaves	Syrup, infusion
8009	<i>Rosmarinus officinalis</i> L.	Lamiaceae	Alecrim	Leaves	Decoction, syrup
8041	<i>Mentha</i> sp.	Lamiaceae	Poejo	Leaves	Infusion
8006	<i>Persea americana</i> Mill.	Lauraceae	Abacate	Leaves	Juice
8043	<i>Senna</i> sp.	Leguminosae	Sene	Leaves	Infusion
8014	<i>Stryphnodendron adstringens</i> (Mart.) Coville	Leguminosae	Barbatimão	Root, leaves, bark	Infusion, decoction
8046	<i>Amburana cearensis</i> (Allemão) A.C.Sm	Leguminosae	Umburana macho, Umburana, amburana, amburana-de- cheiro, imburana, cerejeira-rajada.	Seed, bark	Infusion, decoction
8038	<i>Bauhinia forficata</i> Link	Leguminosae	Pata de vaca	Leaves	Infusion, decoction
6460	<i>Punica granatum</i> L.	Lythraceae	Romã	Leaves	Infusion, decoction
6461	<i>Malpighia emarginata</i> DC.	Malpighiaceae	Acerola	Leaves	Juice
8022	<i>Dorstenia</i> sp.	Moraceae	Contra- erva	Root	Decoction
8036	<i>Myristica fragrans</i> Houtt	Myristicaceae	Nós moscada	Seed, fruit	Juice, decoction
7996	<i>Eugenia uniflora</i> L.	Myrtaceae	Pitanga	Leaves	Decoction
8024	<i>Eucalyptus</i> sp.	Myrtaceae	Eucalpto	Leaves	Infusion, decoction
8021	<i>Averrhoa carambola</i> L.	Oxalidaceae	Carambola	Leaves	Infusion
8033	<i>Passiflora edulis</i> Sims	Passifloraceae	Maracujá	Leaves	Infusion, decoction
8042	<i>Phyllanthus heteradenius</i> Müll. Arg.	Phyllanthaceae	Quebra-pedra	Root, stem, leaves	Decoction
8044	<i>Plantago major</i> L.	Plantaginaceae	Transagem	Leaves, root	Infusion, decoction
6384	<i>Cymbopogon citratus</i> (DC.) Stapf.	Poaceae	Capim santo	Leaves, root	Infusion, decoction, inhalation, juice
8029	<i>Genipa americana</i> L.	Rubiaceae	Genipapo	Fruit	Juice
8032	<i>Citrus limon</i> (L) Osbeck	Rutaceae	Limão	Leaves, fruit	Decoction, juice
8031	<i>Citrus sinensis</i> (L) Osbeck	Rutaceae	Laranja	Leaves	Decoction
8013	<i>Ruta graveolens</i> L.	Rutaceae	Arruda	Leaves	Maceration
8017	<i>Solanum melongena</i> L.	Solanaceae	Beringela	Fruit	Juice
7995	<i>Lippia alba</i> (Mill.) N.E. Brown	Verbenaceae	Erva-cidreira	Leaves, seed, root	Infusion, decoction
8016	<i>Aloe vera</i> L.Burm.f.	Xanthorrhoeaceae	Babosa	Leaves	Scraping, topic contact
8026	<i>Zingiber officinale</i> Roscoe	Zingiberaceae	Gengibre	Root	Decoction

Table 2 Informant consensus factor (ICF) by category of diseases. (ICF is the number of use citations in each category minus the number of species used, divided by the number of use citations in each category minus one).

Category	Species	Citations	ICF
Nervous system	7	43	0.85
Digestive system	17	106	0.84
Respiratory diseases	17	55	0.70
Circulatory system diseases	13	41	0.70
General symptoms/signs ^a	17	47	0.65
Endocrine disorders	4	8	0.57
Genitourinary diseases	10	14	0.30
Skin ailments	3	5	0.5
Neoplasias	1	1	0
Infectious diseases	1	1	0
Diseases of the musculoskeletal system	1	1	0
Diseases of the ear	1	1	0

^asymptoms and signs such as stomachache, inflammation, infection, fever, headache, malaise

Figure 2 Plant parts used for treating diseases/disorders.

In order to find therapeutically potent plants for chemical screening, the Fidelity Level (FL) values of 16 medicinal plants (Table 4) were cited by three or more informants. Two plants were obtained: *Spondias purpurea* and *Bauhinia forficata*, with the highest (100%) FL values for hypertension and diabetes respectively. The

species that gave the highest fidelity level values are considered more promising candidate plants for further pharmacological investigations and deserve priority attention.³⁵ Most plants mentioned in the study, for example, *Bauhinia forficata*, *Punica granatum*, *Matricaria chamomilla* and *Stryphnodendron adstringens* showed high FL, as expected, since literature has plenty of ethnopharmacological studies related to them. However, the plants which mentioned high FL, *Spondias purpurea* and *Amburana cearensis*, are suggested for further study. *Spondias purpurea* was cited for only one treatment (hypertension, 100%), it is exotic (not originally native from Brazil), though it is widely used in northeastern Brazil. *Amburana cearensis* (indigestion, 55.5%) is native from Brazil and has few published information for a therapeutic target, furthermore it has been reported in other studies ethnobotanical.^{12,32} Therefore, these species are promising candidates for pharmacological studies.

Comparison of outstanding medicinal plants with the literature and other studies

Comparison of the pharmacological literature published in different countries with the present ethnobotanical data showed that many of the plants have earlier been reported to have activities against specific diseases. Examples include *Bauhinia forficata*, used to treat diabetes, and *Allium sativum*, used to treat hypertension problems. Such observations hinted at the fidelity of the reports by the residents of São Sebastião, confirming that this knowledge has been preserved as in other communities. Whereas other medicinal plants are used differently, when compared to other traditional practices, for example *Amburana cearensis* was used against indigestion, but it was mentioned in other studies for antibacterial activity, to treat rheumatism, cold, sinusitis⁴⁰ and as anti-inflammatory⁴¹ (Table 5). There is also a species used in this study, *Spondias purpurea*, but not mentioned in the literature for its medicine purpose.

Comparing this study to other ethnobotanical studies on traditional medicinal plants in Northeastern Brazil, although with areas of study in different proportions and methods of research, it was possible to observe that considerable similarity exists with respect to categories of cited diseases (problems of the respiratory system, digestive system, circulatory system).^{4,5} According to this study, *Amburana cearensis* and *Bauhinia* sp are the most common medicinal plants in Northeastern Brazil cited in ethnobotanical surveys.^{5,6,12,23,18}

Table 3 Relative importance (RI) values for medicinal plants used against specific ailments.

Species	Relative importance	Origin*	categories
<i>Lippia alba</i>	2.0	Native	Diseases of the nervous system, genitourinary system, circulatory system, diseases, respiratory system, symptoms and signs involving the digestive system and the abdomen, general signs and symptoms
<i>Cymbopogon citratus</i>	1.87	Exotic	Diseases related to the respiratory system, circulatory and nervous, symptoms and signs involving the digestive system and the abdomen, diseases of the genitourinary system, general signs and symptoms
<i>Foeniculum vulgare</i>	1.43	Exotic	Diseases, symptoms and signs involving the digestive system and abdomen, diseases of the nervous system, diseases of the circulatory system, categories general signs and symptoms, , diseases of the genitourinary system
<i>Aloevera</i>	0.59	Exotic	Neoplasias, diseases of the skin, symptoms and signs involving the digestive system and the abdomen, categories general signs and symptoms
<i>Mentha spicata</i>	0.83	Exotic	Diseases digestive system and abdomen, diseases respiratory system, diseases of the genitourinary system
<i>Dysphania ambrosioides</i>	0.80	Native	Diseases, symptoms and signs involving the digestive system and abdomen, Diseases related to the respiratory system, general symptoms and signs, diseases of the skin
<i>Myristica fragrans</i>	0.59	Native	Symptoms and signs involving the digestive system and the abdomen, general signs and symptoms, diseases of the circulatory system
<i>Punica granatum</i>	0.57	Exotic	Categories respiratory, digestive and genitourinary systems
<i>Passiflora edulis</i>	0.54	Native	Categories nervous system, circulatory, endocrine, nutritional and metabolic diseases

Table Continued...

Species	Relative importance	Origin*	categories
<i>Amburana cearensis</i>	0.51	Native	Diseases digestive system and abdomen, general signs and symptoms
<i>Plectranthus barbatus</i>	0.45	Exotic	Diseases digestive system and abdomen, general signs and symptoms
<i>Plectranthus amboinicus</i>	0.43	Exotic	Diseases respiratory system, infectious diseases
<i>Alliumsativum</i>	0.39	Exotic	Respiratory diseases, diseases of the circulatory system
<i>Stryphnodendron adstringens</i>	0.37	Native	Categories general signs and symptoms, genitourinary, and skin diseases
<i>Citrus limon</i>	0.37	Exotic	Respiratory diseases, diseases of the circulatory system
<i>Bauhinia forficata</i>	0.24	Native	Diseases of the circulatory system, endocrine, nutritional and metabolic diseases
<i>Matricaria chamomilla</i>	0.19	Exotic	Diseases of the nervous system
<i>Ocimum basilicum</i>	0.19	Exotic	Respiratory diseases, diseases of the circulatory system
<i>Spondias purpurea</i>	0.18	Exotic	Diseases of the circulatory system

(RI=NP+NCS) where NP is obtained by dividing the number of properties (reported specific ailments) attributed to a species divided by the total number of properties attributed to the most versatile species (species with the highest number of properties). NBS is the number of body systems (ailment categories) treated by a given species divided by the total number of body systems treated by the most versatile species). Medicinal plant species used by community residents of Vitória da Conquista, recorded three or more times.

*All plants not originally native from Brazil were considered exotic

Table 4 Fidelity level (FL) values of medicinal plants cited by three or more informants. FL= (Ip/Iu x 100), where Ip is the number of informants who independently indicated the use of a species for the same major ailment and Iu the total number of informants who mentioned the plant for any major ailment

Scientific names	Category	Ip	Iu	FL (%)
<i>Spondias purpurea</i>	Hypertension	6	6	100
<i>Bauhinia forficata</i>	Diabetes	5	5	100
<i>Punica granatum</i>	Sore throat	3	4	75
<i>Matricaria chamomilla</i>	Tranquilizer	2	3	66,6
<i>Stryphnodendron adstringens</i>	Inflammation	2	3	66,6
<i>Foeniculum vulgare</i>	Flatulence	20	34	58,8
<i>Mentha spicata</i>	Influenza	10	17	58,8
<i>Amburana cearensis</i>	Indigestion	5	9	55,5
<i>Dysphania ambrosioides</i>	Inflammation	5	9	55,5
<i>Myristica fragrans</i>	Headache	2	4	50
<i>Allium sativum</i>	Hypertension	2	4	50
<i>Aloe vera</i>	Skin diseases	2	4	50
<i>Plectranthus amboinicus</i>	Influenza	3	6	50
<i>Plectranthus barbatus</i>	Indigestion	3	7	42,8
<i>Lippia alba</i>	Flatulence	19	57	33,3
<i>Cymbopogon citratus</i>	Hypertension	17	54	31,4

Table 5 Comparison of the uses of medicinal plants recorded with information gathered from the literature

Scientific Name	Ailments recorded in the present Study	Main use in the Present Study	Some medicines uses reported in the literature
<i>Alliumsativum</i>	Influenza and hypertension	Hypertension	Antifungal, antibacterial, antitumor, anti-inflammatory, antithrombotic, ⁴² anti-hypertensive, ⁴³ attenuated left ventricular diastolic dysfunction and fibrosis without significantly decreasing systolic blood pressure in hypertensive rats ⁴⁴
<i>Aloevera</i>	Hair loss, stomachache, skin inflammation, cancer, Skin diseases ulcer		Lowering LDL, increasing HDL, decreasing blood glucose level, treating genital herpes and psoriasis, ⁴⁵ positive effect on body mass, caecum and tibial bones in the short term ⁴⁶ anti-inflammatory activity. ⁴⁷
<i>Amburana cearensis</i>	Headache, stomach problems, ulcer, intestinal Infection	Indigestion	Used to treat rheumatism, cold, and sinusitis . ⁴⁰ , anti-inflammatory, ⁴¹ antibacterial activity ^{48,49}
<i>Bauhinia forficata</i>	Diabetes	Diabetes	Antidiabetic activity, ⁵⁰ hypoglycemic activity, ⁵¹ antioxidant activity, beneficial in the prevention of diabetes complications associated with oxidative stress ⁵²
<i>Cymbopogoncitratrus</i>	Hypertension, tranquilizer, cramps, stomach problems, influenza, headaches	Hypertension	Cardio protective potential is mainly because of its antioxidant activity, ⁵³ antispasmodic, hypotensive, anticonvulsant, analgesic, antiemetic, antitussive, antirheumatic, antiseptic and treatment for nervous and gastrointestinal disorders and fevers, ⁵⁴ antioxidant activity, antidiabetic activity, ⁵⁵ anti-hypertensive . ⁵⁶
<i>Dysphania ambrosioides</i>	Inflammation, toothache, scarring, ulceration, infection	Inflammation	Antitumor effect, ⁵⁷ anti-inflammatory activity . ⁵⁸ , antifungal activity, ⁵⁹ amoebicidal activity ⁶⁰

Table Continued...

Scientific Name	Ailments recorded in the present Study	Main use in the Present Study	Some medicines uses reported in the literature
<i>Foeniculumvulgare</i>	Intestinal problems, hypertension, urinary infection	Flatulence	Colic in breastfed infant, ⁶¹ antimicrobial properties ⁶² was confirmed in review the used in many parts of the world for the treatment of a number of diseases, for example, abdominal pains, antiemetic, colic in children, constipation, depurative, diarrhea, flatulence and irritable colon ⁶³
<i>Lippiaalba</i>	Stomach and intestinal problems, influenza, tranquilizer, insomnia, hypertension, fever	Tranquilizer	Sedative, anxiolytic activities, ⁶⁴ antibacterial, antifungal, ⁶⁵ vasorelaxant effect was evaluated <i>in vitro</i> in rat superior mesenteric artery rings ³⁷
<i>Matricaria chamomilla</i>	Tranquilizer and insomnia	Tranquilizer	Decreased stress in bovines, associated with inhibition of cortisol production and calming and anxiolytic effects ⁶⁶ sleep aid, sedation ⁶⁷ induce recovery from a polycystic ovary syndrome induced state in rats, but also increase dominant follicles, better endometrial tissue arrangements can be regarded as another therapeutic effect ⁶⁸ protective effects on paraquat-induced damage via oxidative stress in rat lung. ⁶⁹
<i>Mentha spicata</i>	Stomach and intestinal problems, influenza, coughs, inflammations	Influenza	Effective inhibitor of LPS-induced inflammation ⁷⁰ antioxidant effects ⁷¹ effective for antiemetic treatment ⁷²
<i>Myristica fragrans</i>	Hypertension, headache, stomach problems	Headache	Extracts of nutmeg showed a good antidiarrheal effect, with a significant sedative property. ⁷³
<i>Plectranthus amboinicus</i>	Influenza, cough, worms	Influenza	The aqueous extract has the analgesic and anti-inflammatory abilities, ^{74,75} for diabetes, which may be due to the presence of flavonoids ⁷⁶ exhibited antioxidant, diuretic, anti-inflammatory, cytotoxic and antimicrobial activities, ⁷⁷ effectiveness against methicillin-resistant <i>Staphylococcus aureus</i> skin abscesses, ⁷⁸ treatment of rheumatoid arthritis ⁷⁹
<i>Plectranthus barbatus</i>	Stomach problems (indigestion, nausea, vomiting, stomach pain, hangover)	Indigestion	The plant is also used to treat gastritis and intestinal spasms, ⁸⁰ stomach pain, nausea, vomiting and is used as purgatives, carminatives and as anthelmintic ⁸¹ antioxidant activity ⁸²
<i>Punicagranatum</i>	Sore throat, stomach problems, uterine problems	Sore throat	Wound healing activity, ⁸³ anti-inflammatory effects ⁸⁴ might be used as an antibacterial agent in controlling oral infections ⁸⁵
<i>Spondias purpurea</i>	Hypertension	Hypertension	Antimicrobial activities, ⁸⁶ antioxidant activity ⁸⁷
<i>Stryphnodendron adstringens</i>	Inflammation and healing	Inflammation	Anti-inflammatory activity ⁸⁸ antibacterial activity ^{89,90} antifungal activity, ⁹¹ reducing bacterial, ⁹² tyrosinase inhibitory activity, ⁹³ in the treatment of leucorrhea, gonorrhea, gastritis, diarrhea, bleeding, and wound healing ⁹⁴

Conclusion

Ethnobotanical studies in rural areas in Bahia, Brazil, are rare and, when available published in local reports, Brazilian journals or part of Governmental technical reports available mainly in Portuguese, these may contribute to perpetuate local knowledge far from the light of scientific knowledge. Additionally, more studies like this should be done in Bahia, by the cultural and biological diversity, and because it is the biggest state of Northeastern Brazil. The results presented here indicated that medicinal plants are commonly used by São Sebastião rural population, a total of 46 species were documented, the main indications for medicinal plant use were against digestive system, circulatory system and respiratory diseases and of general symptoms/ signs of inflammation. This evidences that the knowledge on the use of medicinal plants in this region has been preserved, since 80% of the species with medicinal properties were cultivated in the houses.

In regard to future studies, it is suggested that plants with high FL deserve attention for pharmacological studies, phytochemical, toxicological and clinical trials so that they might be included in the National Program of Medicinal Plants and Phytotherapies of the Brazilian Sistema Único de Saúde and safely prescribed by regional health teams, because there are still few plants in listed in this program. Thus, this work aimed to contribute in the long-run to improved health care in the regions of study, preservation of local knowledge, as well as search for pharmacologically principles from plants.

Acknowledgements

Work was supported by UESB. GMS was a doctoral student in RENORBIO post-graduation program. We thank Profesor Avaldo Soares Filho from UESB for kindly identifying the plants.

Conflicts of interest

Author declares there are no conflicts of interest.

Funding

None.

References

- Melo J, Santos A, Amorim E, et al. Medicinal plants used as antitumor agents in Brazil: an ethnobotanical approach. *Evid Based Complement Alternat Med*. 2011;2011:365–359.
- Rodal MJ, Barbosa MR, Thomas WW. Do the seasonal forests in northeastern Brazil represent a single floristic unit? *Braz J Biol*. 2008;68(3):467–475.
- Leal I, Silva J, Tabarelli M, Lacher T. Changing the course of biodiversity conservation in the caatinga of northeastern Brazil. *Conserv Biol*. 2005;19(3):701–706.

4. Almeida C, Amorim E, Albuquerque U, et al. Medicinal plants popularly used in the Xingó region a semi-arid location in Northeastern Brazil. *J ethnobiol ethnomed*. 2006;2:1–15.
5. Albuquerque U, Muniz de Medeiros P, Almeida A, et al. Medicinal plants of the caatinga (semi-arid) vegetation of NE Brazil: A new aport to approach. *J Ethnopharmacol*. 2007;114(3):325–354.
6. Ferreira Junior W, Siqueira C, Albuquerque U. Plant stem bark extractivism in the Northeast semiarid region of Brazil: A new aport to utilitarian redundancy model. *Evid Based Complement Alternat Med*. 2012;2012:1–11.
7. Cruz M, Medeiros P, Combariza IS, et al. I eat the manofê so it is not forgotten: local perceptions and consumption of native wild edible plants from seasonal dry forests in Brazil. *J Ethnobiol Ethnomed*. 2014;10(45):1–11.
8. Lozano A, Araújo E, Medeiros M, et al. The apparency hypothesis applied to a local pharmacopoeia in the Brazilian northeast. *J ethnobiol ethnomed*. 2014;10(2):1–17.
9. Santos J, Santos N, Santos M, et al. Honey classification from semi-arid, atlantic and transitional forest zones in Bahia, Brazil. *J Braz Chem Soc*. 2008;19(3):502–508.
10. IBGE. Instituto Brasileiro de Geografia e Estatística, Brazil. 2014.
11. <http://www.cidades.ibge.gov.br/painel/historico.php?lang=&codmun=293330&search=bahia%7Cvitoria-da-conquista%7Cinfograficos:-historico>
12. Cruz da Cunha L, Albuquerque U. Quantitative ethnobotany in an atlantic forest fragment of northeastern Brazil: implications to conservation. *Environ Monit Assess*. 2006;114(1):1–25.
13. Agra M, Freitas P, Barbosa-filho J. Synopsis of the plants known as medicinal and poisonous in Northeast of Brazil. *Braz J Pharmacog*. 2007;17(1):114–140.
14. Braun E, Alves L, Soares M, et al. Technological forecasting on phytotherapies development in Brazil. *International Journal of Medical, Health, Pharmaceutical and Biomedical Engineering*. 2012;6(7):14–18.
15. Christo A, Bruni RG, Araújo F, et al. Evidence for conservation and sustainable use in a fragment of the Atlantic forest in southeastern Brazil by a traditional human group. *SpringerPlus*. 2012;1:1–12.
16. Antonio G, Tesser C, Pires RM. Phytotherapy in primary health care. *Rev Saúde Pública*. 2014;48(3):541–553.
17. Bruning M, Mosegui G, Vianna C. The use of phytotherapy and medicinal plants in primary healthcare units in the cities of Cascavel and Foz do Iguaçu-Paraná: the viewpoint of health professionals. *Cien Saude Colet*. 2012;17(10):2675–2685.
18. Christo A, Guedes-Bruni R, Silva A. Local knowledge on medicinal plant gardens in a rural community near the Atlantic Rain Forest, southeastern Brazil. *Braz J Pharmacog*. 2010;20(4):494–501.
19. Silva H, Caraciolo R, Marangon L, et al. Evaluating different methods used in ethnobotanical and ecological studies to record plant biodiversity. *J ethnobiol ethnomed*. 2014;10(1):48.
20. Phondani P, Bhatt A, Elsarrag E, et al. Ethnobotanical magnitude towards sustainable utilization of wild foliage in Arabian Desert. *J Tradit Complement Med*. 2016;6(3):209–218.
21. Baptista M, Ramos M, Albuquerque U, et al. Traditional botanical knowledge of artisanal fishers in southern Brazil. *J ethnobiol ethnomed*. 2013;9:54.
22. Chinsemu C. Ethnobotanical Study of Plants Used in the Management of HIV/AIDS-Related Diseases in Livingstone, Southern Province, Zambia. *Evid Based Complement Alternat Med*. 2016;2016:1–14.
23. Tufts H, Harris C, Bukania Z, et al. Antioxidant and anti-inflammatory activities of kenyan leafy green vegetables, wild fruits, medicinal plants with potential relevance for kwashiorkor. *Evid Based Complement Alternat Med*. 2015;2015:1–10.
24. Albuquerque U, Sousa Araújo T, Ramos M, et al. How ethnobotany can aid biodiversity conservation: reflections on investigations in the semi-arid region of NE Brazil. *Biodivers Conserv*. 2009;18(1):127–150.
25. Zank S, Hanazaki N. Exploring the Links between ethnobotany, local therapeutic practices, and protected areas in Santa Catarina coastline, Brazil. *Evid Based Complement Alternat Med*. 2012;2012:15.
26. WHO. International Statistical Classification of Diseases and Related Health Problems. 2011.
27. Heinrich M, Ankli A, Frei B, et al. Medicinal plants in Mexico: healers' consensus and cultural importance. *Soc Sci Med*. 1998;47(11):1859–1871.
28. Bennett B, Prance G. Introduced plants in the indigenous pharmacopoeia of northern south America. *Econ Bot*. 2000;54(1):90–102.
29. Friedman J, Yaniv Z, Dafni A, et al. A preliminary classification of the healing potential of medicinal plants, based on a rational analysis of an ethnopharmacological field survey among bedouins in the Negev desert, Israel. *J Ethnopharmacol*. 1986;16(2-3):275–287.
30. Afolayan A, Grierson D, Mbeng W. Ethnobotanical survey of medicinal plants used in the management of skin disorders among the Xhosa communities of the Amathole District, Eastern Cape, South Africa. *J Ethnopharmacol*. 2014;153(1):220–232.
31. Asnake S, Teklehaymanot T, Hymete A, et al. Survey of medicinal plants used to treat malaria by Sidama People of Boricha District, Sidama Zone, South Region of Ethiopia. *Evid Based Complement Alternat Med*. 2016;2016:1–9.
32. Oliveira H, Kffuri C, Casali V. Ethnopharmacological study of medicinal plants used in Rosário. *Brazilian Journal of Pharmacology*. 2010;20(2):256–260.
33. Bieski I, Rios Santos F, Oliveira R, et al. Ethnopharmacology of medicinal plants of the pantanal region (Mato Grosso, Brazil). *Evid Based Complement Alternat Med*. 2012;2012:1–36.
34. Brito M, Senna-Valle L. Diversity of plant knowledge in a 'Caiçara' community from the Brazilian Atlantic Forest coast. *Acta Bot Bras*. 2012;26(4):735–747.
35. Meragiaw M, Asfaw Z, Argaw M. The status of ethnobotanical knowledge of medicinal plants and the impacts of resettlement in Delanta, Northwestern Wello, Northern Ethiopia. *Evid Based Complement Alternat Med*. 2016;2016:1–24.
36. Hassan-Abdallah A, Merito A, Hassan S, et al. Medicinal plants and their uses by the people in the Region of Randa, Djibouti. *J Ethnopharmacol*. 2013;148(2):701–713.
37. Garcia D, Domingues M, Rodrigues E. Ethnopharmacological survey among migrants living in the Southeast Atlantic Forest of Diadema, São Paulo, Brazil. *J ethnobiol ethnomed*. 2010;6(1):29.
38. Maynard L, Santos K, Cunha P, et al. Chemical composition and vasorelaxant effect induced by the essential oil of *Lippia alba* (Mill.) N.E. Brown. (Verbenaceae) in rat mesenteric artery. *Indian J Pharmacol*. 2011;43(6):694–698.
39. Hatano V, Torricelli A, Giassi A, et al. Anxiolytic effects of repeated treatment with an essential oil from *Lippia alba* and (R)-(-)-carvone in the elevated T-maze. *Braz J Med Biol*. 2012;45(3):238–243.
40. Pascual M, Slowing K, Carretero M, et al. Antiulcerogenic activity of *Lippia alba* (Mill.) N. E. Brown (Verbenaceae). *Farmaco*. 2001;56(7):501–504.
41. Farias D, Souza I, Viana M, et al. Antibacterial, antioxidant, and anticholinesterase activities of plant seed extracts from Brazilian semiarid region. *Biomed Res Int*. 2013;2013:1–9.

42. Lopes A, Magalhães T, Uchôa D, et al. Afrormosin, an Isoflavonoid from *Amburana cearensis* A. C. Smith, Modulates the Inflammatory Response of Stimulated Human Neutrophils. *Basic Clin Pharmacol Toxicol.* 2013;113:363–369.
43. Lanzotti V. The analysis of onion and garlic. *J Chromatogr A.* 2006;1112:3–22.
44. Corzo-Martinez M, Corzo N, Villamiel M. Biological properties of onions and garlic. *Trends Food Sci Technol.* 2007;18(12):609–625.
45. Hara Y, Noda A, Miyata S, et al. Effects of Aged Garlic Extract on Left Ventricular Diastolic Function and Fibrosis in a Rat Hypertension Model. *Experiment Anim.* 2013;62(4):305–310.
46. Ahlawat K, Khatkar B. Processing, food applications and safety of *Aloe vera* products: a review. *J Food Sci Technol.* 2011;48(5):525–533.
47. Beyaa W, Davidson B, Erlwangera K. The effects of crude aqueous and alcohol extracts of *Aloe vera* on growth and abdominal viscera of suckling rats. *Afri J Tradit Complement Altern Med.* 2012;9(4):553–560.
48. Lee Y, Ju H, Kim Y, et al. Enhancement of anti-inflammatory activity of *Aloe vera* adventitious root extracts through the alteration of primary and secondary metabolites via salicylic acid elicitation. *PLoS one.* 2013;8(12):1–13.
49. Figueredo F, Ferreira E, Lucena B, et al. Modulation of the antibiotic activity by extracts from *Amburana cearensis* AC Smith and *Anadenanthera macrocarpa* (Benth) Brenan. *Biomed Res Int.* 2013;2013:1–5.
50. Sá M, Ralph M, Nascimento D, et al. Phytochemistry and preliminary assessment of the antibacterial activity of chloroform extract of *Amburana cearensis* (Allemão) AC Sm against *Klebsiella pneumoniae* Carbapenemase-Producing Strains. *Evid Based Complement Alternat Med.* 2014;2014:1–7.
51. Pepato M, Keller E, Baviera A, et al. Anti-diabetic activity of *Bauhinia forficata* decoction in streptozotocin-diabetic rats. *J Ethnopharmacol.* 2002;81:191–197.
52. Menezes F, Minto A, Ruela H, et al. Hypoglycemic activity of two Brazilian *Bauhinia* species: *Bauhinia forficata* L. and *Bauhinia monandra* Kurz. *Braz J Pharmacog.* 2007;17(1):8–13.
53. Khalil N, Pepato M, Brunetti I. Free radical scavenging profile and myeloperoxidase inhibition of extracts from antidiabetic plants: *Bauhinia forficata* and *Cissus sicyoides*. *Biol Res.* 2008;41:165–171.
54. Gayathri K, Jayachandran K, Vasanthi HR, et al. Cardioprotective effect of lemon grass as evidenced by biochemical and histopathological changes in experimentally induced cardiotoxicity. *Hum Exp Toxicol.* 2011;30(8):1073–1082.
55. Shah G, Shri R, Panchal V, et al. Scientific basis for the therapeutic use of *Cymbopogon citratus* stapf (Lemon grass). *J Adv Pharm Technol Res.* 2011;2(1):3–8.
56. Mirghani M, Liyana Y, Parveen J. Bioactivity analysis of lemongrass (*Cymbopogon citratus*) essential oil. *Int Food Res J.* 2012;19(2):569–575.
57. Olorunnisola S, Asiyambi H, Hammed A, et al. Mini Review Biological properties of lemongrass: An overview. *Int Food Res J.* 2014;21(2):455–462.
58. Nascimento F, Cruz G, Pereira P, et al. Ascitic and solid Ehrlich tumor inhibition by *Chenopodium ambrosioides* L. treatment. *Life Sci J.* 2006;78(22):2650–2653.
59. Ibrinke G, Ajiboye K. Studies on the anti-inflammatory and analgesic properties of *Chenopodium ambrosioides* leaf extract in rats. *Int J Pharmacol.* 2007;3(1):111–115.
60. Sousa Z, Oliveira F, Conceição A, et al. Biological activities of extracts from *Chenopodium ambrosioides* Lineu and *Kielmeyera neglecta* Saddi. *Ann Clin Microbiol Antimicrob.* 2012;11(20):1–7.
61. Avila-Blanco M, Rodríguez M, Moreno Duque J, et al. Amoebicidal activity of essential oil of *Dysphania ambrosioides* (L.) Mosyakin & Clemants in an amoebic Liver abscess hamster model. *Evid Based Complement Alternat Med.* 2014;2014:1–7.
62. Savino F, Cresi F, Castagno E, et al. A randomized double-blind placebo-controlled trial of a standardized extract of *Matricariae recutita*, *Foeniculum vulgare* and *Melissa officinalis* (ColiMil) in the treatment of breastfed colicky infants. *Phytother Res.* 2005;19(4):335–340.
63. Kaur G, Arora D. Bioactive potential of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi* belonging to the family Umbelliferae-Current status. *J Med Plant Res.* 2010;4(2):87–94.
64. Badgujar S, Patel V, Bandivdekar A. *Foeniculum vulgare* Mill : A review of its botany, phytochemistry, pharmacology, contemporary application and toxicology. *Biomed Res Int.* 2014;2014:1–32.
65. Hennebelle T, Sahpaz S, Joseph H, et al. Ethnopharmacology of *Lippia alba*. *J Ethnopharmacol.* 2008;116(2):211–222.
66. Ara N, Amran M, Wahid M, et al. *In vitro* antimicrobial and cytotoxic of leaves and flowers extracts from *Lippia alba*. *Pak J Biol Sci.* 2009;12(1):87–90.
67. Reis L, Pardo P, Oba E, et al. *Matricaria chamomilla* CH₁₂ decreases handling stress in Nelore calves. *J Vet Sci.* 2006;7(2):189–192.
68. Srivastava J, Shankar E, Gupta S. Chamomile: A herbal medicine of the past with bright future. *Mol Med Rep.* 2010;3(6):895–901.
69. Zangeneh Z, Minaee B, Amirzargar A, et al. Effects of chamomile extract on biochemical and clinical parameters in a rat model of polycystic ovary syndrome. *J Reprod Infertil.* 2010;11(3):169–174.
70. Ranjbar A, Mohsenzadeh F, Chehregani A, et al. Ameliorative effect of *Matricaria chamomilla* L. on paraquat: Induced oxidative damage in lung rats. *Pharmacognosy Res.* 2014;6(3):199–203.
71. Pearson W, Fletcher R, Kott L, et al. Protection against LPS-induced cartilage inflammation and degradation provided by a biological extract of *Mentha spicata*. *BMC Complement Altern Med.* 2010;10:1–19.
72. Arumugam P, Ramesh A. In-vivo antioxidant effects of ethyl acetate fraction of *Mentha spicata* L. on 4-Nitroquinoline-1-Oxide Injected Mice. *Iran J Pharm Res.* 2011;10(4):787–793.
73. Tayarani-Najaran Z, Talasaz-Firooz E, Nasiri R, et al. Antiemetic activity of volatile oil from *Mentha spicata* and *Mentha piperita* in chemotherapy-induced nausea and vomiting. *Ecancermedicalscience.* 2013;7:290.
74. Grover J, Khandkar S, Vats V, et al. Pharmacological studies on *Myristica fragrans* antidiarrheal, hypnotic, analgesic and hemodynamic (blood pressure) parameters. *Methods Ind Exp Clin Pharmacol.* 2002;24(10):675–680.
75. Chiu Y, Huang T, Chiu C, et al. Analgesic and antiinflammatory activities of the aqueous extract from *Plectranthus amboinicus* (Lour.) Spreng. Both *In Vitro* and *In Vivo*. *Evid Based Complement Alternat.* 2012;2012:1–11.
76. Ravikumar V, Dhanamani M, Sudhaman T. *In-vitro* anti- inflammatory activity of aqueous extract of leaves of *Plectranthus amboinicus* (Lour.) Spreng. *Anc Sci Life.* 2014;28(4):7–9.
77. Koti B, Gore A, Thippeswamy AH, et al. Alcoholic leaf extract of *Plectranthus amboinicus* regulates carbohydrate metabolism in alloxan-induced diabetic rats. *Indian J Pharmacol.* 2011;4(3):286–290.
78. El-hawary S, El-sofany R, Abdel-Monem A, et al. Polyphenolics content and biological activity of *Plectranthus amboinicus* (Lour.) spreng growing in Egypt (Lamiaceae). *Phcog J.* 2012;4(32):45–54.
79. Oliveira F, Torres A, Gonçalves T, et al. Efficacy of *Plectranthus amboinicus* (Lour.) Spreng in a murine model of methicillin-resistant *Staphylococcus aureus* skin abscesses. *Evid Based Complement Alternat Med.* 2013;2013:1–9.

80. Chang J, Chengv C, Hung L, et al. Potential Use of *Plectranthus amboinicus* in the treatment of rheumatoid arthritis. *eCAM*. 2010;7(1):115–1120.
81. Câmara C, Nascimento N, Macêdo-Filho C, et al. Antispasmodic effect of the essential oil of *Plectranthus barbatus* and some major constituents on the guinea-pig ileum. *Planta Med*. 2003;69(12):1080–1085.
82. Likhoba C, Simmonds M, Paton A. *Plectranthus*: A review of ethnobotanical uses. *J Ethnopharmacol*. 2006;103:1–24.
83. Maioli M, Alves L, Campanini A, et al. Iron chelating-mediated antioxidant activity of *Plectranthus barbatus* extract on mitochondria. *Food Chem*. 2010;122(1):203–208.
84. Murthy K, Veigas V, Murthy U. Study on wound healing activity of *Punica granatum* peel. *J Med Food*. 2004;7(2):256–259.
85. Lansky E, Newman R. *Punica granatum* (pomegranate) and its potential for prevention and treatment of inflammation and cancer. *J Ethnopharmacol*. 2007;109(2):177–206.
86. Abdollahzadeh S, Mashouf R, Mortazavi H, et al. Antibacterial and antifungal activities of *Punica granatum* peel extracts against oral pathogens. *J Dent (Tehran)*. 2011;8(1):1–6.
87. Miranda-Cruz E, Espinosa-Moreno J, Centurión-Hidalgo D, et al. Actividad antimicrobiana de extractos de *Psidium friedrichsthalianum* L., *Pterocarpus hayesii* L., *Tynanthus guatemalensis* L. y *Spondias purpurea* L. *Bol Latinoam Caribe Plant Med Aromat*. 2012;11(4):354–361.
88. Gregoris E, Pereira Lima G, Fabris S, et al. Antioxidant properties of Brazilian tropical fruits by correlation between different assays. *Biomed Res Int*. 2013;2013:1–8.
89. Lima J, Martins D, De Souza P. Experimental evaluation of stem bark of *Stryphnodendron adstringens* (Mart.) Coville for antiinflammatory activity. *Phytother Res*. 1998;12:218–220.
90. Audi E, Eduardo C, Toledo M, et al. Biological activity and quality control of extract and stem bark from *Stryphnodendron adstringens*. *Acta Farm Bonaerense*. 2004;23(3):328–333.
91. Costa M, Ishida K, Kaplum V, et al. Safety evaluation of proanthocyanidin polymer-rich fraction obtained from stem bark of *Stryphnodendron adstringens* (Barbatimao) for use as a pharmacological agent. *Regul Toxicol Pharmacol*. 2010;58(2):330–335.
92. Ishida K, Rozental S, De Mello J, et al. Activity of tannins from *Stryphnodendron adstringens* on *Cryptococcus neoformans*: effects on growth, capsule size and pigmentation. *Ann Clin Microbiol Antimicrob*. 2009;8:29.
93. Cavalcante G, Resende R, Leite J, et al. Potential of ethyl acetate fractions of *Stryphnodendron adstringens* shells and fruit extracts of *Caesalpinia ferrea* to control bacterial leaf speck and on the potentiation of defense enzymes in tomato. *Trop Plant Pathol*. 2014;39(4):267–274.
94. Souza P, Elias S, Simeoni L, et al. Plants from brazilian cerrado with potent tyrosinase inhibitory activity. *PloS one*. 2012;7(11):1–7.
95. Costa M, Palazzo de Mello J, Kaneshima E, et al. Acute and chronic toxicity of an aqueous fraction of the stem bark of *Stryphnodendron adstringens* (Barbatimão) in Rodents. *Evid Based Complement Alternat Med*. 2013;2013:9.