

# Acute effect of Quantec therapy on cardiovascular risk factors and autonomous modulation of heart rate

## Abstract

**Background:** Quantec is a new instrumental bioinformation therapy, that acts on the patient's electromagnetic fields to restore the body's normal functions. Heart rate variability (HRV) is an important biomarker applied in research to measure physiological effects.

**Method:** Approved by the Ethics Committee (1068/2014). 10 individuals, mean age 60.4±6.4 years, with at least one cardiovascular risk. An instrumental biocommunication device (Quantec 6.0-R2-08, Munich, Germany) was used for an acute application of 12 seconds of treatment. The recording of the HR and the RR intervals (Polar RS800CX, Polar Electro Oy, Kempele, Finland) was done with the volunteer sitting in a state of rest for 10 minutes before application and 5 minutes afterwards. Only the RRI series with more than 95% sinus beats were analyzed, having selected the 256 most stable points and calculated the HRV indices (Software Kubios HRV, version 2.0, University of Kuopio, Finland).

**Results:** Δ variation between pre and post moments of the LH/HF ratio had a significant negative correlation ( $p = 0.006$ ). Glycemia showed a significant reduction ( $p = 0.010$ ) as did the Triglycerides ( $p = 0.024$ ).

**Discussion:** Quantec had a positive effect on blood concentration values of cholesterol, triglycerides and glycemia. In the altered LH/HF values, it showed greater reductions in sympathetic activity and an increase in parasympathetic activity.

**Conclusion:** Acute treatment with Quantec was effective in reducing metabolic imbalances.

**Keywords:** hypertension, additional therapies, dyslipidemia, diabetes mellitus, holistic health

Volume 11 Issue 3 - 2018

Jose Alfredo Ordenes Mora,<sup>1,2</sup> Eduardo Federighi Baisi Chagas,<sup>3</sup> Robison Jose Quiterio<sup>1,2</sup>

<sup>1</sup>Physical Therapy and Occupational Therapy Department, São Paulo State University (UNESP) - School of Philosophy and Sciences, Brazil

<sup>2</sup>Laboratory of Investigation in Biocommunication, Physical Exercise and Cardiac Autonomic Modulation (LIBEM)

<sup>3</sup>Physical Education Department, Marília University (UNIMAR), Brazil

**Correspondence:** Jose Alfredo Ordenes Mora, Physical Therapy and Occupational Therapy Department, São Paulo State University (Unesp) - School of Philosophy and Sciences – Marília, Av Hygino Muzzi, 737, Marília, São Paulo, Brazil, Tel +5514 3402 1331, Email jordenesmora@gmail.com

**Received:** April 20, 2018 | **Published:** May 18, 2018

## Introduction

Despite advances in scientific research, the prevalence of dyslipidemias (DLP), Hypertension (SAH) and Diabetes Mellitus (DM) continues to increase worldwide.<sup>1-6</sup> These are the main risk factors for the cardiovascular diseases that generally affect the heart's autonomic nervous system, this being the subject of intensive scientific research. These studies suggest that a decrease in the vagal tone may precede the development of one of these critical risk factors for cardiovascular diseases. Thus, over the last few years this system has been the subject of intensive research. One of the outstanding methods used in the research was the Heart Rate Variability (HRV) study, due to its great disease forecasting power, a lower HRV being associated with greater cardiovascular morbidity and mortality.<sup>7-10</sup> As metabolic and cardiovascular diseases have reached epidemic proportions worldwide and are difficult to cure, complementary medicine based on quantum physics has awakened great interest and opened up new research perspectives and areas, such as magnetobiology and biomagnetism.<sup>11,12</sup>

The theory proposes acting on diseases through the transmission of specific electronic vibrations, to bring the atom back to the normal vibration; thereby eliminating the disease.<sup>13</sup> This technique involves

using instrumental biocommunication devices that are capable of interpreting imbalances in the body's electromagnetic fields, re-sending electronic vibration patterns in order to restore normal patterns. Quantec is an instrumental bioinformation equipment, which through the white noise diode can capture the subtle bodies and connect with the consciousness. It has been used as a direct interface between an information generator and the patient's subconscious to perform treatment modifying altered patterns of functioning through change in the internal information of each individual.<sup>14,15</sup>

As the heart generates a huge source of bioinformation for the whole body and is responsible for the being's functioning as a whole, it delivers biological information through heart sounds, pulse and the heart rate,<sup>16</sup> this study's hypothesis is that a vibration therapy session by instrumental biocommunication increases the heart's parasympathetic modulation and shows an improvement in the clinical characteristics of individuals with hypertension, diabetes and dyslipidemia.

## Methods

The study concerned 10 individuals of both sex, aged between 52 and 67 with a previous clinical diagnosis of imbalance in at least one of the risk factors listed: SAH, DLP and DM. It did not include those

with respiratory, neurological and metabolic diseases of a different nature. All were informed about the experimental procedures and signed the free and informed consent form. Research study approved under the number 1068/2014. The assessments were performed during the morning, in order to formally record the circadian variations in the organism<sup>17</sup>. The experiment was performed in an air-conditioned environment, temperature at  $23\pm 2^{\circ}\text{C}$  and air relative humidity at  $50\pm 10\%$ .

### Application of Instrumental Bioinformation

On the first day the volunteer's personal details were recorded in the Instrumental biocommunication device (Quantec 6.0-R2-08, Munich, Germany) including a photograph of the patient's entire face and body taken without the use of flash or artificial light, with a 12-megapixel camera. The camera was positioned two meters' away from the volunteer, who was positioned on a standardized white wall away from any electromagnetic network within a 2-meter radius. Then, a sentence was written on *Healing Sheet* page: "I am very grateful for the rebalancing of my blood's cholesterol, triglyceride and glycemia levels. Thank you very much!". *HealingSheet* is an input mask that contains useful information for correcting the desired imbalances, by means of a statement that sets out the objective. The photo of the patient's entire body was then included in the *HealingSheet* page. Once the *Healing Sheet* was defined, the morphogenetic field selection page was opened and the "Metabolic Treatment" spreadsheet selected, which contained a set of different frequencies in order to restore the body's normal vibration patterns. Next, the generator carried out a search of the body's imbalances, through the white noise diode, a reading was performed and established a correlation between the two photographs and the patient's hand, that was positioned on the gold clip. The results were ordered according to the percentage of correlation between the photographs and the patient's hand, and the potency and intensities were automatically and individually defined by the device.

On the second evaluation, 2 days after the last detection, the patient was asked to remove any kind of device or instrument that might generate any kind of electromagnetic interference (mobile, watch, iPad, etc). The patient's blood pressure was measured by the auscultatory method, after remaining in a seated position for 20 minutes. The pressoric levels were classified according to the 6th Brazilian Arterial Hypertension Directive<sup>18</sup>; Cholesterol, triglycerides and glycemia by means of a reflectance spectrophotometry biochemical analyzer (Accutrend Plus, Roche Diagnostics, 2007). The total cholesterol and triglycerides concentrations were classified according to the 5th Brazilian Dyslipidemia and Atherosclerosis Prevention Directive,<sup>19</sup> and the glycemia was classified according to the Brazilian Cardiology Society Directive.<sup>20</sup>

Then the RR interval (RRi) transmitter was placed on the patient's chest at the height of the xiphoid process (Polar RS800CX, Polar Electro Oy, Kempele, Finland). The receiver was held by the researcher and the patient remained seated and held on the palm of his hand the device's gold clip, which contains the white noise diode.

The RRi was recorded for 10 minutes, with the volunteer breathing spontaneously and after stabilization of the HR. The Quantec treatment starts at the minute 10 and lasted for 12 seconds. The recording of the RRi was interrupted at minute 10 and the BP and

the blood biochemistry measured again. These measurements were repeated after 30 minutes.

### Analysis of the HRV data

The instantaneous HR and RRi were converted into text files and analyzed (*Kubios HRV*, version 2.0, University of Kuopio, Finland). Only the series with over 95% of sinus rhythm beats and the 256 most stable points of the stretches were selected: before and after the application of bioinformation.

In the time domain, the following statistical calculations were carried out: average and standard deviation of the HR values in beats per minute (bpm) and of the RRi (ms); standard deviation of the square root of the mean of the squares of the differences between the successive normal intervals (RMSSD), expressed in milliseconds (ms); this last index representing the parasympathetic modulation.<sup>21,22</sup>

We also calculated the SD1 and SD2 indices of the *Poincaré plot*, which is a map of points in Cartesian coordinates, where every point is represented on the horizontal X axis (abscissae) by the previous normal RRi and on the Y axis (ordinate) by the following RRi. The standard deviation of the perpendicular points and along the length of the identity line give rise to the SD1 and SD2 indices respectively. The SD1 index measures the standard deviation of the distances of the points towards the diagonal  $y=x$ , is related to the short-term variability, is influenced by the respiratory sinus arrhythmia and represents the parasympathetic activity. The SD2 measures the standard deviation of the distances from the points towards the straight line  $y=-x+RRm$ , where  $RRm$  is the mean of the RRi, with long-term variability and reflects the overall variability. The relationship between them (SD1/SD2) shows the ratio between the short and long variations of the RR intervals.<sup>22,23</sup>

For the spectral analysis, an autoregressive model was used and the following spectral indices calculated: high frequency (HF – 0.15 at 0.4Hz) which corresponds to the respiratory modulation and to the parasympathetic modulation of the vagus nerve on the heart; low frequency (LF–0.04 at 0.15Hz), which represents sympathetic and parasympathetic modulation, with sympathetic predominance; and the LF/HF ratio, which represents the sympathovagal balance.<sup>19</sup> The LF data was presented in absolute values ( $\text{ms}^2$ ) and the HF as a standardized unit (un). The standardized data was calculated by dividing the spectral power density of a given band (i.e. HF) by the total power, subtracting the very low frequency band and multiplying by 100.<sup>24</sup>

### Statistical analysis

The results were organized in the descriptive statistical form, with mean, standard deviation (SD) and confidence interval (CI 95%) values. The standard distribution was checked by the *Shapiro-Wilk* with Lillifors correction. In order to analyze the variation between the pre (M1) and post operation (M2) moments, and also 30 minutes after the end of the operation (M3), the delta variation ( $\Delta$ ) (post-pre) was calculated. The variation in relation to the initial values are graphically represented with their respective moment correlation x Pearson product coefficients. To analyze the acute effect of the intervention on the HRV indices, the *t student* test was applied for paired samples and their non-parametric similar (Wilcoxon) when the data did not

represent the normal distribution. An ANOVA was constructed for repetitive measurements in order to test the time effects on SBP, DBP, GL, TG and TC. Mauchly's test was used to test the sphericity hypothesis. If the sphericity hypothesis was rejected, the analyses were based on the Greenhouse-Geisser multivariate test. When the time effect was statistically significant, the multiple LSD comparison test was carried out. For all the analyses the SPSS software version 19.0 Windows version was used, adopting a significance level of 5%.

## Results

The average age of the sample elements was  $60.4 \pm 6.4$ , with minimum age 52 and maximum age 67 years. The sample was composed of 6 men (60%) and 4 women (40%). SAH (80%) was the most prevalent factor, followed by DM (40%) and DLP (40%). Table 1 presents the systolic arterial pressure (SBP) values, Diastolic blood pressure (DBP), GL, TG and TC for moments M1, M2 and M3. A significant reduction was identified between M1 and M2, and between M1 and M3 for SBP. GL presented similar reduction behavior ( $p=0.010$ ), being significant between M1 and M2, and M1 and M3. The variation between the moments was significant ( $p=0.024$ ) for the TG values, a significant difference being observed between moments M2 and M3.

**Table 1** Arterial Pressure and biochemical variables for the initial pre-operation moment (M1), after the operation (M2) and 30 minutes after the operation (M3)

	M1	M2	M3	
	MEAN $\pm$ PD	MEAN $\pm$ PD	MEAN $\pm$ PD	p-value
SBP	125.0 $\pm$ 7.4 <sup>a</sup>	117.8 $\pm$ 8.5 <sup>b</sup>	121.2 $\pm$ 5.7 <sup>b</sup>	0.053
DBP	82.2 $\pm$ 4.6	80.6 $\pm$ 7.7	80.0 $\pm$ 2.9	0.530
GL	95.1 $\pm$ 22.3 <sup>a</sup>	81.2 $\pm$ 22.3 <sup>b</sup>	72.7,4 $\pm$ 19.3 <sup>b</sup>	0.010*
TG	210.8 $\pm$ 55.7	203.6 $\pm$ 48.0 <sup>a</sup>	224.2 $\pm$ 59.3 <sup>b</sup>	0.024*
TC	174.9 $\pm$ 21.1	169.0 $\pm$ 17.9	171.9 $\pm$ 18.7	0.324

Note: \*  $p \leq 0.05$  significant difference between the moments (Anova repeated measurements test). Different letters overwritten indicate significant differences ( $p \leq 0.05$ ) between the moments, and the same letters overwritten indicate the absence of significant differences between the moments, (multiple LSD comparison test).

TC, Cholesterol Total; DB, Diastolic Blood Pressure; GL, Glycemia; SBP, Systolic Blood Pressure; TG, Triglycerides

The  $\Delta$  variations of intervals M2-M1 and M3-M1 in relation to the initial M1 value for the variable, TC, GL and TG, are graphically represented with their respective moment correlation x Pearson product coefficient ( $r$ ) and significance value ( $p$ ) in Figure 1. The significant correlation between the initial values and the  $\Delta$  variation were not verified. This relationship with the initial values was only non-identifiable for the TG values in the analysis of the  $\Delta$  variation between the M3 and M1 moments in relation to the initial M1

values. There are good correlation values despite a reduced size of the sample, which does not permit statistical significance. The  $\Delta$  variations of intervals M2-M1 and M3-M1 in relation to the initial M1 value for the variable, SBP and DBP, are graphically represented with their respective moment correlation x Pearson product coefficient ( $r$ ) and significance value ( $p$ ) in Figure 2. The  $\Delta$  variation between the M1 and M3 moments of the DBP presented negative and significant correlation ( $p=0.012$ ) with the initial M1 values.

Table 2 presents average and standard deviation (SD) of the HRV indices in the pre and post operation moments, followed by the average delta ( $\Delta$ ) variation (post - pre) value and its respective Confidence Interval of 95% (lower | higher). Statistically significant differences between the pre and post operation moments were not observed. Insofar as the delta variation values indicate a reduction trend in the HRV indices measured. Figure 3 present the pre and post-intervention moments in relation to the initial values for the HRV indices referring to parasympathetic and sympathetic values, with their respective Pearson product moment correlation values and significance value ( $p$ ). A significant correlation between the delta variations and the initial values of the parasympathetic (Figure 1) and sympathetic indices (Figure 2) of the HRV was not observed, indicating that the variations observed were independent of the initial values.


The  $\Delta$  variations of the post-pre intervals in relation to the initial values of the LH/HF ratio are graphically represented with their respective moment correlation x Pearson product coefficient ( $r$ ) and significance value ( $p$ ) in Figure 4. The  $\Delta$  variation between the pre and post-intervention moments of the LH/HF ratio presented negative and significant correlation ( $p=0.006$ ) with the initial values.

**Table 2** HRV indices pre and post intervention followed by the average delta ( $\Delta$ ) variation (post - pre) value and its respective Confidence Interval (CI) of 95% (lower|higher)


	Pre	Post	$\Delta$ (post-pre)		
	MEAN $\pm$ SD	MEAN $\pm$ SD	MEAN	$\Delta$ CI 95%	P value
MeanHR	65.4 $\pm$ 9.4	65.0 $\pm$ 9.3	-0.34	-0.91   0.22	0.200 <sup>c</sup>
RMSSD <sup>w</sup>	27.9 $\pm$ 19.5	27.6 $\pm$ 21.2	-0.28	-5.42   4.86	0.721 <sup>w</sup>
pNN50	13.0 $\pm$ 21.6	11.5 $\pm$ 22.4	-1.42	-5.44   2.58	0.442 <sup>c</sup>
LH	223.7 $\pm$ 153	217 $\pm$ 169	-6.70	-105   91.8	0.881 <sup>c</sup>
HF	52.8 $\pm$ 25.5	51.2 $\pm$ 22.5	-0.80	-9.51   7.91	0.840 <sup>c</sup>
LH/HF <sup>w</sup>	1.58 $\pm$ 1.76	1.36 $\pm$ 1.08	-0.21	-1.13   0.70	0.959 <sup>w</sup>
SD1	19.7 $\pm$ 13.8	18.5 $\pm$ 13.7	-1.20	-6.25   3.85	0.604 <sup>c</sup>
SD2	42.0 $\pm$ 14.0	37.3 $\pm$ 18.2	-4.69	-12.3   2.99	0.201 <sup>c</sup>

Note: \* $p \leq 0.05$  significant difference between M1 and M2. <sup>c</sup> student t test. <sup>w</sup>Wilcoxon test.

SD, standard deviation; RMSSD, root mean square of successive differences; pNN50, the proportion of NN50 divided by total number of NNs; LH, low frequency; HF, High frequency; LF/HF, sympathovagal balance; SD1, standard deviation of instantaneous beat-to-beat interval variability; SD2, the continuous long-term R/R interval variability.




**Figure 1** Delta variation ( $\Delta$ ) of the Total Cholesterol (TC), Glycemia (GL) and Triglycerides (TG) for the moment immediately after the operation (M2-M1) and 30 minutes after the intervention (M3-M1) in relation to the initial values (M1).




**Figure 2** Delta variation ( $\Delta$ ) of the Systolic Blood Pressure (SBP) and Diastolic Blood Pressure (DBP) for the moment immediately after the intervention (M2-M1) and 30 minutes after the operation (M3-M1) in relation to the initial values (M1).

## Discussion

All the variables studied (SBP, DBP, GL, TG, TC) decreased after the acute intervention, and according to Table 1, only GL and TG reached statistical significance between the initial moment and at the moments immediately afterwards (M2) and the late moment (M3). There was also a decrease in the sympathetic modulation of the heart rate. SBP presented a significant difference between M1 and M2 and between M1 and M3, demonstrating an effect that was maintained



**Figure 3** Delta ( $\Delta$ ) (post-pre) variation of the values of the parasympathetic and sympathetic activity indices of the HRV, where the delta is represented on the Y axis and the initial values at the pre-intervention moment on the X axis.



Note:  $r$  = Pearson correlation coefficient, and  $p$  = significance value corresponding to the Pearson correlation test.

**Figure 4** Delta ( $\Delta$ ) (post-pre) variation of the sympathovagal balance index of the HRV, where the delta is represented on the Y axis and the initial values at the pre-intervention moment on the X axis.

for 30 minutes. DBP did not presents significant difference between the moments, despite the reduction trend. Glycemia presented a significant reduction between M1 and M2, and M1 and M3. TC did not present a significant difference, despite the reduction trend. TG presented a significant increase between the M2 and M3 moments, but not between M1 and M3, suggesting that the effects of the intervention were not maintained 30 minutes after the treatment. The  $\Delta$  variation between the M1 and M3 moments of the DBP presented a negative and significant correlation ( $p=0.012$ ) with the initial M1 values,

demonstrating that despite the fact that it did not present significant difference in the group, Quantec was selective for the reduction of DBP only in patients who presented alterations in this variable.

In Figure 4 a negative correlation was observed between the initial values and the delta variation of the values of the LH/HF ratio that represents the sympathovagal balance. These results indicate that patients with very high initial LH/HF values present more reduction in sympathetic activity and an increase in parasympathetic activity. Patients without alterations did not change their initial values, suggesting that the intervention had an individualized positive effect on the sympathovagal balance of each participant. We therefore propose that the observed effects result from a balancing of the biological information of each individual, given that research studies with other forms of bioinformation managed to balance alterations in blood pressure.<sup>25,26</sup>

Quantec had a positive effect on the values of the blood concentrations of cholesterol, triglycerides and glycemia, demonstrating the ability to act at a cellular level. We noticed that deeper studies are required to understand how each shift in value takes place and why this is different in each Glycemia, Triglyceride and cholesterol molecule, thus making another study with a prolonged treatment is necessary to look at Quantec's efficacy through a constant modulation in the electromagnetic field. Future studies must examine whether the large-scale implementation of these techniques based on the circulatory and autonomous apparatus have a significant impact on reducing the mortality and morbidity risk in patients.

## Conclusion

Instrumental bioinformation using Quantec had an effective approach for reducing Blood Pressure, Glycemia, triglycerides and cholesterol and offers a new treatment method with encouraging results for those with cardiovascular risk factors.

## Acknowledgements

My thanks to Ricardo Yanasse, Jonathan Salinas, Cristiano Sales da Silva, Eduardo Chagas, Robison Quiterio and Sergio Areias for his crewmanship and made this research study possible.

## Conflict of interests

Authors declare that there is no conflict of interest.

## References

- Dimitropoulos G, Tahrani AA, Stevens MJ. Cardiac autonomic neuropathy in patients with diabetes mellitus. *World J Diabetes*. 2014;5(1):17–39.
- Guimarães AC, Lima A, Mota E, et al. The cholesterol level of a selected Brazilian salaried population: biological and socioeconomic influences. *CVD Prev*. 1998;1:3306–317.
- Lewington S, Clarke R, Qizilbash N, et al. Age specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 2002;360(9349):1903–13.
- Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics—2011 update: a report from the American Heart Association. *Circulation*. 2011;123(4):e18–e209.
- Vanderlei, Pastre CM, Hoshi RA, et al. Noções básicas de variabilidade da frequência cardíaca e sua aplicabilidade clínica. *Rev Bras Cir Cardiovasc*. 2009;24(2):205–17.
- Wild S, Roglic G, Green A, et al. Global prevalence of diabetes: estimates for year 2000 and projections for 2030. *Diabetes Care*. 2004;27(5):1047–53.
- Kimura T, Matsumoto T, Akiyoshi M, et al. Body fat and blood lipids in postmenopausal women are relating to retting autonomic nervous system activity. *Eur J App Physiol*. 2006;97(5):542–47.
- Kraus WE, Houmard JA, Duscha BD, et al. Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med*. 2002;347(19):1483–92.
- Liao D, Sloan RP, Cascio WE, et al. Multiple metabolic syndrome is associated with lower heart rate variability. *Diabetes Care*. 1998;21(12):2116–28.
- Must A, Spadano J, Coakley EH, et al. The disease burden associated with overweight and obesity. *JAMA*. 1999;282(16):1523–9.
- Zeiger, Bischof. The quantum vacuum and its significance in biology. In: Proceedings of the 3rd International Hombroich Symposium on Biophysics. Germany: Neuss; 1998.
- Carneiro A, et al. Biomagnetism: Instrumental Aspects and Applications. *Rev Bras Ens Fis* 2000; 22(3).
- Tansley D. La radiónica y la anatomía sutil del hombre. 1st ed. Malaga: Sirio S.A; 1972.
- Areias S. Bioinformação: O elo perdido da medicina. 1st ed. São Paulo: CPR; 2013.
- Buegner PV. Física y Tiempo de Ensueño. 2nd ed. Altkirchen: Alemanha; 2010.
- Burleson O, Schwartz E. Cardiac torsion and electromagnetic fields: The cardiac bioinformation hypothesis. *Med Hypotheses*. 2005;64:1109–16.
- Rüger M, Scheer F. Effects of circadian disruption on the cardiometabolic system. *Rev Endocr Metab Disord*. 2009;10(4):245–60.
- Sociedade Brasileira de Cardiologia, Sociedade Brasileira de Hipertensão, Sociedade Brasileira de Nefrologia. VI Diretrizes Brasileiras de Hipertensão. *Arq Bras Cardiol*. 2010;95(1 supl):1–51.
- Sociedade Brasileira de Cardiologia. *V Diretriz Brasileira de Dislipidemias e Prevenção a Aterosclerose*. *Arq Bras Cardiol* 2013.
- Sociedade Brasileira de Diabetes. Diretrizes da Sociedade Brasileira de Diabetes Brasil. 3rd ed. Itapevi, SP: A Araújo Silva Farmacêutica; 2009.
- Rajendra, Paul Joseph K, Kannathal N, et al. Heart rate variability: a review. *Med Bio Eng Comput*. 2006;44(12):1031–51.
- TASK FORCE of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Heart rate variability: standards of measurements, physiological interpretation and clinical use*. *Circulation* 1996;93:1043–65.
- Brennan M, Palaniswami M, Kamen P. Do existing measures of Poincare plot geometry reflect nonlinear features of heart rate variability. *IEEE Trans Biomed Eng*. 2001;48 (11):1342–47.
- Malliani A. The Pattern of Sympathovagal Balance Explored in the Frequency Domain. *News Physiol Sci*. 1999;14:111–17.
- Abdullah A. Coherence: A Novel Nonpharmacological Modality for Lowering Blood Pressure in Hypertensive Patients. *Global Adv Health Med*. 2012;1(2):56–64.
- Prinsloo GE, Rauch HG, Karpul D, et al. The Effect of a Single Session of Short Duration Heart Rate Variability Biofeedback on EEG: A Pilot Study. *Appl Psychophysiol Biofeedback*. 2013;38:45–56.