

Cannabis and the role of the endocannabinoid system in epilepsy treatment

Abstract

The aim of this review is to evaluate the effects, both beneficial and adverse, of medical cannabinoids; and the role of the endocannabinoid system in epilepsy treatment in adults and children. Despite the introduction of new antiepileptic drugs, the therapeutic response for patients with epilepsy remains poor; and the quality of life using such drugs has not substantially improved. Notwithstanding several advantages, new pharmacological antiepileptic drugs have not satisfactorily reduced the number of refractory patients. Cannabis has been used as a medicine for millennia. Current research has shown encouraging results for its use in the treatment of epilepsy. In cannabis, Δ^9 -tetrahydrocannabinol (THC) is the main psychoactive ingredient while cannabidiol (CBD) is the main non-psychoactive ingredient. THC is showing anticonvulsant properties in most animal models and many case reports. THC exerts its neural effects through the activation of CB1 receptors (CB1Rs) which are a part of the endocannabinoid system. Due to the psychotropic effects of THC, which have detrimental effects on cognition and behavior, its application for children and adolescents appear to be limited. To overcome this, the non-psychoactive compound, CBD, is being researched. CBD has shown anticonvulsant properties in numerous case reports, and in preclinical and clinical studies. However, CBD, as yet, does not have a completely clarified mechanism of action. Numerous case reports, preclinical studies and clinical studies indicate that CBD and CBD-containing medications are able to help manage epilepsy in adults and children affected by refractory seizures, such as occur in Dravet syndrome and Lennox-Gastaut syndrome. Clinical trials are limited however. Although the data is promising, further preclinical and clinical studies are needed to better evaluate the potential therapeutic profile of CBD in epilepsy.

Keywords: cannabidiol, childhood epilepsy, endocannabinoid system, epilepsy, refractory form of epilepsy, Δ^9 -tetrahydrocannabinol

Volume 11 Issue 6 - 2018

Inga Cicenaitė-Merz,¹ Nicholas A Kerna,^{2,3}
Tony L Brows⁴

¹Clear Lake Regional Hospital, Webster, Texas, USA

²College of Medicine, University of Science, Art and Technology, BWI

³Major General Hugh G Robinson Center for Medical Studies, USA

⁴Harvard University, USA

Correspondence: Nicholas A Kerna, College of Medicine, University of Science, Art and Technology, 7583 Sourdough Dr, Morrison, CO 80465, USA, Email nicholas.kerna@usat.edu

Received: May 27, 2017 | **Published:** December 06, 2018

Abbreviations: CBD, cannabidiol; CB1R, CB1 cannabinoid receptor; CB2R, CB2 cannabinoid receptor; CW, charlotte's web; GPDR, G-protein-coupled receptor; THC, Δ^9 -tetrahydrocannabinol; 2-AG, 2-arachidonoylglycerol

Introduction

According to the American Epilepsy Society, more than three million Americans live with epilepsy.¹ Availability of numerous antiepileptic drugs has not substantially reduced the proportion of patients with medically refractory epilepsy. Their quality of life and therapeutic response to currently available medications remains poor. Patients need new treatments that control seizures and have fewer side effects. This gap in epilepsy treatment—especially in refractory forms of childhood epilepsy—has led patients to seek alternative treatments. To this purpose, emphasis has been given to phytocannabinoid treatment which has become more accepted. Phytocannabinoids are naturally occurring cannabinoids derived from the cannabis plant. Researchers around the world are trying to understand how phytocannabinoids work, and how they might help in epilepsy.

Discussion

Cannabis sativa is one of world's most ancient medical plants. It has been used for over 5,000 years in many cultures for different therapeutic purposes. Marijuana has deeply American roots, too: George Washington, the first president of the United States, grew hemp at his estate at Mount Vernon. For most of America's history, cannabis

was legal; commonly found in tinctures and extracts.^{1,2} Marijuana's recreational use and abuse became common in Western societies in 1960s, approximately the same time scientists started to look for explanations of therapeutic and behavioral effects of Cannabis sativa. The major psychoactive ingredient of cannabis was first isolated in 1964, and then identified as Δ^9 -tetrahydrocannabinol (THC) in 1967.³ Another major chemical component, structurally related to THC but non-psychoactive, cannabidiol (CBD) was identified in 1963.⁴ These discoveries launched the search for the pharmacological effects of the chemical class of natural derivatives, phytocannabinoids. Now, over 500 compounds have been isolated from Cannabis sativa with approximately 105 being cannabinoids.⁵ Of those 105 compounds, THC has been determined as the primary constituent; it plays the central role in the discovery of a new signaling system in human body known as the endocannabinoid system. Two major cannabinoid receptors, CB1 (CB1R) and CB2 (CB2R), were identified as mediators of the biological effects of THC.⁶ In addition, two lipid molecules, anandamide and 2-arachidonoylglycerol (2-AG), are recognized as being the endogenous ligands for these G-protein-coupled receptors (GPDRs). These, together with several metabolic enzymes and endogenous cannabinoids, constitute the endocannabinoid system.⁷ Numerous detailed research studies of the last three decades have confirmed the fundamental role of the endocannabinoid system in multiple physiological and pathophysiological processes. Most research has aimed to manage the activity of endocannabinoid pathways for medical therapeutic purposes.

CB1R has now been mapped in many species through auto radiographic methods, Immunohistochemistry and in situ hybridization. CB1R is highly expressed in the brain, predominantly on neurons. The distribution of receptors is consistent between species and correlates well with the known behavioral effects of cannabinoids. Results show that cannabinoid receptor binding sites in the human brain are found mainly in the forebrain areas associated with higher cognitive functions; in the forebrain, midbrain and hindbrain areas associated with the control of movement; and in hindbrain areas associated with the control of motor and sensory functions of the autonomic nervous system. High levels of receptors are observed in the hippocampus; in basal ganglia output nuclei (globus pallidus and substantia nigra); the cerebellum; and cortical association regions; while moderate levels are observed in primary cortical regions, thalamic nuclei, striatum, and low levels in brain stem and spinal cord regions.⁸ Retrograde activation of presynaptic CB1R on axon terminals is now widely accepted as the general signaling mode by which a postsynaptic neuron can temporarily or continuously limit the strength of its afferent inputs.⁹ CB2Rs are located primarily on immune cells. Upon activation, they can modulate immune cell migration and cytokine release.⁸ Some neurons are expressing CB2Rs; however, the role of those neuro CB2Rs is not yet established.

More than three million Americans live with epilepsy. One-third of these people have ongoing treatment-resistant seizures.¹ The anecdotal reports of the positive effects of the marijuana plant for some individuals with epilepsy are giving hope for all epilepsy patients and, especially, those with treatment-resistant epilepsy. Modern medical literature is filled with case reports on surveys of different cannabis products. Those reports generally found that these products act as anticonvulsants in the majority of the cases. Proper randomized, double-blind, placebo-controlled trials are not permitted due to the fact that medical and recreational use of marijuana is still illegal in the many of the states in the U.S., and in most countries. Despite laws that permit use of medical marijuana in 40 states, cannabis is still illegal under federal law and is classified as a Schedule I drug. The federal government, therefore, views cannabis as highly addictive and having no medical value.¹⁰ Legal and ethical issues are making epidemiological studies on cannabis with high content of THC almost impossible. For now, researchers should consider present animal studies and case reports as valuable and useful sources of information regarding the potential benefits and limitations of cannabis treatment of seizures. These studies can indicate directions for cannabis research in near future, especially when considering the strong association between high THC/low CBD-containing cannabis and the chances of first time psychosis.¹¹

Another important consideration in this type of research is how long-term cannabis use for seizure control may affect the nervous system in epileptic patients upon cessation of marijuana smoking. A first, well-documented, case study described the reappearance of seizures in two patients whose focal epilepsy had previously been controlled by medical marijuana, directly after stopping marijuana use. A dramatic increase in seizure frequency together with underlying abnormal electrical brain activity was documented with electroencephalogram video telemetry.¹²

Due to the high prevalence of childhood epilepsy, it is a very important to research antiepileptic effects of cannabis. Russ et al. performed cross-sectional analysis of data from the National Survey of Children's Health (2007) on 91,605 children ages birth

to 17years, including 977 children reported by their parents to have been diagnosed with epilepsy/seizure disorder.¹³ Results showed that estimated lifetime prevalence of epilepsy/seizure disorder was 10.2 per 1000 (95% confidence interval) or 1%; and current epilepsy/seizure disorder was 6.3 per 1000 (95% confidence interval), or 0.6%.¹³ Based on these findings, it is estimated that 466,000 patients in the US and 765,000 childhood patients in Europe are currently diagnosed with epilepsy. Those numbers, together with the large numbers of refractory forms of epilepsy, make it imperative to find safe and efficient cannabinoid-based treatments. However, the concept of long-term administration of cannabis with higher concentration of THC is not favorable due to susceptibility in the developing prenatal, postnatal, and adolescent brain.¹⁴ For instance, THC was reported to impair short-term memory and learning abilities, to cause neuronal cell death of the hippocampus after chronic exposure, and to increase the risk of developing psychotic disorders with maturity.¹⁵ Therefore, research efforts were focused on using cannabis enriched in CBD. This cannabis preparation showed anticonvulsant effects in multiple animal studies and in several preclinical and some clinical studies.¹⁶ The antiepileptic mechanisms of CBD are not known; it has little affinity for either CBR1 and CBR2,¹⁷ but it has high antagonist effects at CBR1 and CBR2.¹⁸ However, this mechanism of CBD action cannot explain its antiepileptic properties because CBR1 activation is a primary protective measure against seizures.¹⁹ Moreover, administration of low doses of THC or administration of synthetic cannabimimetics, are all anticonvulsant in vitro and in vivo; and these effects can be blocked by CBR1 antagonists. When applied alone, high doses of CBR1 antagonists have usually been shown to increase seizure duration and frequency.¹⁵ Mechanism of action of CBD may include effects on the equilibrative nucleoside transporter; the orphan G-protein-coupled receptor GPR55; the transient receptor potential of melastatin type 8 channel; the 5-HT_{1A} receptor; the α 3 and α 1 glycine receptors; and the transient receptor potential of ankyrin type 1 channel. CBD has neuroprotective and anti-inflammatory effects.²⁰ It is not entirely clear how CBD works to control seizures; more research needed.

With the increasing legal availability of medical marijuana in some U.S. states, parents have started to use CBD cannabis preparations as an adjunctive therapy for their children with very serious, debilitating forms of childhood epilepsies. One of Colorado's biggest cannabis success stories is the recent case report of Charlotte Figi, who suffers from Dravet syndrome (a severe form of epilepsy caused by a mutation in the SCN1A gene encoding a voltage-gated sodium channel subunit). Charlotte's seizures started at age 3months as a prolonged status epilepticus. She had frequent febrile and afebrile status epilepticus as well as tonic, tonic-clonic, and myoclonic seizures. She began losing milestones, had significant cognitive and motor delays, required a feeding tube for nutrition and water, struggled to walk and talk, and required full assistance with her activities of daily living. At one point, she was experiencing up to 50 generalized tonic-clonic seizures per day. After unsuccessful treatment including all conventional anticonvulsant drugs (levetiracetam, oxcarbazepine, topiramate, zonisamide, valproate, clobazam, clonazepam, and valium) and the ketogenic diet, her parents tried CBD enriched/low THC content cannabis oil made from a specific cannabis strain, now named after her as Charlotte's Web (CW). This extract, slowly titrated over weeks and given sublingually in addition to her existing antiepileptic drug regimen, reduced Charlotte's seizure frequency to 2–3 nocturnal convulsions per month. This effect persisted for the

20 months, and Charlotte was successfully weaned from her other antiepileptic drugs. Then, her parents started to slowly decrease her CW and they reached 2 mg of CBD/lb per day (from her steady dose of 4 mg CBD/lb per day). Charlotte's seizures started coming back; when she was completely off the CW, her seizures returned to 5–10 seizures per day for 3 days, at which time parents restarted CW. Now, Charlotte is feeding and drinking orally and on her own, and her autistic behavior has lessened. She is finally able to walk and talk. Her seizure frequency is 2–3 nocturnal convulsions per month, a level consistently maintained over the last 3 years.²¹

A survey organized by Stanford University, among parents who are using CBD cannabis preparations to treat their children with epilepsy, reported significant improvement in their children. Nineteen children participated in this survey; they ranged in age from 2 to 16 years. Thirteen children had Dravet syndrome, 4 children had Doose syndrome (epilepsy with myoclonic astatic seizures), and one each had Lennox-Gastaut syndrome (multiple different types of seizures, particularly tonic and atonic) and idiopathic early-onset epilepsy. In all cases, except one, the children experienced treatment-resistant epilepsy for more than 3 years before trying CBD-enriched cannabis. The doses of CBD the parents reported providing ranged from less than 0.5 mg/kg/day to 28.6 mg/kg/day, whereas THC dosage was below 0.8 mg/kg/day in every case. The duration of CBD-enriched cannabis administration ranged from two weeks to over one year. Sixteen (84%) of the 19 parents reported a reduction in their child's seizure frequency. Two parents reported that their child became seizure-free after more than 4 months of CBD-enriched cannabis use. Three parents reported no change. Beneficial effects of CBD-enriched cannabis, other than reduced seizures, included better mood (15/19, 79%), increased alertness (14/19, 74%), better sleep (13/19, 68%) and decreased self-stimulation (6/19, 32%). Negative side effects included drowsiness (7/19, 37%) and fatigue (3/19, 16%).²²

The new trend in medical cannabis use in children with epilepsy is raising significant safety issues. Cannabis extracts are often inaccurately labeled, and can contain highly variable levels of CBD and THC. Medical application of cannabis poses risks due to a lack of standardization and regulation, inexact dosing, possible adverse effects, and medication interactions. More valid data regarding use of cannabis is coming from randomized clinical studies of CBD-containing medications. A new drug Epidiolex (GW Pharmaceuticals) contains about 98% CBD with only trace amounts of other cannabinoids, and is approved as an investigational drug. In 2015, very promising results for Epidiolex were reported at the American Epilepsy Society Annual meeting.²³ In a prospective observational study, data was obtained from 25 children with drug-resistant epilepsy. Patients were started on 5 mg/kg/day of CBD, and titrated weekly by 5 mg/kg increments until a maximum dose of 25 mg/kg/day, in addition to anti-epileptic drug therapy. The addition of CBD treatment was associated with more than a 50% seizure reduction in one-third of the patients at 3 months. This improvement in seizure control was maintained in 40% of patients for 12 months. The best results were obtained in Dravet syndrome patients. Twelve patients discontinued CBD due to lack of efficacy, and one patient discontinued due to a significant increase in seizures while on CBD treatment.²³

More hope to patients with Lennox-Gastaut Syndrome is given by results of a phase 3 randomized, double-blind placebo-controlled trial on the same drug Epidiolex; announced in June 2016. Data was obtained from 171 patients over a 14-week treatment period. Patients

were age 2–55; the average age of trial participants was 15 years, and 34% were 18 years or older. Median reduction in monthly drop seizures of 44% was reported, compared with a reduction of 22% in placebo patients; the difference between treatments was statistically significant ($p=0.0135$).²⁴ In addition, in March 2016, GW Pharmaceuticals announced positive results of the first pivotal phase 3 trial of Epidiolex in Dravet syndrome.²⁴

Conclusion

Some evidence suggests that cannabis and cannabinoid-based medicines may be beneficial in some patient populations with epilepsy, but may not be effective in other populations; and aggravation of epileptic seizures can occur in some cases. Further research is needed. The American Epilepsy Society has stated, "At present, the epilepsy community does not know if marijuana is a safe and effective treatment. Nor do we know the long-term effects that marijuana will have on learning, memory, and behavior, especially in infants and young children." The American Epilepsy Society has also urged the U.S. Drug Enforcement Agency (DEA) to change marijuana from its current Schedule I status to allow researchers to conduct studies more efficiently.²⁵

Acknowledgments

None

Conflicts of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. American Epilepsy Society. AES position on cannabis as a possible treatment for epileptic seizures. 2016.
2. National Geographic. 2015.
3. Mechoulam R, Gaoni Y. The absolute configuration of δ^1 -tetrahydrocannabinol, the major active constituent of hashish. *Tetrahedron Letters*. 1976;8(12):1109–1111.
4. Mechoulam R, Shvo Y. Hashish-I: the structure of cannabidiol. *Tetrahedron Letters*. 1963;19(12):2073–2078.
5. Husni AS, McCurdy CR, Radwan MM, et al. Evaluation of phytocannabinoids from high potency using bioassays to determine structure-activity relationships for cannabinoid receptor 1 and cannabinoid receptor 2. *Med Chem Res*. 2014;23(9):4295–4300.
6. LA Matsuda, SJ Lolait, MJ Brownstein, et al. Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature*. 1990;346(6284):561–564.
7. Blankman JL, Cravatt BF. Chemical probes of endocannabinoid metabolism. *Pharmacological Reviews*. 2013;65(2):849–871.
8. Glass M, Dragunow M, Faull M. Cannabinoid receptors in the human brain: A detailed anatomical and quantitative autoradiographic study in the fetal, neonatal and adult human brain. *Neuroscience*. 1997;77(2):299–318.
9. Castillo PE, Younts TJ, Cha'vez AE, et al. Endocannabinoid signaling and synaptic function. *Neuron*. 2012;76(1):70–81.
10. Americans for safe Access. Advancing Legal Medical Marijuana. Therapeutics and Research. Federal marijuana Law; 2016.

11. Di Forti M. Proportion of patients in south London with first-episode psychosis attributable to use of high potency cannabis: A case-control study. *Lancet psychiatry*; 2015.
12. Hegde M, Santos-Sanchez C, Hess CP, et al. Seizure exacerbation in two patients with focal epilepsy following marijuana cessation. *Epilepsy and Behavior*. 2012;25(4):563–566.
13. Russ SA, Larson K, Halfon N. A national profile of childhood epilepsy and seizure disorder. *Pediatrics*. 2012;129(2):256–264.
14. Hadland SE, Knight JR, Harris SK. Medical marijuana: review of the science and implications for developmental-behavioral pediatric practice. *J Dev Behav Pediatr*. 2015;36(2):115–123.
15. Rudenko V, Rafiuddin A, Leheste JR, et al. Inverse relationship of cannabimimetic (R₁)WIN 55, 212 on behavior and seizure threshold during the juvenile period. *Pharmacol Biochem Behav*. 2012;100(3):474–484.
16. Jones NA, Glyn SE, Akiyama S, et al. Cannabidiol exerts anti-convulsant effects in animal models of temporal lobe and partial seizures. *Seizure*. 2012;21(5):344–352.
17. Bisogno T, Hanus L, De Petrocellis L, et al. Molecular targets for cannabidiol and its synthetic analogues: effect on vanilloid VR1 receptors and on the cellular uptake and enzymatic hydrolysis of anandamide. *Br J Pharmacol*. 2001;134(4):845–852.
18. Thomas A, Baillie GL, Phillips AM, et al. Cannabidiol displays unexpectedly high potency as an antagonist of CB1 and CB2 receptor agonists in vitro. *Br J Pharmacol*. 2007;150(5):613–623.
19. Citraro R, Russo E, Ngomba RT, et al. CB1 agonists, locally applied to the cortico-thalamic circuit of rats with genetic absence epilepsy, reduce epileptic manifestations. *Epilepsy Research*. 2013;106(1-2):74–82.
20. Devinsky O, et al. Cannabidiol: Pharmacology and potential therapeutic role in epilepsy and other neuropsychiatric disorders. *Epilepsia*. 2014;55(6):791–802.
21. Maa E, Figi P. The case for medical marijuana in epilepsy. *Epilepsia*. 2014;55(6):783–786.
22. Porter BE, Jacobson C. Report of a parent survey of cannabidiol-enriched cannabis use in pediatric treatment-resistant epilepsy. *Epilepsy Behav*. 2013;29:574–577.
23. Oldham M, Sullivan J, Singhal N, et al. Long term efficacy and tolerability of add-on cannabidiol for drug resistant pediatric epilepsies. American Epilepsy Society annual meeting abstract; 2015;2.296.
24. GW Pharmaceuticals. GW Pharmaceuticals Announces Positive phase 3 Pivotal Trial Results for Epidiolex (cannabidiol) in the Treatment of Lennox-Gastaut Syndrome; 2016.
25. American Epilepsy Society AES. AES Position on Medical Marijuana. 2016.