

Active compounds in EAhy 926 endothelial cells, investigating cashew extract and molecules as natural health product approach to reducing the formation of advanced glycation endproducts

Abstract

Cashew, *Anacardium occidentale* Linn of the family Anacardiaceae, has been documented as traditional plant treatment for diabetes and hypertension. In our previous studies using *A. occidentale* leaf and seed extracts, we demonstrated the effective improvement in renal or testis function and reduced lesions associated with diabetic state in STZ-diabetic rats and stimulating glucose transport in C2C12 muscle cells.¹⁻³ We then wondered what molecular compounds were involved in these potent anti-diabetic activities. In the present study, using antioxidant activity-guided fractionation (DPPH advanced glycation endproducts (AGEs) essays), four extracts (E1 to E4) of *A. occidentale* leaves were studied for possible antioxidant capacities and vascular activity. We also compared the protective effects of cashew leaf extracts against high glucose mediated toxicity in cultured bovine aortic endothelial cells (EAhy 926 cells). Total antioxidant activity of the extracts of *A. occidentale* and identified compounds was determined using DPPH free radical scavenging assay. Two extracts of cashew's leaves, E3 and E4, were found to possess significant DPPH radical-scavenging activity with equivalent trolox values of 218±8 and 760±3 mg trolox/g dried extract, respectively. The order of the radical scavenging potency is E4>E3>E2>E1.

In vitro AGEs generation after 24h revealed that E3 had the strongest inhibitory activity against AGEs formation and its IC₅₀ value was 0.070mg/ml compared to that of aminoguanidine as standard (0.826mg/ml). A significant ($p<0.05$) increase in fluorescence of AGEs in 25.0mM glucose (HG) was partially corrected by E3, E4 and agathisflavone (AF) ($p<0.05$) vs. 25mM glucose alone. Additionally, cotreatment with E3, E4 or AF significantly attenuated HG induced EAhy as judged by overexpression of Mn-SOD and NF-kB.

These results suggest that *A. occidentale* MetOH extracts together with its principal compound, identified as agathisflavone reduces high glucose-induced toxicity by attenuating AGEs, MnSOD, NF-kB generation in endothelial cells, and therefore may be of significant therapeutic benefits against oxidative stress in diabetic complications and hypertension.

Keywords: *Anacardium occidentale*, leaves extracts, hyperglycaemic conditions, glycation, endothelial cells, oxidative stress

Volume 12 Issue 1 - 2019

Leonard Tedong,¹ Desire Paul D Dzeufiet,² Danielle Bilanda,² SD Sockeng FM Kanmangne,¹ PD Chuisseu,¹ P Haddad,³ Andrea Rosanoff,^{4,5}

¹Department of Physiology, High Institute of Health, University Cameroon, Cameroon

²Department of Animal Biology and Physiology, University of Yaounde, Cameroon

³Department of Pharmacology, University of Montreal, Montreal, Canada

⁴Department of Animal Biology and Physiology, University of Ngaoundere, Cameroon

⁵Center for Magnesium Education & Research Pahoehoe, USA

Correspondence: Leonard Tedong, Department of Physiology, High Institute of Health, University "des Montagnes", Bangante, Cameroon, Tel +237 75 89 76 09, +237 691 144 920, Email ltetong@yahoo.fr

Received: December 28, 2018 | **Published:** February 25, 2019

Introduction

Diabetes is the most common non-communicable diseases worldwide.⁴ It has become one of the leading causes of morbidity and mortality in developed countries, and the management of hyperglycaemia remains a major therapeutic concern. In the fourth edition of International Diabetes Federation Atlas,⁴ it has been stated that the estimated diabetes prevalence for 2010 has risen to 280million, representing 6.6% of the world's population. Although diagnosed on the basis of hyperglycaemia, diabetes mellitus is associated with broad metabolic abnormalities that contribute to microvascular and macrovascular complications.⁵ Studies based on cultured endothelial cells have clearly shown that endothelial cell growth is inhibited by hyperglycemia.⁶ The mechanism of glucose inhibiting endothelial cell growth has not been clearly defined but may be related to generation of Reactive Oxygen Species (ROS). Previous studies have shown

that diabetes mellitus increases oxidative stress through a number of mechanism. First, hyperglycaemia induces lipid peroxidation; second, glucose can undergo auto-oxidation via numerous intermediates that can react with proteins to generate Advanced Glycation End-products (AGEs) leading to gradual deterioration in the structure and function of tissue proteins and contributing to diabetes complications.⁷ AGEs can induce ROS production during oxidative stress and inflammatory state such that key enzymes that catabolise ROS (SOD, catalase and glutathione) and antioxidants such as Vitamine E, Vit C and carotenoids are reduced in patients with diabetes.⁸ Patients treated with antihyperglycemic agents often fail to reach glycemic normalization or, once achieved, do not maintain these goals long term.⁹ Great effort has been devoted to searching for antioxidants without toxicity and side effects, e.g. traditional crude drugs, medicinal prescriptions and functional foods. To that, various types of agents have been developed,

but medicines that completely ameliorate diabetes mellitus have not yet been founded. The specie *Anacardium occidentale* Linn (*A. occidentale*) commonly known as cashew tree, belongs to the family Anacardiaceae which includes the mango, *Mangifera indica* L., and the *Spondias* genus. Cashew plant is cultivated for its edible fruit and nut in addition to its traditionally medicinal values. The economic importance of this special tree is such that it is now widely distributed throughout the tropics, particularly in many parts of Africa and Asia. Cashew seeds are rich in flavonoids and monounsaturated fatty acids such as linoleic acid (Omega-3) which is associated with a lower prevalence of coronary artery disease.¹⁰ It is also reported that the cashew nut extract and some isolated compounds have anti-oxidant properties.¹¹ The cashew tree also produces a pseudo-fruit called the cashew apple that is commonly consumed in the northern Cameroon. Studies on the biological activities of fresh cashew apple juice (CAJ) have shown that not only does it exhibit antibacterial and antitumor activities; it also has antioxidant and antimutagenic properties.¹² Previously, studies have demonstrated anti-diabetic effects of *A. occidentale* leaf, stem-bark and seed extracts.^{13,14} Other studies in our research team also reported the ability of extracts of *A. occidentale* leaves to provide pancreatic protection against streptozotocin-induced diabetes in rats.¹³ More recently, we demonstrated a stimulating effect of cashew seeds extract on glucose transport in C2C12 muscle cells.³

To date, there have been no studies of the effect of *cashew's* leaf extracts and related compounds on protein glycation inhibitory activity and oxidative stress induced by hyperglycemia. Therefore, the aim of the present study was: (1) to search for potential antioxidant and anti-AGEs effects of *A. occidentale's* leaf extracts for application in the treatment of diabetes and hypertension (2) to determine whether treatment with *A. occidentale's* extracts may improve high glucose-induced endothelial dysfunction and if so, to investigate the signalling pathway involved.

Materials and methods

Reagents

The human EA.hy 926 endothelial cells lines were purchased from American type culture collection (University of North Carolina, USA). Dulbecco's modified Eagle's medium (DMEM) containing 25mM (4.5g/l) glucose, Foetal Bovine Serum (FBS), Trypsin EDTA, were obtained from Lonza (Verviers, Belgium). The following chemicals: 4-hydroxy-tempo, L-glucose, DPPH (1, 1-diphenyl-2-picrylhydrazyl) as Nitrate/Nitrite assay kits, superoxide anion assay kits, glutathione assay kits, monoclonal antibody (anti-catalase) were purchased from Sigma Aldrich Co. (St. Louis, MO, USA).

Plant material

Anacardium occidentale were collected in May 2007 in Garoua (Northern Province), Cameroon. The plant samples were identified in the National Herbarium Cameroon (NHC), Obili-Yaounde, Cameroon. A voucher specimen of the collected plant sample was also deposited in the herbarium and given a specimen number 65604/NHC. Leaves were separated and dried under shade for three days. Dried leaf samples were ground into a uniform powder using a blender.

Preparation of extracts

880g of the dried powdered Cashew tree leaf samples were taken separately in a paper cone and placed into Soxhlet apparatus. Cyclohexane (4L), an organic solvent was taken in the round bottom flask

attached to the Soxhlet apparatus. A condenser was attached to this set up (with a temperature set in the range of 25-30°C). The whole set up was placed on a heating mantle. Cyclo-hexane gets vaporized and rises up to the condenser where it condenses back into liquid. This process was continued until maximum extraction from the plant by cyclohexane. The same procedure was repeated with three other solvents: dichloromethane (DCM), ethyl acetate EtOAc, and methanol (MeOH). The extracts from each of the above overnight evaporations were stored in vials.

Compound isolation & identification

Analytical TLC followed by flash column chromatography of EtOAc and MeOH extracts was carried out on pre-cut plates (Si gel F 254 from EM science). The structures of compounds were identified and confirmed by their TLC profiles as well as their HPLC and NMR spectra.

In vitro studies

Free-radical scavenging activity: The potential antioxidant activity of cashew leaf extracts and isolated compounds from E3 and E4 were assessed on the basis of scavenging activity of stable DPPH (1-phenyl-2-picrylhydrazyl) radical free radical. In its radical form, DPPH has an absorption band at 520nm which disappears upon reduction by antiradical compound(s). The reaction mixture (3.15mL) contained 3mL of daily prepared DPPH solution (0.025g/L) and different concentrations of cyclo-hexane, DCM, EtOAc and MeOH extracts (E1, E2, E3 and E4, respectively) of *A. occidentale* or tested compounds dissolved in methanol. After 30minutes at room temperature, the absorbance was recorded at 520nm. The DPPH concentration in the reaction medium was calculated from a calibration curve analysed by linear regression. The percentage of disappeared DPPH was calculated as follows: $(1 - [\text{DO}]_{\text{test group}} / [\text{DO}]_{\text{reference without any antioxidant}}) \times 100$. A Trolox calibration curve was constructed by measuring the reduction in absorbance of the DPPH solution in the presence of different concentrations of Trolox (0–400μM). The DPPH radical scavenging activity of the extracts was measured using the Trolox standard curve and results were expressed as mg Trolox equivalent (TE) antioxidant capacity per g dried plant sample. Measurements were performed at least in triplicate; means and S.D. were calculated. Rosmarin and chlorogenic acid were used as positive controls.

Chemical determination of total AGEs by fluorescence measurement: A Maillard fluorescence-based assay was developed and optimized for screening of crude extracts (E1, E2, E3 and E4) and chemical compounds (Agathisflavone) that are able to inhibit the formation of AGEs. The assay involved incubating BSA (10mg/mL) with D-ribose (0.5M) and the tested compounds (3.10-6 to 3.10-3M) or extracts (10-6 to 1mg/mL) in a phosphate buffer 50mM pH 7.4. Solutions were incubated in microplate (96 wells) at 37°C for 24hours in a closed system. AGEs fluorescence (excitation: 370nm; emission: 440nm) was measured using a spectrofluorimeter Infinite M200 (Tecan, Lyon, France). To avoid quenching phenomena, the fluorescence resulting from the incubation in the same conditions of BSA (10mg/mL) and the tested compounds or extract (10-6 to 1mg/mL) was subtracted for each measure. Negative control, indicated as 100% inhibition of AGEs formation consisted of wells with only BSA. Positive control, with no inhibition of AGEs formation consisted of wells with BSA and D-ribose. The final volume assay was 100μL. The percentage inhibition of AGEs formation was = $[1 - (\text{fluorescence}$

of test group/fluorescence of the control group)] x 100% (Wu and Yen, 2005). The IC₅₀ (defined as concentration of 50% inhibition AGEs) values for each sample were derived (??). From dose-response curves using Microsoft-Excel computer software. Tests were performed in triplicate.

Cell culture and treatment: Regarding the high antiradical, the anti-glycation and the vasorelaxant activity of *A. occidentale* extracts, we decided to continue our investigation with E3 and E4 on cell culture. EAhy 926 cells were cultured at a density of 1x10⁶ per 75-cm² flask (Lifes Sciences, Croning, NY, 14831, USA) in DMEM (Dulbecco's modified Eagle's medium: Ham's F-12, 1:1; Lonza) supplemented with 1% L-glutamine, 1% non-essential amino acids, 1% Na-pyruvate, 1% streptomycin/penicillin (Lonza, Belgium), 1% hypoxanthine, aminopterin, thymidine (Sigma-Aldrich) and 10% of heat-inactivated foetal bovine serum (Invitrogen, Cergy Pontoise, France). The culture was maintained at 37°C in a humidified incubator gassed with 5% CO₂.

Protection of EAhy 926 cells from glucose toxicity induced advanced glycosylation end-products (AGEs). After reaching confluence, EA.hy 926 cells were seeded at a density of 1.5x10⁵ cells/well in 6-well tissue culture plates (Lifes Sciences, Croning, NY, 14831, USA). To test for cytoprotective activities against glucose toxicity, viability assays were performed on EA.hy 926 cells subjected to chronically elevated glucose in the presence of extracts or vehicle (0.1% DMSO). Cells were seeded in 96-well plates at a density of 6.25x10³ cells/well and cultured for 24h at 37°C. Complete medium was then replaced with serum-free medium adjusted to 25mM glucose (approx. 5-fold greater than normal concentration for this cell line) and supplemented with 0.025% BSA, with extracts or vehicle (DMSO) for 7 days. While these conditions feature a supra-physiological concentration of glucose (the concentrations observed in severe unmanaged diabetes can attain levels 7-fold greater than normal), they are necessary to induce apoptosis in approximately 40% of the cells over the 4-day experimental duration and to provide adequate experimental resolution. Under this paradigm, toxicity is due to glucose per se and not osmotic stress since the substitution of L-glucose by D-glucose abolishes toxicity.

Groups of cells were formed each one receiving the following fresh media every 48h: normal glucose medium (NG; 5mmol/L D-glucose), high glucose medium (HG; 25mmol/L D-glucose), or hyperosmotic control medium (5.5mmol/L D-glucose plus 19.5mmol/L L-glucose) for up to 7 days. In experiments, crude extract of *A. occidentale* leaves or isolated compounds solution, 100mg/ml and 100mmol/L respectively were prepared in 100% DMSO or water and stored at -20°C. Each solution was then diluted and added as supplemented to cell culture media containing high glucose. The final DMSO concentration in the medium was 0.1%. Experimental groups of cells were formed each one receiving EtOAc extract or MeOH extract of cashew leaves at doses of 7, 12.5 and 25µg/ml while isolated compounds groups received 5, 10µmol/l and 20µmol/l for 7 days. Control assays were performed without extracts, but equivalent volume of DMSO was added. When needed, pharmacological chemical, tempol (10mmol/l) were incubated in parallel group and used as positive control.

Effect of *A. occidentale* leaf extracts on cell viability: At the end of experiment, the cells were harvested after trypsinized with 0.05% trypsin (ml/well) for 5 min. DMEM (2ml/well) was added to neutralise the trypsin. The cells suspensions (20µl) were mixed with 0.2% trypan blue (20µl) and counted using a malassez chamber.

In another set of experiment, cell viability was determined using a 3-[4,5-dimethylthiazol-2-yl] K₂,5-diphenyl tetrazolium bromide (MTT), quantitative colorimetric method as previously described.¹⁵ The MTT assay was performed in triplicate for each drug used.

Western blotting: To investigate a possible antioxidant mechanism of cashew leaves's action, NF-κB, catalase and MnSOD expression was evaluated by Western blot analysis as we previously described.¹

Statistical analysis

All values are expressed as mean±SD. Statistical significance was determined using analysis of variance (ANOVA) followed by Tukey's test. P<0.05 was considered statistically significant.

Results

Chemical characterization and Identification of a flavonoid from EtOAc and MeOH extracts of cashew leaves

For leaves of *A. occidentale* extracted with EtOAc and MeOH, the HPLC "fingerprint" of the extracts showed major peaks at retention times (min) of 4.84, 21.41 and 27.90 at wavelength of 254nm (Figure 1A). Purification was achieved by a combination of polyamide and sephadex LH-20 columns. Figure 1B shows, as an example, the HPLC-UV chromatogram of a purified flavonoid extract obtained from the EtOAc extract of *A. occidentale*. Phytochemical and semi-preparative studies resulted in the isolation of two compounds: Agathisflavone, determined by UV, MS, and NMR spectra (Figure 1B & Figure 2), plus a second compound we named Q13 that was not completely identified (data not shown).




Figure 1A HPLC chromatogram of cashew leaves' EtOAc (Blue line) and MeOH extracts (Red line) (from Cameroon). Peak 30, Agathisflavone.

Chemical assays

Antioxidant activity of *A. occidentale* extracts using scavenging of 1,1-diphenyl-2-picrylhydrazyl (DPPH)

The radical scavenging activities of the extracts of *A. occidentale* were estimated using the Trolox standard curve and results were expressed as milligram Trolox equivalent (TE) antioxidant capacity per g dried plant sample. Except cyclohexane extract, all other extracts show free radical scavenging activity at different potencies. The highest scavenging activity of DPPH was observed for the MeOH extract, with 760±32mg trolox equivalent/g dried extract followed by the EtOAc and DCM extracts with values of 217±8 and 24±3mg

Trolox equivalent/g dried extract, respectively, compared to that of rosmarin, 64±3mg TE/g dried extract (Table 1).




Figure 1B HPLC chromatogram: UV absorption spectra of Agathisflavone.




Figure 2 Structure of biflavonoid, Agathisflavone major compound isolated from EtOAc extract leaves of *A. occidentale*.

Table 1 Inhibiting effect of different extracts of *A. occidentale* leaves and isolated compounds on free radical generation in vitro, compared to Rosmarin and chlorogenic acid

A occidentle leaves extracts 0.1g/L	mg Trolox equivalent/g dried extract	Compounds 50µM	TE/µg
Cyclohexane (E1)	12±1	Not detected	-
Dichloromethane (E2)	24±3	Not detected	-
Ethyl acetate (E3)	217±8	Agathisflavone	16±1
Methanol (E4)	760±3	Q13 (from E3 or E4)	642±3
Rosmarin	64±7	Chlorogenic acid	53±2

Inhibitory effect of *A. occidentale* extracts and isolated compounds on chemical AGEs formation

The formation of total AGEs was assessed by monitoring the production of fluorescent products at excitation and emission maxima of 440nm, as previously explained. Figure 3A & 3B shows the effects of *A. occidentale* leaf extracts and major identified compound

(Agathisflavone) on ribose-BSA AGEs formation during 24h of BSA incubation at 37°C. The fluorescence intensity of this glycoprotein which is characteristic of AGEs, was highly increased through incubation of BSA during 24h. The results showed that *A. occidentale* leaf extracts inhibited the AGEs formation in the following order: EtOAc>MeOH>DCM> Cyclohexane (Figure 3A). As it is evident in the same figure, EtOAc extract at different concentrations (10-100 mg/ml) has significantly quenched the fluorescence and the result is more pronounced than that of aminoguanidine, the positive control compound known as an inhibitor of glycation process (91% vs 80% at a concentration of 3mg/ml). The higher inhibitory effect of EtOAc extract is also evident with the IC₅₀ (defined as the concentration of extract or test compound required to produced 50% inhibition) value which is higher than that of Cyclohexane, DCM and MeOH or aminoguanidine (0.070 vs 0.374, 0.221, 0.153 and 0.826 mg/ml), Table 2. The effect of agathisflavone shown in Figure 3B demonstrated that this major compound isolated from EtOAc and MeOH extracts inhibited the fluorescence intensity induced by glycation process. Agathisflavone significantly suppressed fluorescence intensity more than did aminoguanidine in a dose-dependent manner; agathisflavone's IC₅₀ was 9.5x10⁻² µmol/l versus 516.9x10⁻²µmol/l for aminoguanidine, the positive control (Table 2).

Table 2 The IC₅₀ of *A. occidentale* leaves extracts and isolated compound on AGEs formation

Extracts	IC ₅₀ (mg/ml)	Compounds	IC ₅₀ x 10 ⁻² (µmol/l)
Cyclohexane	0.374	-	-
DCM	0.221	-	-
EtOAc	0.07	Agathisflavone	5.2
MeOH	0.153		-
Aminoguanidine	0.826	Aminoguanidine	0.096

In vitro bioassays

Anacardium occidentale leaf extracts or an isolated biflavonoid (Agathisflavone) protection against glucose induced advanced glycosylation end-products (AGEs) in EAhy 926 cells.

EA.hy 926 endothelial cells were cultured for 7days in experimental media, after which spontaneous fluorescence was measured as a non-specific marker of AGEs production. Fluorescence was increased more in 25.0mM glucose than in 5.5mM glucose (Figure 4) (p<0.05). This augmentation was partially corrected by both the EtOAc extract (E3) and the MeOH extract (E4) as well as agathisflavone (AF) (p<0.05) vs. 25mM glucose alone). The fluorescence levels obtained with E3, E4 and Agathisflavone added to 25mM glucose did not differ significantly between them.

Protection of Anacardium occidentale leaves extracts or an isolated biflavonoid (Agathisflavone) against cell proliferation in EAhy 926 cells.

EtOAc or MeOH extracts (E3 or E4) to a lesser extent protect against glucose-induced inhibition of cell proliferation (see above and Figure 4). Proliferation of endothelial cells (cell viability) exposed on 25mmol/l glucose for 7days was reduced to 78±1% of that in physiological (5.5mmol/l) glucose (Figure 5). In the presence of 7 and 12.5µg/ml of EtOAc extract of *A. occidentale* leaves (E3), proliferation of endothelial cells grown in 25mmol/l glucose was reduced to 91% and 93%, respectively, of that in cells grown in

5.5mmol/l glucose alone. MetOH extract (E4) at the doses of 7µg/ml and 12.5µg/ml increased the proliferation of endothelial cells to 84% and 89%, respectively. Agathisflavone, a major compound isolated from EtOAc t and MeOH extracts, was particularly effective in protecting against the anti-proliferative effects of high glucose. In the presence of 5µmol/l of agathisflavone and 5µmol/l of tempol (positive control), proliferation of endothelial cells grown in 25mmol/l of glucose for 7days was 93±5%, of that in cells grown in 5.5mmol/l glucose (Figure 6). The proliferation of cells grown in 5.5mmol/l of glucose was not different of that of cells grown in 19.5mmol of L-Glucose and 5.5mmol/l of glucose taken as osmotic control (results not shown).




Figure 3 A Inhibitory effect of *A. occidentale* extracts on fluorescence intensity of glycated BSA in terms of % AGEs inhibition. (B) Inhibitory effect of Agathisflavone on fluorescence intensity of glycated BSA in terms of % AGEs inhibition.




Figure 4 Fluorescence of advanced glycosylation end-products (AGEs) in EAhy 926 cells cultured in physiological or high glucose concentrations, supplemented with *A. occidentale* leaves extracts or an isolated biflavonoid.

Glu, D-Glucose; L, Glu; L, glucose; AF, Agathisflavone; AG, Aminoguanidine




Figure 5 Effect of *A. occidentale* leaves extracts on high glucose-induced loss of cell viability. Eahy cells were cotreated with the indicated concentrations of *A. occidentale* and high glucose (25mmol/l) for 7days.




Figure 6 Proliferation of endothelial cells cultured in 5.5 or 25mmol/l glucose, *A. occidentale* leaves extract (A) or isolated compound (B).

Effect of *Anacardium occidentale* leaf extracts or an isolated biflavonoid (Agathisflavone) on high glucose-induced loss of EAhy 926 cells viability. Cytotoxicity assay based on MTT revealed that none of the extracts of plant or identified compound at the doses tested was toxic to the EA.hy 926 endothelial cells after 7days exposure (data not shown).

Total cells lysate nitrates/nitrites

As shown in Figure 7, exposure of endothelial cells to hyperglycaemic (25mM) conditions for 7days caused a significant ($p<0.05$) increase of the level of total nitrates/nitrites in cell lysate compared to normoglycemic control group (13.17 ± 0.46 vs $4.95\pm0.46\mu\text{mol/ml}$). Co-treatment of hyperglycaemic-exposed cells with *A. occidentale* leaf ethyl acetate extract (E3) or methanol extract (E4) for 7days significantly ($p<0.05$) reduced the total level of nitrates/nitrites compared with the hyperglycemic group by 27.7% and 33.1% respectively. When L-glucose (19.5mM) was used as osmotic control, EA.hy cells did not show any increase in total nitrate/nitrite. The 15% and 21% reduction of total nitrates/nitrites respectively with agathisflavone (identified compound) and non-identified compound (Q13) was not significant while tempol failed to reduce the total level of nitrate/nitrite in cell lysate (Figure 7).

Effects of EtOAc and MeOH extracts and their major compounds on hyperglycaemia induced on NF-kB, Mn-SOD and catalase expressions. *A. occidentale* leaf extracts and identified compounds at different concentrations, maintained in DMEM containing 5.5 or 25mM glucose for 7days, were harvested and NF-kB, Mn-SOD and catalase expression were measured.

High glucose treatment (25mM) but not 19.5mM L-Glucose markedly increased expression of NF-kB (125%) (Figure 8). E3 or Q13 cotreatment with HG, largely prevented NF-kB expression and its expression was 98 and 101% respectively.

High glucose treatment markedly enhanced EA.hy 926 endothelial cells' Catalase expression (by 1.37 fold) compared to normal glucose. HG effect on catalase expression was not attenuated by either Cashew's extracts (E3 and E4) or compounds isolated from these extracts.

MnSOD protein expression and activity were determined in EA.Hy926 cells cultured in high and normal glucose concentrations to determine whether hyperglycemia affects this enzyme and cellular defense mechanism against oxidative stress. Following 7days exposure to HG, the mitochondrial superoxide generation (MnSOD) were significantly elevated in comparison to the NG incubations (Figure 8). All treatment with plant extracts and compounds for 7days significantly attenuated the HG-induced superoxide generation.




Figure 7 Effect of glucose on total cells lysate nitrates/nitrites.

NG, Normoglycemic, 5.5mmol/l glucose; HG, Hyperglycemic, 25mmol/l glucose; control, NG+19.5mmol/l L-glucose (osmotic control). The levels of nitrate/nitrite cultured in NG, HG, control medium or different treatments (E3, *A. occidentale* leaves ethyl acetate extract, 12µg/ml; E4, *A. occidentale* methanol extract, 12µg/ml; AF, Agathisflavone 10µM; Q13, non identified fraction, 10µg/ml; T, tempol, 10µM) for 7 days were measured as described in the Materials and Methods. Results are expressed as mean±SE from three separate experiments.

*p<0.05 difference compared to NG group and #p<0.05 difference compared to HG group.





Figure 8 A Representative Western blots showing the protein expression of NFkB, catalase and Mn-superoxide dismutase (SOD) in EAhy endothelial cells cultured under NG, normoglycaemic (lanes 1; 5.5mmol/l) and HG, hyperglycaemic (25mmol/l glucose) conditions, L-Glu, control medium (NG+19.5mmol/l L-glucose) or different treatments (HG+E3, *A. occidentale* leaves ethyl acetate extract, 12µg/ml or +E4, *A. occidentale* methanol extract, 12µg/ml or AF, Agathisflavone 10µM or Q13, non identified fraction, 10µg/ml or T, tempol, 10µM) for 7 days. (B) Histogram showing the fold differences in protein expression (B1, Catalase; B2, NF-kB; B3, Mn-SOD) observed in EAhy cultured under hyperglycaemic vs normoglycaemic conditions or vs different treatments.

Discussion

Anacardium occidentale Linn (*A. occidentale*) is commonly cultivated for its edible fruits and nut which have high medicinal value particularly on diabetic and hypertension diseases. Diabetic state has been shown to enhance oxidative stress, however, very little is known in regards of *A. occidentale* antioxidant impact on this metabolic disease. In attempting to correct this, in the present study we demonstrated that extracts of *A. occidentale* leaves possess antioxidant properties. In addition, one identified compound (Agathisflavone) and one unidentified compound (Q13) isolated from these extracts showed similar properties.

Assays such as the scavenging of DPPH and anti-glycation by cyclohexane (E1), dichloromethane (E2), ethyl acetate (E3) and methanol (E4) extracts of *A. occidentale* leaves plus an identified compound (Agathisflavone) and an unidentified compound (Q13) isolated from these extracts were performed. Our data clearly established that E3 and E4 extracts of *A. occidentale* possessed significant DPPH radical-scavenging activity with Trolox equivalence values of 218 ± 8 and 760 ± 3 mg trolox/dried extract, respectively, and the DPPH scavenging value of E3 and E4 was significantly stronger than that of standard antioxidant, Romarin (64 ± 7 mg trolox/dried extract). Our results imply that there are abundant antioxidative phytochemicals present in the acetate and methanol extracts of *A. occidentale* leaves. This was confirmed by testing the scavenging of DPPH by Q13, a non-identified compound isolated from E3, which compared well to chlorogenic acid as standard (642 ± 3 vs 53 ± 2 mg trolox/dried extract).

In this study, synthetic DPPH was utilized to assess radical scavenging abilities of *A. occidentale* extracts in order to identify the potent extract among E1, E2, E3 and E4. This method is widely used for the determination of the radical scavenging abilities of plants. Although these radicals may not be biologically relevant, they provide an indication of hydrogen/electron-donating capacity of plants and are hence a useful measure of in vitro antioxidant activity.

Previous phytochemical and pharmacological investigations of *A. occidentale* leaves have shown the presence of myricetin, agathisflavone, robustaflavone, amentoflavone, quercetin, kaempferol, apigenin, quercetin 3-O-rhamnoside, and quercetin 3-O-glucoside. The biological properties of flavonoids have been focused on extensively because they are natural antioxidant products with little cytotoxicity. They have also been recognized to possess anti-inflammatory, antiviral and anti-carcinogenic antidiabetic and anti-hypertensive activities (Fujii et al., 2006). Flavonoids can exhibit their antioxidant activity in several ways: (i) radical scavenging activity (ii) prevention of the transition metal-catalyzed production of reactive species (iii) interaction with other antioxidants (such as cooperative actions), localization, and mobility of antioxidant at the microenvironment.¹⁶

Since *A. occidentale* extracts were potent on the radical scavenging abilities, we hypothesized that crude extracts with powerful antioxidant properties may inhibit Advanced Glycation End products (AGEs). To determine any link between antioxidant activity of *A. occidentale* leaf extracts and glycation and oxidation, we evaluated antiglycation activity of the 4 different extractions of *A. occidentale*'s leaves in vitro. Of the 4 extracts tested, the EtOAc fraction (E3) exhibited the strongest inhibitory activity against AGEs formation in the BSA-ribose assay, and its IC₅₀ value (0.070 mg/ml) was less

than one tenth that of aminoguanidine as standard (0.826 mg/ml) (Table 1). This anti-AGEs effect could be explained by the presence of Agathisflavone, the biflavonoid extracted and identified from the EtOAc (E3) extraction which exhibited a significant inhibitory effect on AGEs formation compared to the E3 crude extract. This effect was more potent than that of aminoguanidine as a standard. Our results also demonstrated that protein glycation and oxidation could be inhibited effectively when flavonoids were added during the glycoxidative process in endothelial cells. AGEs breakers may help to provide a protective effect against hyperglycaemia-mediated protein damage and subsequently improve the pathophysiology of diabetes and its complications.¹⁷

In the second set of our study, we used endothelial cells to characterize metabolic disturbances that occur following chronic exposure to high levels of glucose. We then appreciated the abilities of the EtOAc and MeOH extracts and the principle identified compound to protect endothelial cells against high glucose and AGE-induced cellular toxicity. To demonstrate glucose induced inhibition of cell proliferation in vitro we used EA.hy 926 endothelial cells which have been recognized to have higher sensitivity to ox-LDL-induced injury associated with weaker enzymatic antioxidant defences.¹⁸ Circulating concentrations of glucose can range from around 10 mmol/l to >50 mmol/l in severely untreated diabetic patients. For an in vitro model system of vascular endothelial cells, we chose glucose concentrations from 5.5–25 mmol/l to represent a range corresponding to normal and diabetic conditions. The normal growth medium was supplemented with an additional 25 mmol/l for the majority of experiments as this concentration induced a reduction of proliferation activity of endothelial cells. This concentration is similar to that used in other similar studies investigating the effects of glucose on protein glycation and oxidative stress in macrovessel and microvessel endothelial cells.¹⁹ In our study, E3 and E4 were both effective in protecting cells against glucose- and AGE-mediated toxicity except 25 µg/ml of E3. On the other hand, coinubation of identified compound agathisflavone with high glucose for 7 days showed proliferation of endothelial cells (93%) higher than that of cells grown in high glucose alone (71%). In agreement with other studies, it has been demonstrated that our isolated and identified biflavonoid, Agathisflavone, also increased cell survival and RA-induced neuronal differentiation in pluripotent stem cells.²⁰ The flavonoid isolated from *A. occidentale* leaf extracts, namely Agathisflavone, should be added to the list of other 10 flavonoids isolated from *P. linteus* which have been shown by Lee et al.,²¹ to protect cultured endothelial cells following exposure to bovine serum albumin derived advanced glycation endproducts.

It could thus be of great health benefit to propose that administrations of naturally occurring flavonoids are beneficial for the prevention of protein glycation in high glucose environments such as diabetes. Our data on Agathisflavone's anti-glycation activities is in agreement of those of Manual et al.,²² who demonstrated that daflon 500, a clinical drug which is made up of flavonoids, had attenuated effect on HbA1C and protein glycation in a group of twenty eight Type 1 diabetic patients. This probably indicates that the *A. occidentale* extracts can either scavenge hydroxyl radicals as justified by the reduction of the total nitrate/nitrite observed after cotreatment of endothelial cells with high glucose and different treatment (E3, E4, Agathiflavone, Q13 and Tempol); the crude extracts could also act by chelating transition metals leading to less hydroxyl radical production, or they may have both effects. Increasing evidence suggests that glucose and AGEs, at

concentrations reflecting those found in the plasma of diabetic patients, exerts profound effects on vascular function, secondary to inhibition of endothelial cell NO generation.²³ Natural compounds, such as agathisflavone tested in our study, which have combined anti-glycation and antioxidant properties are likely to offer maximum therapeutic potential. The effect of high glucose levels on cellular metabolism and cell growth is likely to be mediated by intracellular signaling. We have founded in the current study that exposure endothelial cells culture to *A. occidentale* for 7days elicits significant increases in nuclear factor kB (NF-kB), Manganese Superoxide Dismutase (MnSOD) and Catalase expression. What is the link between all these components and diabetes? In diabetes, NF-kB is an oxidant-sensitive transcription factor, and hyperglycemia induced oxidative stress may play a key role in the pathogenesis of diabetic vascular disease. In this study, we demonstrated that prolonged hyperglycemia can lead to the formation of AGEs which could act through specific receptors on vascular cells to directly activate NF-kB. The nuclear factor kB in vascular smooth muscle cells can induce pathological changes in gene expression found in atherosclerosis and diabetic²⁴ complications. Removal of toxic oxygen metabolites is the putative function of antioxidant enzymes such as CuZnSOD, MnSOD, GPX and Catalase. To test whether E3 and E4 extracts and identified compound, Agathisflavone, impact the cell viability exposed endothelial cells to high glucose to mimic diabetic conditions.

Co-incubation of EAhy cells with E3 and E4 caused a significant decrease in the NF-kB and Mn-SOD expression in cells treated with high glucose. Agathisflavone, the identified compound, failed to alter proteins expression but the non-identified compound, namely Q13, significantly decreased NF-kB Catalase and MnSOD expression in cells treated with high glucose ($P < 0.05$ vs High Glucose). The reduction of the NF-kB expression by these cashew leaf extracts could be explained by the presence of Q13 (non- identified compound) in our extracts or other compounds such as anacardic acid which has been shown to suppresses expression of NF-kB-regulated gene products in cancer cells (Sun et al., 2006). Glucose autoxidation products can attach to specific receptors from the surface of endothelial cells and change their properties, contributing to early formation of atherosclerotic lesions.²⁵ Among the pathways studied, MnSOD appears to play the most crucial role in diabetic complications. In this study, we observed that MnSOD was not altered after EA.hy endothelial cells were expose to high glucose and different extracts (E3 and E4). MnSOD is an enzyme stimulating conversion of superoxide to hydrogen peroxide (H₂O₂) and previous studies have suggested that H₂O₂ plays a central role in NF-kB activation in coronary artery endothelial cells. Hyperglycemia-induced superoxide generation might also favour increased expression of iNOS through the activation of NF-kB, which increases the generation of nitric oxide (NO).²⁶ It therefore appears reasonable to hypothesize that elevated expression and activity of antioxidant enzymes is an adaptive response of endothelial cells to meet the biological demand exerted by hyperglycaemic-induced oxidative stress.^{27–32}

Conclusion

Collectively, we have made novel observations in our present experiment. First, the MeOH extract of *A. occidentale* leaves is the most potent in scavenging the DPPH. Second, we found out that the EtOAc extract of *A. occidentale* leaves possesses strong glycation activities and the free radical scavenging of its isolated compound, Agathisflavone may contributes in part to their antiglycation effects.

The extracts of *A. occidentale* leaves and the principal compound identified also protect endothelial cells against glucose toxicity by attenuating High Glucose-induced MnSOD and NF-kB generation. In this regard, EtOAc and/or MeOH extract of *A. occidentale* leaves may protect endothelial cells against free radical-induce cell damage in diabetes complications.

Acknowledgments

This study was facilitated by the Grants commission of Association of Francophone Universities, Cameroon which provided us the postdoctoral fellowship for scientific research (N° 1001PP/224/BAC 2008). Authors gratefully acknowledge the Department of Integrated Neurovascular Biology (UMR CNRS 6214-INSERM 771) of University of Angers, France, for research support and material to carry out this project.

Conflicts of interest

Author declares no conflicts of interest.

References

1. Tedong L, Dimo T, Dzeufiet DPD, et al. Antihyperglycemic and renal protective activities of *Anacardium occidentale* (Anacardiaceae) leaves in Streptozotocin induced diabetic rats. *Afr J Traditional, Complementary and alternative Medicines*. 2006;3(1):23–35.
2. Tedong L, Dimo T, Dzeufiet DPD, et al. Effect of leaf *Anacardium occidentale* L. (Anacardiaceae) hexane extract on reproductive function in streptozotocin-induced diabetic rats. *Phototherapie*. 2007;5:182–193.
3. Tedong L, Padma M, Martineau LC, et al. Hydro-ethanolic extract of cashew tree (*Anacardium occidentale*) nut and its principal compound, anacardic acid, stimulate glucose uptake in C2C12 muscle cells. *Mol Nutr Food Res*. 2010;54(12):1753–1762.
4. International Diabetes Federation (IDF). 4th edn. IDF Atlas; 2009. p. 7.
5. Buse JB, Rosenstock J, Sesti G, et al. Liraglutide once a day versus exenatide twice a day for type 2 diabetes: a 26-week randomised, parallel-group, multinational, open-label trial (LEAD-6). *Lancet*. 2009;374(9683):39–47.
6. Curcio F, Ceriello A. Decreased cultured endothelial cell proliferation in high glucose medium is reversed by antioxidants: new insights on the pathophysiological mechanisms of diabetic vascular complications. *In Vitro Cell Dev Biol*. 1992;28(11-12):787–790.
7. Song P, Wu Y, Xu J, et al. Reactive Nitrogen species by hyperglycemia suppress Akt signalling and triggers apoptosis by upregulating phosphatase PTN (Phosphatase and tensin homologue deleted on chromosome 10) in an LKB1-Dependant manner. *Circulation*. 2007;116(14):1585–1595.
8. Marchetti Piero, Matthews David. Pancreatic β -cell in type 2 diabetes. Birth, life, failure and rescue. 1st edn. Oxford, UK: Ed Wiley-Blackwell; 2009. p 75–113.
9. Williams-Herman D, Johnson J, Teng R, et al. Efficacy and safty on combinaison therapy with sitagliptin and metformin in patients with type 2 diabetes: a 54-week study. *Cur Med Res Opi*. 2009;25(3):569–583.
10. Konan NA, Bacchi EM. Antiulcerogenic effect and acute toxicity of hydroethanolic extract from the cashew (*Anacardium occidentale* L.) leaves. *J Ethnopharmacol*. 2007;112(2):237–242.
11. Kamath V, Rajini PS. The efficacy of cashew nut (*Anacardium occidentale* L.) skin extract as free radical scavenger. *Food Chemistry*. 2007;103(1):428–433.

12. Melo-Cavalcante AA, Picada J N, Rubensam G, et al. Antimutagenic activity of cashew apple (*Anacardium occidentale* Sapindales, Anacardiaceae) fresh juice and processed juice (*cajuína*) against methyl methanesulfonate, 4-nitroquinoline N-oxide and benzo[a]pyrene. *Genet Mol Biol.* 2008;31(3):751–766.
13. Sokeng DS, Kamtchouing P, Watcho P, et al. Hypoglycemic activity of *Anacardium occidentale* L. Aqueous extract in normal and streptozotocin-induced diabetic rats. *Diab Res.* 2008;36:001–009.
14. Ojewole JA. Laboratory evaluation of hypoglycemic effect of *Anacardium occidentale* Linn (Anacardiaceae) stem-bark extracts in rats. *Methods Find Exp Clin Pharmacol.* 2003;25(3):199–204.
15. Gondwe M, Kamadyaapa DR, Tufts M, et al. Sclerocarya birrea [(A. Rich.) Hochst.] [Anacardiaceae] stem-bark ethanolic extract (SBE) modulates blood glucose, glomerular filtration rate (GFR) and mean arterial blood pressure (MAP) of STZ-induced diabetic rats. *Phytomedicine.* 2008;15(9):699–709.
16. Apak R, Güçlü K, Demirata B, et al. Comparative evaluation of various total antioxidant capacity assays applied to phenolic compounds with the CUPRAC assay. *Molecules.* 2007;12(7):1496–1547.
17. Ardestani A, Yazdanparast R. Inhibitory effects of ethyl acetate extract of *Teucrium polium* on in vitro protein glycoxidation. *Food Chem Toxicology.* 2007;45(12):2402–2411.
18. Claise C, Chalas J, Edeas M, et al. Comparison of oxidized low-density lipoprotein toxicity on EA.hy 926 cells and human vein endothelial cells: influence of antioxidant systems. *Cell Mol Life Sci.* 1997;53(2):156–161.
19. Padayatti PS, Jiang C, Glomb MA, et al. A high concentration of glucose induces synthesis of argpyrimidine in retinal endothelial cells. *Curr Eye Res.* 2011;23(2):106–115.
20. Paulsen BS, Souza CS, Chicaybam L, et al. Agathisflavone enhances retinoic acid-induced neurogenesis and its receptors α and β in pluripotent stem cells. *Stem Cells Dev.* 2011;20(10):1711–1721.
21. Lee YS, Kang YH, Jung JY, et al. Protein Glycation Inhibitors from the Fruiting Body of *Phellinus linteus*. *Biol Pharm Bull.* 2008;31(10):1968–1972.
22. Manuel y Keenoy B, Vertommen J, De Leeuw I. The effect of flavonoid treatment on the glycation and antioxidant status in Type 1 diabetic patients. *Diabetes Nutr Metab.* 1999;12(4):256–263.
23. Xu B, Chibber R, Ruggerio D, et al. Impairment of vascular endothelial nitric oxide synthase activity by advanced glycation end products. *FASEB.* 2003;17(10):1289–1291.
24. Sung B, Pandey MK, Ahn KS, et al. Anacardic acid (6-Nonadecyl Salicylic Acid), an inhibitor of histone acetyltransferase, suppresses expression of NF- κ B-regulated gene products involved in cell survival, proliferation, invasion and inflammation through inhibition of I κ B α kinase, leading to potentiation of apoptosis. *Blood.* 2008;111(10):4880–4891.
25. Goycheva P, Gadjeva V, Popov B. Oxidative stress and its complications in diabetes mellitus. *Trakia Journal of Sciences.* 2006;4(1):1–8.
26. Pacher P, Beckman J, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev.* 2007;87(1):315–424.
27. Cracowski JL, Devillier P, Durand T, et al. Vascular biology of the isoprostanes. *J Vasc Res.* 2001;38(2):93–103.
28. Dogne JM, Hanson J, Domenico P. Thromboxane, prostacyclin and isoprostanes: therapeutic targets in arterogenesis. *Trends Pharmacol Sci.* 2005;26(12):639–644.
29. Fujii H, Yokozawa T, Kim YA, et al. Protective effect of grape seed polyphenols against high glucose-induced oxidative stress. *Biosci Biotechnol Biochem.* 2006;70(9):2104–2111.
30. Kubo I, Muroi H, Himejima M, et al. Structure-antibacterial activity relations of anacardic acids. *J Agric Food Chem.* 1993;41(6):1016–1019.
31. Li SP, Zhang GH, Zeng Q, et al. Hypoglycemic activity of polysaccharide, with antioxidation, isolated from cultured *Cordyceps mycelia*. *Phytomedicine.* 2006;13(6):428–433.
32. Morrow JD. Quantification of isoprostanes as indices of oxidant stress and the risk of atherosclerosis in humans. *Arterioscler Thromb Vasc Biol.* 2005;25(2):279–286.