

Morphometric characteristic and growth patterns of Climbing perch (*Anabas testudineus*) from Sungai Batang River, Indonesia

Abstract

This study provides scientific evidence on the length-weight relationship, condition factor and growth patterns of the Climbing perch (*Anabas testudineus*) from Sungai Batang River, Indonesia. The fish were obtained from local fishermen. A total of 156 individuals of *A. testudineus* consisted of 77 males and 79 females (55-125mm total length and 3-47g weight) were analyzed using SPSS-16 software. There were no statistically significance differences in the total length (TL), body depth (BD), and body weight (W) between male and female, as well as the W/TL and BD/TL ratio values of the fish ($P>0.05$). Most of total catch falls within the range of 80 and 89mm total length and more than 40% of total catch weighed between 10 and 19g. The W/TL ratio value of *A. testudineus* in the present study was more or less identical to that of *A. testudineus* from other different geographical areas. The fish samples were in good condition and grew negatively allometric ($b=2.7766-2.8791$). Outcomes of this study could be useful for fisheries management and conservation measures in this river.

Keywords: allometric, condition factor, length-weight, climbing perch, sungai batang river

Introduction

Like other freshwater fish species, the Climbing perch (*Anabas testudineus*) belongs to family Anabantidae is also widely distributed and commercially sold particularly in Indonesia,¹ Lao PDR,² Cambodia,³ Vietnam,⁴ Thailand,⁵ Malaysia,⁶ the Philippines,⁷ Bangladesh,⁸ and India.⁹ It is rich in iron and copper, which are essentially needed for haemoglobin synthesis¹⁰ and has high quality poly-unsaturated fats and many essential amino acids. This species is locally and seasonally common throughout its range. The presence of this species beneficially supports fish farming business, however, its presence in Australia, Papua New Guinea and India, is adversely affected native freshwater and estuarine species such as birds, reptiles, animals and predatory fish due to their sharp dorsal and opercular spines.¹¹⁻¹³ Like other labyrinth fishes i.e. Kissing gourami,¹⁴ Snakeskin gourami¹⁵ and Snakehead,¹⁶ this species can also be cultured in the earthen pond, tanks and cages.^{9,17,18} It can tolerate low dissolved oxygen, low pH and temperature changes.^{10,19,20} Pollution, overfishing and wetland conversion may potentially threaten to this species.²¹

Numerous studies on *A. testudineus* have been dedicated to describe, for example: food and feeding habits,²²⁻²⁴ reproductive biology,^{8,25} boldness,²⁶ growth performance and survival rate,^{27,28} length-weight relationship and condition factor,^{29,30} morphometric and meristic variation,³¹ water quality assessment,²⁰ phototactic responses,^{32,33} conservation and aquaculture,³⁴ and fishing activity,³⁵⁻³⁷ as well as overview of business prospect for this species.^{38,39} It is generally accepted that the length-weight relationship is the most common scientific approach that used for analyzing growth pattern of an individual fish species⁴⁰ and advanced techniques for morphometric analysis was recently presented.⁴¹ It is also useful for understanding condition factor, survival, maturity and reproduction,^{7,42} of various species from different geographical regions, as well as for comparing local and interregional, morphological and life historical among species and populations.^{14,43}

Volume 3 Issue 4 - 2019

Ahmadi

Faculty of Marine and Fisheries, Lambung Mangkurat University, Indonesia

Correspondence: Ahmadi, Faculty of Marine and Fisheries, Lambung Mangkurat University, Jalan Achmad Yani km 36, Banjarbaru 70714, South Kalimantan, Indonesia, Email ahmadizarigani@gmail.com

Received: May 17, 2019 | **Published:** July 29, 2019

Fishing activity in Sungai Batang waters is open throughout the year regardless of seasonal periods, which is done by both villagers and beyond. The use of unfriendly fishing method (e.g. electrofishing and poisoning) is still happening beyond the control particularly during nighttime. If it is allowed without forbidden, it will adverse to fish habitat and socio-economic as the whole. To manage the Climbing perch fishery resource rationally, it is therefore needed in-depth knowledge of its biology, feeding habit and ecology. As part of the research, we started to investigate the length-weight relationship and condition factor of *A. testudineus* to provide some fundamental suggestions for better fisheries management.

Materials and Methods

The research was conducted in Sungai Batang River, Martapura of South Kalimantan Province (Figure 1), located on 03°22'36" S and 114°49'29" E, determined by GPS-60 Garmin, Taiwan. The river supports the local economic activities such as fishery, agriculture and irrigation. The village consists mostly of wetland area with water level fluctuation between 0.5 and 2m. The Climbing perch locally is called *papuyu* (Figure 2). For local people, it is generally accepted that the fish are quite difficult to catch during rainy season (October-April) since fish spread out in the wetland, but very easy to collect them during dry season (May-September) because they are being concentrated on the sludge holes backwater or shallow water. The fish are being caught from the swamp using *lukah* (fish pot), *tempirai* (stage-trap) and electrofishing.

Lukah is an elongated tube-shaped made of bamboo (150cm) diameter of 20cm containing one entry funnel mounted on the inside of conical-shape and tapering inside to about 2.5cm, called *hنجap* (one-way valve, made of elastic rattan; about 40 cm one to other), and containing one exclusion funnel at the opposite side. Thus, fish can enter easily but it is difficult to escape. *Lukah* are deployed in the swamp under highly vegetated habitats with slow or no current at morning and retrieved at afternoon. *Lukah* are submerged partly

at an oblique angle of about 15° so that fish can take oxygen on the water surface. *Tempirai* is similar to Pengilar but bigger in the size. It is made of heart-shaped bamboo, 52cm high, 37cm width, and 5cm wide opening of the entrance slit. A small trap door on the top allowed for removal of catches. The snail (*Achanita* sp.) is used as bait and placed inside the trap. The trap is set in the riverbank before sunset and retrieved the next morning or mounted on a high tide and removed after low tide. The size of *Tempirai* is typically smaller than that of *Tempirai* used in Bangkau swamp. We are not able to describe the detailed electrofishing devices used in this study due to a technical barrier (unwillingness of fishermen).




Figure 1 The location of sampling site in Sungai Batang River, Indonesia.



Figure 2 A fish sample of Climbing perch from Sungai Batang River.

A total of 156 individuals of *A. testudineus* comprising 77 males and 79 females were bought from local fishermen during April 2017. All fishes were individually identified for sex, and measured for total length (TL) and body depth (BD) and weight (W). Total length was taken from the tip of the snout to the extended tip of the caudal fin. Body depth was measured from the dorsal fin origin vertically to the ventral midline of the body. The total length and body depth of each

individual were measured with a ruler to the nearest mm, while whole body weight was determined with a digital balance to an accuracy of 0.01 g (Dretec KS-233, Japan). The size distribution of fish sampled was set at 5-interval class for total length and weight, and was stated in percent. The length-weight relationship of fish can be expressed in either the allometric form.⁴⁴

$$W = aL^b \quad (1)$$

Where: W is the total weight (g), L is the total length (mm), a is the constant showing the initial growth index and b is the slope showing growth coefficient. The b exponent with a value between 2.5 and 3.5 is used to describe typical growth dimensions of relative wellbeing of fish population.⁴⁵ King⁴⁶ observed that the marked variability in the value of b may reflect changes in the condition of individual related to feeding, reproductive or migratory activities. This b value has an important biological meaning; if fish retains the same shape and grows increase isometrically (b=3). When weight increases more than length (b>3), it shows positively allometric. When the length increases more than weight (b<3), it indicates negatively allometric. Isometric growth indicates that the body increases in all dimension in the same proportion of growth whereas the negative allometry indicates that the body become more rotund as it increases in length and a slimmer body.⁴⁷ Determination coefficient (R^2) and regression coefficient (r) of morphological variables between male and female were also computed. The condition factor of fish was estimated using the following formula.⁴⁸

$$K = 100(W / L^3) \quad (2)$$

Where K is the Fulton's condition factor, L is total length (cm) and W is weight (g). The factor of 100 is used to bring K close to a value of one. The metric indicates that the higher the K value the better the condition of Climbing perch. The K value is used in assessing the health condition of fish of different sex and in different seasons. In addition, the t-test was applied to compare the body sizes and condition factor between male and female. All tests were analyzed at the 0.05 level of significance using SPSS-16 software.

Results

All estimated length-weight relationship and the ratio of body sizes of *A. testudineus* from Sungai Batang River are presented in Tables 1&2. A total of 156 individuals comprising 77 males and 79 females were analyzed. The body size of male ranged from 57 to 125mm (90.95 ± 13.87 mm) total length and from 5 to 47g (15.94 ± 7.81 g) weight. While the body size of female varied from 55 to 111mm (87.92 ± 10.52 mm) total length and from 3 to 26 g (14.04 ± 4.49 g) weight, with the ratio of male to female was 1:1.

Significant differences were observed at length-weight relationship of male and female (Figure 3A), while b values implied that the body shape displays a negative allometric growth pattern (b<3), which means that the length increases more than weight. The estimated b values obtained from allometric equations were 2.8791 for male and 2.7766 for female; with the R^2 values ranged from 0.8771 and 0.8383 indicating that more than 83% of variability of the weight is explained by the length. The index of correlation (r) of male and female were 0.9365 and 0.9156, found to be higher than 0.5, showing the length-weight relationship is positively correlated. Statistical analysis showed that there were no significant differences in the total length, body depth and body weight between female and male ($P > 0.05$). It was also clearly demonstrated in Figure 3B, no statistically significant

difference in the mean ratio of body weight to total length between male and female was observed ($P>0.05$). The ratio of W/TL for male ranged from 0.0769 to 0.3760 (0.17 ± 0.06), while for female varied from 0.0545 to 0.2342 (0.16 ± 0.04). As described in Table 2, the R^2 values were found between 0.6798 and 0.7524 indicating that more than 67% of variability of the ratio is expounded by the length. The 'r' values of male and female were 0.8674 and 0.8245, found to be higher than 0.5, showing the ratio relationship is highly correlated. The increased of body depth was directly proportional to the total length (Figure 4A). The exponent values obtained from the equation curve were 0.8876 for male and 0.8878 for female. There was no significance difference in the mean ratio of body depth to total length between male and female ($P>0.05$) (Figure 4B). The BD/TL ratio for male ranged from 0.2373 to 0.5882 (0.32 ± 0.02), while for female varied from 0.2600 to 0.4024 (0.32 ± 0.04) as given in Table 2.




Figure 3(A) Climbing perch grew negatively allometric; **(B)** No significant difference in the W/TL ratio values between male and female was observed.

The size distribution of *A. testudineus* samples in the present study is displayed in Figure 5. The highest number of catch was distributed between 80 and 84mm TL (20.78%) for male and between 85 and 89mm TL (18.99%) for female. A low number of catch was observed for larger size class more than 114mm TL. The heaviest catches of male (40.26%) and female (45.57%) weighted between 10 and 19 g. Considering the number of catch on the bar chart, the percentages of fish size distribution both length and weight was more visible in male than in female. We could find none of female catch between 30 and 49 g weight sizes. Regardless the sex, we observed that there was a difference in the exponent value for smaller length class as compared to larger ones indicating that the species has two growth levels (Figure 6). The smaller individuals (< 85 mm TL) grew isometrically with the exponent being equal to the cubic value ($W=0.00001 TL^{3.0877}$,

$R^2=0.9296$). While the larger individuals (≥ 85 mm TL) grew negatively allometric with the exponent lower than the cubic value ($W=0.00003 TL^{2.9206}$, $R^2=0.9747$), indicating that more than 92% of variability of the fish growth is explained by the length. Dealing with the condition factor (K) of the fish, there was no statistically significant difference in the mean K values between male and female ($P>0.05$) as presented in Figure 7. The mean K values obtained ranging from 0.7304 to 3.0226 (1.99 ± 0.30) for male and from 1.1000 to 3.2646 (2.01 ± 0.33) for female.




Figure 4(A) The body depth of Climbing perch increases proportionally to the total length; **(B)** No significant difference in the BD/TL ratio values between male and female was identified.

Discussion

Local people in Sungai Batang village made a point of balancing between economic, social and environmental sustainability of commercially important fish species including Climbing perch fishery resource. It is quite reasonable because fish have high economic value around IDR 45,000-60,000 per kg and have good acceptance by the consumers. Like other fish species, Climbing perch is also vulnerable to destructive fishing practices. Therefore, it is categorized by the International Union for Conservation of Nature and Natural Resources (IUCN) as a vulnerable species.⁴⁹ On the other word, Climbing perch formed a major part of the fish catch in the Sungai Batang River. Contrariwise, some researchers found none of Climbing perch during their sampling periods, for example in Betwa and Gomti Rivers, India.⁵⁰ in the Keniam River of Taman Negara Pahang,⁵¹ in River Orogodo of Niger Delta, Nigeria,⁵² in Malaysian Waters,⁵³ in Bukit Merah reservoir, Malaysia,⁵⁴ in Lubuk Lampam floodplain of South Sumatera, Indonesia,⁵⁵ in Shahrbijar River, Iran,⁴⁰ as well as

in Narreri Lagoon of Sindh, Pakistan.⁴³ This is likely attributable to sample size variation, typical fishing gear used, and environmental factors.




Figure 5 The length size (A) and weight size (B) distribution between male and female of Climbing perch taken from Sungai Batang River.




Figure 6 The circle mark on the curve indicates the intersection points of smaller (<85 mm TL) and larger individuals (≥ 85 mm TL) of Climbing perch growth pattern.

The maximum size (125mm TL) of *A. testudineus* in the present study was larger than *A. testudineus* collected from Chandpur, Matlab of Kalipur, Bangladesh: 93mm,⁴⁴ but it was lower than maximum size of *A. testudineus* from Kausalyaganga of Orissa, India: 175mm,⁹ from Deepar Beel of Assam, India: 134mm,²⁹ from Mohanpur of West Bengal, India: 170mm,⁵⁶ or from a fish pond Banjarbaru of South Kalimantan, Indonesia: 170mm.⁵⁷ During fishing season, it is very likely to collect *A. testudineus* smaller than 55mm TL and 3g weight

using the nets, but fishermen prefer release them back to the river rather than sold them with no or lower price, conversely the smaller fish might be untrappable because of fishing gear selectivity.⁵⁸ The other way, it is also possible for fishermen to collect fish with the size larger than 125mm TL in the study; however, it is beyond our investigation due to the transactional selling of fish is usually occurred in early morning before the fish transported to the local market.




Figure 7 The relationship of the W/TL ratio and condition factor between male and female of Climbing perch. No statistically significant difference in this relationship was observed.

In the present study, *A. testudineus* grew negatively allometric ($b < 3$), which is similarly documented in the previous studies.^{9,21,30,32,44} Our finding is contrary to *A. testudineus* collected from Deepar Beel of Assam, India that showed a positive allometric growth pattern.²⁹ Dealing with variation of these growth types, some of other studies may provide much lower b values than these findings.⁵⁹ The weight-length relationships are not constant over the entire year and vary according to the following factors: food availability, feeding rate, gonad development and spawning period,^{60,61} fecundity,⁵⁶ temperature,⁶² salinity⁵ and inherited body shape.⁶³

From the exponent values obtained, we can say that *A. testudineus* male ($b = 2.8791$) grow faster than female ($b = 2.7766$). Such growth pattern was similarly found in our previous study.⁵⁷ The exponent value ($b = 2.9206$) for the larger individuals (85-125mm TL) was lower as compared to other fish species like *M. cavasius* ($b = 3.150$) from Indus River, Pakistan observed at the maximum size of 148mm.⁶⁴ It meant that such ontogenetic variations in the cubic law differ from other fish species; however, a more detailed analysis regarding the reasons contributing to such variations is necessary. There was a variability of the ratios of body weight to total length among *A. testudineus* species from different geographical areas (Table 3). The average W/TL ratio of *A. testudineus* in the present study (0.167) was relatively higher than that of *A. testudineus* from Deepar Beel of Assam, India: 0.162,²⁹ from Chandpur, Matlab of Kalipur, India: 0.069⁴⁴ or from Tetulia River, Bangladesh: 0.083.²¹ However, this ratio was considerably lower as compared to those sampled from Kausalyaganga of Orissa, India: 0.390,⁹ from Kuttanad of Kerala, India: 0.363,³⁰ or from Mohanpur of West Bengal, India: 0.353mm.⁵⁶

The average K value obtained (2.00) for *A. testudineus* in the present study (Table 3) was slightly lower than that of *A. testudineus* from Kuttanad of Kerala, India: 2.06³⁰ or from Kausalyaganga of

Orissa, India: 2.07.⁹ Barnham & Baxter⁶⁵ suggested that if the K value is 1.00, the condition of the fish is poor, long and thin. A 1.20 value of K indicates that the fish is of moderate condition and acceptable to many anglers. A good and well-proportioned fish would have a K value that is approximately 1.40. Based on this criterion, the Climbing perch from Sungai Batang River were in good condition although fish grew negatively allometric. Regardless the sex, Bagenal & Tesch⁶⁶ documented the K value between 2.9 and 4.8 for mature freshwater

fish species; meanwhile the K values in the current study ranged from 0.7304 to 3.2646, indicating that some of individual fish are being mature when captured. Variation in the value of the mean K may be attributed to biological interaction involving intraspecific competition for food and space between species including sex, stages of maturity, state of stomach contents and availability of food,^{50,66} as well as season and other environmental conditions.⁶⁷

Table 1 Total length, weight and condition factor of Climbing perch sampled from Sungai Batang River

Sex	N	Total length(mm)			Weight(g)			a	b	R ²	r	Growth pattern	K Mean±SD
		Min	Max	Mean±SD	Min	Max	Mean±SD						
Males	77	57	125	90.95±13.87	5	47	15.94±7.81	0.00003	2.8791	0.8771	0.9365	A-	1.99±0.30
Females	79	55	111	87.92±10.52	3	26	14.04±4.49	0.00005	2.7766	0.8383	0.9156	A-	2.01±0.33
Pooled	156	55	125	89.42±12.34	3	47	14.97±6.40	0.00004	2.8353	0.8625	0.9287	A-	2.00±0.31

a, constant; b, exponent; R², determination coefficient; r, regression coefficient; A-, negative allometric; K, condition factor.

Table 2 The ratio of body sizes of Climbing perch from Sungai Batang River

Sex	n	W/TL	a	b	R ²	r	BD/TL	a	b	R ²	r
Male	77	0.12±0.02	0.00003	1.8791	0.7524	0.8674	0.32±0.02	0.526	-0.112	0.0652	0.2553
Female	79	0.16±0.04	0.00005	1.7766	0.6798	0.8245	0.32±0.04	0.5221	-0.112	0.0306	0.1749
Pooled	156	0.16±0.05	0.00004	1.8353	0.7244	0.8511	0.32±0.02	0.518	-0.11	0.0432	0.2078

a, constant; b, exponent; R², determination coefficient; r, regression coefficient; BD, body depth; W, body weight; TL, total length

Table 3 Comparative parameters of length-weight relationships and growth patterns of Climbing perch from different geographical areas

Locations	Country	n	W/TL	a	b	R ²	r	Growth pattern	Average K	References
Sungai Batang River	Indonesia	156	0.167	0.00004	2.8353	0.8625	0.9287	A-	2.000	Present study
Banjarbaru, South Kalimantan	Indonesia	608	0.216	0.0005	1.8049	0.4339	0.6587	A-	2.115	Ahmadi (2018b) ⁵⁷
Kuttanad, Kerala	India	246	0.363	0.0003	2.8452	0.9556	0.9775	A-	2.060	Kumary and Raj (2016) ³⁰
Mohanpur, Nadia District	India	30	0.353	-2.025	2.316	0.5396	0.7346	A-	-	Ziauddin et al. (2016) ⁵⁶
Kausalyaganga, Orissa	India	544	0.390	-1.432	2.7201	0.0821	0.9264	A-	2.070	Kumar et al. (2013) ⁹
Deepar Beel, Assam	India	120	0.162	-2.540	3.645	0.8500	0.9220	A+	1.000	Rahman et al. (2015) ²⁹
Chandpur, Matlab, Kalipur	Bangladesh	73	0.069	-1.518	2.423	0.9660	0.9828	A-	1.085	Begum and Minar (2012) ⁴⁴
Tetulia River	Bangladesh	176	0.083	0.0220	2.90	0.9740	0.9869	A-	-	Hossain et al. (2015) ²¹

a, constant; b, exponent; R², determination coefficient; r, regression coefficient; A, allometric; K, condition factor.

In the investigated area, local fishermen mostly used *lukah* (fish pot), *tempirai* (stage-trap) and also electrofishing for catching *A. testudineus* from its natural habitats, while other workers used the seine nets in the cultured ponds, Bangladesh,¹¹ fish nets in local ditches in Adra Purulia, West Bengal India,²³ gillnets in flood plain swamp, South Sumatera Province or in Batang Kerang Floodplain, Balai Ringin, Sarawak,^{68,69} *Khepla Jal* (cast net) in the Chalan Beel, Bangladesh,⁷⁰ *tampirai* (wire stage-trap) in Sebangau River, Central

Kalimantan Province,⁷¹ a conical trap in the ponds, West Bengal India,²⁴ *lalangit* (horizontal set gillnet) in Bangau swamp, South Kalimantan Province,³⁶ *pukot* and electrofishing in floodplain Lakes of Agusan Marsh, Philippines,⁷² light traps in the cultured ponds, South Kalimantan Province.^{32,33} According to Aminah & Ahmadi,³⁷ the catchability of the gear is higher during the dry season compared to the wet season, because the fish are being concentrated on the sludge holes or shallow areas and allow for catching them easily.

While Kamaruddin et al.⁷³ found the opposite condition, with the reason is the wet season indicates the main feeding and growing time for the fish,⁵² and it is very likely to catch the fish. In addition, female climbing perch showed a marked increase in their food intake early in the wet season.⁷ Variation in the catches may be attributed to the catchability of the gears, species target, fishing ground characteristic, fishing operation, and the abundance of the fish in that area.^{70,74} There are three main constraints being faced in this area of study: the first, there is no daily record for *A. testudineus* catch in quantity (e.g. number, length and weight) because fishes are directly sold to traders or consumers in some places; secondly, fishing activity is on-going throughout the year regardless of seasonal periods, thus resulted in the ratio of the fish exploitation rate to the fish growth rate in this river is still unpredictable; and the thirdly, the use of electrofishing is still beyond the control because it is usually undertaken at the night. So, it is a great challenge for Marine and Fisheries Services of Banjar District to improve the quality of inland fishery statistical data and manage the fishery resources through EAFM program.

Conclusion

The climbing perch sampled from Sungai Batang River were in good condition and grew negatively allometric. The scientific information on the length-weight relationship and condition factor of climbing perch could be used to estimate the stock population and to take conservation measures.

Acknowledgments

None.

Conflicts of interest

The author declares that there are no conflicts of interest.

References

1. Akbar J, Mangalik A, Fran S. Application of fermented aquatic weeds in formulated diet of climbing perch (*Anabas*). *Int J Engineer Res Sci*. 2016;2(5):240–243.
2. Sokheng C, Chhea CK, Viravong S, et al. Fish migrations and spawning habits in the Mekong mainstream: a survey using local knowledge (basin-wide). *Assessment of Mekong fisheries: Fish Migrations and Spawning and the Impact of Water Management Project (AMFC)*. AMFP Report 2/99. Vientiane, Lao, P.D.R. 1999.
3. Sverdrup JS. Fisheries in the lower Mekong basin: status and perspectives. MRC Technical Paper No. 6, Mekong River Commission. Phnom Penh. 2002;8–23.
4. Van KV, Hoan VQ. Intensive nursing climbing perch (*Anabas testudineus*) in hapas using pellet feed at different protein levels. *J Sci Dev*. 2009;7:239–242.
5. Chotipuntu P, Avakul P. Aquaculture potential of climbing perch, *Anabas Testudineus*, in brackish water. *Walailak J Sci Tech*. 2010;7(1):15–21.
6. Zalina I, Saad CR, Christianus A, et al. Induced breeding and embryonic development of climbing perch (*Anabas testudineus*, Bloch). *J Fish Aquat Sci*. 2012;1–16.
7. RAD Bernal, Aya FA, de Jesus-Ayson EGT, et al. Seasonal gonad cycle of the climbing perch *Anabas testudineus* (Teleostei: Anabantidae) in a tropical wetland. *Ichthyol Res*. 2015;62(4):389–395.
8. Uddin S, Hasan MH, Iqbal MM, et al. Study on the reproductive biology of Vietnamese climbing perch (*Anabas testudineus*, Bloch). *Punjab Univ J Zool*. 2017;32(1):1–7.
9. Kumar K, Lalrinsanga PL, Sahoo M, et al. Length-weight relationship and condition factor of *Anabas testudineus* and *Channa* species under different culture systems. *World J Fish Mar Sci*. 2013;5(1):74–78.
10. Sarma K, AK Pal, Ayyappan S, et al. Acclimation of *Anabas testudineus* (Bloch) to three test temperatures influences thermal tolerance and oxygen consumption. *Fish Physiol Biochem*. 2010;36(1):85–90.
11. Storey AW, Roderick ID, REW Smith, et al. The spread of the introduced climbing perch (*Anabas testudineus*) in the Fly River system, Papua New Guinea, with comments on possible ecological effects. *Int J Ecol Envi Sci*. 2002;28:103–114.
12. Hitchcock G. Climbing Perch (*Anabas testudineus*) (Perciformes: Anabantidae) on Saibai Island, northwest Torres Strait: first Australian record of this exotic pest fish. *Memoirs of the Queensland Museum*. 2008;52(2):207–211.
13. Paliwal GT, Bhandarkar SV. Diversity of exotic fishes in Navegaonbandh Reservoir with reference to negative impact of *Anabas* (Anabantidae) on Biodiversity. *Int J Curr Microbiol App Sci*. 2014;3(8):592–597.
14. Zohrah HS, Haji Kasim HD. Pond aquaculture of Kissing gourami *Helostoma temminckii* (Pisces: Helostomatidae) in Bukit Udal, Tutong. A preliminary investigation. *Bruneiana: Anthol Sci Artcl*. 2002;3:34–41.
15. Tate M, Mcgoran R, White CR, et al. Life in a bubble: the role of the labyrinth organ in determining territory, mating and aggressive behaviours in anabantoids. *J Fish Biol*. 2017;91(3):723–749.
16. Xie H, Lü X, Zhou J, et al. Effects of acute temperature change and temperature acclimation on the respiratory metabolism of the snakehead. *Turkish J Fish Aquat Sci*. 2017;17:535–542.
17. Long DN, Hieu NT, Tuan NA. Trials on intensive culture system of Climbing perch (*Anabas testudineus*) in earthen ponds in Long An province. *Sci J Can Tho Univ*. 2006;93–103.
18. Mondal MN, Shahin J, Wahab MA, et al. Comparison between cage and pond production of Thai climbing perch (*Anabas testudineus*) and Tilapia (*Oreochromis niloticus*) under three management systems. *J Bangladesh Agri Univ*. 2010;8(2):313–322.
19. Kohinoor AHM, Jahan DA, Khan MM, et al. Culture potentials of Climbing perch, *Anabas testudineus* (Bloch) under different stocking densities at semi-intensive management. *Bangladesh Fish Res*. 2009;13(2):115–120.
20. NV Be, NT Giao, TH Đan, et al. Water quality in intensive Climbing perch ponds (*Anabas testudineus*) and suggestion for better management of wastewater discharge. *Imperial J Interdiscipl Resh*. 2017;3(9):665–671.
21. Hossain MY, Hossen MA, Pramanik MNU, et al. Threatened fish of world: *Anabas testudineus* (Bloch, 1792) (Perciformes: Anabantidae). *Croatian J Fish*. 2015;73:128–131.
22. Roy D, Masud AA, Bhouiyana NA, et al. Food and feeding habits of Climbing perch *Anabas testudineus* (Bloch) and indigenous cat fish Rita rita (Hamilton). *Int J Bio Res*. 15(1):1–6.
23. Bhattacharjee I, Chandra G. Food and feeding habits of three air-breathing fish in its natural habitat. *Int J Fish Aquat Stud*. 2016;4(3):586–589.
24. Patra BC, Patra S, Bhattacharya M. Evaluating the nutritional condition of an Indian Climbing perch, *Anabas testudineus* fingerlings by the RNA/DNA, Ca/P ratio and Protein Bio-Synthesis in Liver and Muscle. *Fish Aqua J*. 2017;8(1):1–8.
25. Hafijunnahar, Rahman MA, Hossain MMM. An investigation on breeding biology of Vietnam strain of climbing perch, *Anabas testudineus* (Bloch) reared in a commercial hatchery. *Int J Fish Aquat Stud*. 2016;4(1):08–12.
26. Binoy VV. Comparative analysis of boldness in four species of freshwater teleosts. *Indian J Fish*. 2015;62(1):128–130.

27. Rahman MA, Marimuthu K. Effect of different stocking density on growth, survival and production of endangered native fish climbing perch (*Anabas testudineus*, Bloch) fingerlings in nursery ponds. *Adv Envi Biol.* 2010;4(2):178–186.

28. Putra DF, Fanni M, Muchlisin ZA, et al. Growth performance and survival rate of climbing perch (*Anabas testudineus*) fed *Daphnia* sp. enriched with manure, coconut dregs flour and soybean meal. *AACL Bioflux.* 2016;9(5):944–948.

29. Rahman A, Talukdar K, Rahman W, et al. Length-weight relationship and relative condition factor of *Anabas testudineus* (Bloch) of Deepar Beel (wetland) of Assam, India. *Int J Appl Res.* 2015;1(11):956–958.

30. Kumary KSA, Raj S. Length-weight relationship and condition of climbing perch *Anabas testudineus* Bloch population in Kuttanad, Kerala. *Int J Adv Res Biol Sci.* 2016;3(9):21–26.

31. Hossen MB, Sharker MR, Rahman MA, et al. Morphometric and meristic variation of indigenous and Thai Koi, *Anabas testudineus* available in coastal region of Bangladesh. *Int J Innov Res.* 2017;2(1):01–08.

32. Ahmadi, Muhammad, E Lilimantik. Phototactic response of climbing perch *Anabas testudineus* to different colors and light pattern of LED light traps. *AACL Bioflux.* 2018;11(3):678–689.

33. Ahmadi. Phototactic response and morphometric characteristic of climbing perch *Anabas testudineus* (Bloch, 1792) under culture system. *Croatian J Fish.* 2018a;76(4):164–173.

34. Sarkar UK, Depak PK, Kapoor D, et al. Captive breeding of climbing perch *Anabas testudineus* (Bloch, 1792) with Wova–FH for conservation and aquaculture. *Aquac Res.* 2005;36(10):941–945.

35. Iwata A, Ohnishi N, Kiguchi Y. Habitat use of fishes and fishing activity in plain area of southern Laos. *Asian Afr Area Stud.* 2003;3:51–86.

36. Irhamsyah, Ahmadi, Ruspiliyansari. Fish and fishing gears of the Bangkau Swamp, Indonesia. *J Fish.* 2017;5(2):489–496.

37. Aminah S, Ahmadi. Experimental fishing with LED light traps for Three-spot gourami (*Trichogaster trichopterus*) in Martapura, Indonesia. *Int J Fish Aquat Stud.* 2018;6(1):37–42.

38. Khatune-Jannat M, Rahman MM, Bashar MM, et al. Effects of stocking density on survival, growth and production of Thai Climbing Perch (*Anabas testudineus*) under fed ponds. *Sains Malaysiana.* 2012;41(10):1205–1210.

39. Izmaniar H, Mahyudin I, Agusliani E, et al. The business prospect of Climbing perch fish farming with biofloc technology at De' Papuya Farm Banjarbaru. *Int J Envi Agri Biotechnol.* 2018;3(3):1145–1153.

40. Asadi H, Sattari M, Motalebi Y, et al. Length-weight relationship and condition factor of seven fish species from Shahrivar River, Southern Caspian Sea basin, Iran. *Iranian J Fish Sci.* 2017;16(2):733–741.

41. Mojekwu TO, Anumudu CI. Advanced techniques for morphometric analysis in fish. *J Aquac Res Dev.* 2015;6:354.

42. Buragohain A. Study of absolute fecundity of *Mystus carcio* (Hamilton, 1822) of Puthimari River (Hajo) of Kamrup (R) District of Assam, India. *Int J Zool Stud.* 2018;3(2):268–270.

43. Awan KP, Qamar N, Farooq N, et al. Sex ratio, length weight relationships and condition of eight fish species collected from Narreri Lagoon, Badin, Sindh, Pakistan. *J Aquac Mar Biol.* 2017;5(4):00130.

44. Froese R. Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *J Appl Ichthyol.* 2006;22(4):241–253.

45. Bagenal T. *Methods for assessment of fish production in freshwaters*. 3rd edn. Oxford: Blackwell Scientific Publication. Oxford, London. 1978;365.

46. King EP. Length weight relationship of Nigerian coastal water fishes. *Fishbyte.* 1996;19(4):53–58.

47. Jobling M. *Environmental factors and rates of development and growth*. In: Hart PJ & Reynolds JD, editors. *Hand book of fish biology and fisheries*: Vol.1. Fish Biology. Oxford: Blackwell Publishing Ltd. 2008;97–122.

48. Weatherley AH, Gill HS. *The biology of fish growth*. Academic Press: London. 1987;14–21.

49. Singh YB, Saha H, Mandal B, et al. Breeding of Climbing perch (*Anabas testudineus* Bloch, 1792) induced with Ovatide. *Israeli J Aquac Bamidgeh.* 2012;64:766–769.

50. Sani R, Gupta BK, Sarkar UK, et al. Length-weight relationships of 14 Indian freshwater fish species from the Betwa (Yamuna River tributary) and Gomti (Ganga River tributary) rivers. *J Appl Ichthyol.* 2010;26(3):456–459.

51. Azham Yahya M, Singh HR. An assessment of the distribution of the freshwater fishes of the Taman Negara Pahang, Malaysia. IEEE Symposium on Business. *Engineering and Industrial Applications.* 2012;175–180.

52. Meye JA, Ikomi RB. Seasonal fish abundance and fishing gear efficiency in River Orogodo, Niger Delta, Nigeria. *World J Fish Mar Sci.* 2012;4(2):191–200.

53. Khairul Adha AR, Esa Y, Arshad A. The influence of alien fish species on native fish community structure in Malaysian waters. *Kuroshio Sci.* 2013;7(1):81–93.

54. Mohd Shafiq Z, Amir Shah Ruddin MS, Zarul HH, et al. The effect of seasonal changes on freshwater fish assemblages and environmental factors in Bukit Merah reservoir (Malaysia). *Transylvanian Rev System Ecol Res.* 2014;16(1):97–108.

55. Jubaedah D, Kamal MM, Muchsin I, et al. The trophic status of the Lubuk Lampam floodplain in South Sumatera, Indonesia. *Makara J Sci.* 2016;20(2):61–70.

56. Ziauddin G, Behera S, Kumar S, et al. Morphometrical and gonadal studies of a threatened fish, *Anabas testudineus* with respect to seasonal cycle. *Int J Fish Aquac Sci.* 2016;6(1):7–14.

57. Ahmadi. Phototactic responses of climbing perch (*Anabas testudineus*) to various intensities and colors of LED underwater lamps: Recommendation for future research. *Adv Biol Earth Sci.* 2018b;3(2):114–129.

58. Ahmadi, Rizani A. Light traps fishing in Sungai Sipai flood swamp of Indonesia: Recommendations for future study. *Kasetsart Univ Fish Res Bull.* 2013;37(2):17–30.

59. Froese R, Pauly D. Fish Base. *World Wide Web electronic publication.* 2019.

60. Amin SMN, Arshad A, Siraj SS, et al. Reproductive seasonality and maturation of the sergestid shrimp, *Acetes japonicus* (Decapoda: Sergestidae) in coastal waters of Malacca, Peninsular Malaysia. *Afr J Biotechnol.* 2010;9(45):7747–7752.

61. Wong BY, Ong HKA, Khoo G. Length-weight relationships of *Acetes* spp. sampled along the West Coast of Peninsular Malaysia. *Sains Malaysiana.* 2015;44(3):379–386.

62. Lopez-Martinez J, Arreguin-Sánchez F, Hernandez Vazquez S, et al. Inter-annual variation of growth of the brown shrimp *Farfantepenaeus californiensis* and its relation to temperature. *Fish Res.* 2003;61(1–3):95–105.

63. Yousuf F, Khurshid S. Length-weight relationship and relative conditions factor for the halfbeak *Hamirampus far* Forsskal, 1775 from the Karachi coast. *Univ J Zool Rajashahi Univ.* 2008;27:103–104.

64. Muhammad H, Iqbal Z, Bashir Q, et al. Length weight relationship and condition factor of catfish species from Indus River, Pakistan. *Punjab University Journal of Zoology*. 2017;32(1):35–38.
65. Barnham C, Baxter A. Condition Factor, K, for Salmonid Fish. *Fish Notes*. 1998;1–3.
66. Bagenal TB, Tesch FW. In *Bagenal and Tesch: Methods for the assessment of fish production in fresh waters*. 3rd edn. Oxford: Blackwell Scientific Publications. 1978;93–123.
67. Braga FMS. Comprimento paraalgun speixes marinhos. *Rev Brasil Biol*. 1986;46(2):339–346.
68. Muthmainnah D, Gaffar AK. Fish and fisheries in flood plain swamp in middle part of Musi River. *Indonesian J Envi Man Sustain*. 2017;1(1):1–5.
69. Khairul Adha AR, Daud SK, Siraj SS, et al. Freshwater fish diversity and composition in Batang Kerang Floodplain, Balai Ringin, Sarawak. *Pertanika J Tropic Agri Sci*. 2009;32(1):7–16.
70. Sultana N, Islam MN. A study of Khepla Jal (Cast Net) in the Chalan Beel, Bangladesh. *Int J Modern Engineer Res*. 2017;7(11):34–40.
71. Thornton SA, Dudin, SE Page, et al. Peatland fish of Sebangau, Borneo: diversity, monitoring and conservation. *Mires and Peat*. 2018;22(4):1–25.
72. Jumawan JC, Seronay RA. Length–weight relationships of fishes in eight floodplain Lakes of Agusan Marsh, Philippines. *Philippine J Sci*. 2017;146(1):95–99.
73. Kamaruddin IS, Mustafa Kamal AS, Christianus A, et al. Fish community in Pengkalan Gawi – Pulau Dula Section of Kenyir Lake, Terengganu, Malaysia. *J Sustain Sci Man*. 2011;6(1):89–97.
74. Ahmadi. The catching efficiency of light traps and morphometric characteristics of the native species from Barito River, Indonesia. *Ege J Fish Aquat Sci*. 2018c;35(1):63–72.