

Mir-34a-5p and mir-34a-3p contribute to the signaling pathway of p53 by targeting overlapping sets of genes

Abstract

In contrary to the common belief that only one strand of the pre-miRNA is active (usually the 5p one) while the second one is discarded, functional 5p and 3p have been observed for many miRNAs. Among those miRNAs is miR-34a which is a target gene of the tumor suppressor p53. In this paper we have re-examined the role of miR-34a-5p and miR-34a-3p in the signaling pathway of p53. We found that they target overlapping sets of genes (MDM2 and THBS1). By a GO enrichment analysis we found that THBS1 is involved in cancer and metastasis relevant processes. We have also deduced that p53, MDM2 and miR-34a form a I1-FFL that can speed the response of p53 to external stress signals.

Keywords: cancer, tumor suppressor, micro RNAS

Volume 3 Issue 1 - 2018

Abderrahim Chafik

Ministry of Employment and Social Affairs, Morocco

Correspondence: Abderrahim Chafik, The Abdus Salam International Center for Theoretical Physics (ICTP), Strada Costiera, 111 - 34151 Trieste, Italy, Email ab.chafik@gmail.com

Received: November 01, 2017 | **Published:** January 18, 2018

Introduction

P53 has been dubbed “guardian of the genome”¹ or gatekeeper² due to its central role in maintaining the genomic stability and tumor suppression.³⁻⁵ It has been found to be mutated in about half of the human cancers.^{6,7} Since its discovery p53 has been subject to a tremendous amount of work making it one of the most extensively studied gene. Its tumor suppressive role consists in inducing anti-proliferative cellular responses to a variety of stress signals, namely a cell-cycle arrest, senescence or an apoptosis. P53 can then be activated in response to DNA damage, hypoxia, or aberrant growth signals resulting from deregulated expression of oncogenes.⁸⁻¹⁰ Under normal conditions p53 is expressed at low level as a result of the action of the ubiquitin-protein ligase MDM2, that repress p53 through ubiquitin-mediated degradation.^{11,12} In response to a stress signal as DNA damage p53 undergoes post-transcriptional modifications (phospho-rylation and acetylation) by various kinases, including ATM, ATR, DNA-PK, Chk1, and Chk2.¹³ These modifications activate p53 by inhibiting MDM2 from binding to its N-terminal and allow it to carry out its major function as a transcription factor that binds to specific DNA sequences and activate the transcription of adjacent genes. By inducing the expression of its target genes p53 induces cell-cycle arrest or apoptosis. MicroRNAs (miRNAs) are small (20-25 nucleotides) non-coding RNAs that negatively regulate gene expression post transcriptionally by base-pairing to the 3'UTR of target mRNAs leading to repression of protein production or mRNA degradation (For a recent introduction see for example.¹⁴ A single miRNA can target hundreds of different mRNAs, and a single mRNA can be coordinately suppressed by multiple different miRNAs, they are involved in many biological processes, in particular cancer-relevant processes such as proliferation, cell cycle control, apoptosis, differentiation, migration and metabolism. A miRNA can have either oncogenic or tumor suppressive function. Examples of tumor suppressor miRNAs are let-7 family, miR-29, miR-34 and miR-15, and of oncogene miRNAs are miR-17-92, miR-155 and miR-221. The biogenesis of miRNAs can be summarized as follows: a primary transcript (PRI-miRNA) is first generated by RNA polymerase as separate transcriptional units or embedded within the introns

of protein coding genes. Then the primary transcript is processed by the microprocessor complex containing the RNase III enzyme Drosha to an approximate 70-nucleotide (nt) pre-miRNA hairpin (the precursor-miRNA).¹⁵⁻¹⁷ Pre-miRNAs are subsequently exported to the cytoplasm by exportin 5(XPO5)^{18,19} where their terminal loops are excised by the RNase III Dicer to give rise to a double-stranded 22nt stem composed of 5'âĀĤ and 3'âĀĤ strands representing 5p and 3p respectively. While one of the two strands (the passenger strand miRNA*) is discarded, the other one (the guide) is then embedded into the RISC (RNA Induced Silencing Complex) to complementary target mRNA for post-transcriptional gene silencing.²⁰

Since it was believed that according to the thermodynamic stability of the pre-miRNA cells preferentially select the less stable one of the two strands (the guide) and destroy the other one (the miRNA*), early works on miRNAs have focused on the guide strand (which was usually considered as the 5p one because it was found to be more abundant than its counterpart miRNA* in humans.²¹ However, even though miRNA* are less abundant they are often present and remain functional because they conserve their seed sequences and have been isolated from RISC.^{22,23} It has been shown by profiling analyses that both strands could be co-accumulated in some tissues while being subjected to strand selection in others.²⁴⁻²⁶ The interplay between the 5p and 3p strands from the same precursor has been shown either in arm-switching where the dominant miRNA is switched from one arm of the precursor to the other²⁷ or by targeting overlapping sets of genes when the abundances of the two strands are similar.²⁸ Functional 5p and 3p have been characterized for many miRNAs. Examples include miR-9,²⁹ miR-17,³⁰ miR-19,²³ miR-28,³¹ miR-30c,³² miR-125a,³³ miR-142,³⁴ miR-155,³⁵ miR-199,^{36,37} miR-223,³⁸ miR-342,³⁹ miR-2015,⁴⁰ miR-18a,⁴¹ miR-582.⁴² In some cases the two arms function in opposing ways (e.g. miR-28,³¹ and miR-125³³), while in others they function in joint fashion (e.g. miR-199³⁷ and miR-155³⁵). miRNAs have been shown to be important components in the p53 network Their interactions with p53 have been demonstrated through the identification of several miRNAs as direct target genes of p53. By inducing the expression of specific miRNAs that have a tumor suppressive function a novel mechanism for tumor suppression for

p53 has been then revealed. In particular the role of the miR-34 family has been reported in several studies.⁴³ In mammals miR-34 family consists of miR-34a, miR-34b and miR-34c, that are encoded by two different genes. miR-34a is encoded by an individual transcript in chromosome 1 and expressed in a majority of tissues, while miR-34b and miR-34c share a common primary transcript in chromosome 11 and are mainly expressed in lung tissues. Several studies have reported that the members of miR-34 family were direct target genes of p53 and their up regulation induces apoptosis and cell-cycle arrest.⁴⁴⁻⁵⁰ Indeed, ectopic expression of miR-34 induces cell cycle arrest in both primary and tumor-derived cell lines.⁵⁰ Inactivation of miR-34a strongly attenuates p53-mediated apoptosis in cells exposed to genotoxic stress mir-34b/mir-34c were also down-regulated in p53-null human ovarian carcinoma cells and both cooperate in suppressing proliferation of neo plastic epithelial ovarian cells.⁴⁵ Since cell-cycle arrest and apoptosis are the responses of p53 to the stress signals, these facts imply that miR-34 mediate the tumor suppressive function of p53. On the other hand the members of the miR-34 family can have decreased expression in cancer because of the inactivating mutations of p53 or the expression of viral inhibitors of p53, but also as a consequence of their own mutational or epigenetic inactivation.

Since 30% of all genes and the majority of the genetic pathways are regulated by miRNAs,⁵¹⁻⁵³ we can expect some miRNAs to regulate p53 and its pathway. This hypothesis has been verified and some miRNAs have been identified as regulators of p53. miR-504 can negatively regulate p53 expression through its binding to two binding sites in human p53 3' UTR.⁵⁴ It has also been reported that miR-125b is another miRNA targeting p53.⁵⁵ Another regulator is miR-29 that was identified as a positive regulator of p53 that up-regulate p53 protein levels and induce p53-mediated apoptosis through repression of p85^Îs.⁵⁶ Furthermore, miR-34a, which is a transcription target of the p53 protein, was also found to positively regulate p53 activity and function in apoptosis through its direct negative regulation of SIRT1.⁵⁷ SIRT1 is a negative regulator of p53, which physically interacts with p53 and de acetylates Lys382 of p53.⁵⁸ The purpose of this paper is the study of the role of both miR-34a-5p and miR-34a-3p in the signaling pathway of the p53.

Materials and methods

We begin by using DIANA-miRPath v3. 0 which is a miRNA pathway analysis web-server, providing accurate statistics utilizing predicted or experimentally validated miRNA interactions derived from DIANA-TarBase.⁵⁹ We first perform a KEGG reverse search for miRNAs in the p53 signaling pathway (hsa04115) by choosing the tar Base v 7.0 for method. In the same web-server we obtain the target genes of the different miRNAs that we found. The distribution of the target genes in the pathway can also be displayed. Then we use the cytoscape 3.4.0 software to visualize the interactions where the relevant genes are involved.⁶⁰ We take the int Act¹ to be our source of interactions. We filter by taxonomy identifier to restrict the obtained network to the human case. In order to form clusters of genes that share a similar function we use the cluster Maker, a Cytoscape application with the GLayer Community Clustering algorithm. The BiNGO, a Cytoscape Plugin⁶¹ gives the Gene Ontology (GO) terms that are significantly overrepresented in each cluster.

Results and discussion

The result of the KEGG reverse search was a list containing 660 different miRNAs including hsa-miR-34a-5p (p-value=1.840530e-100)

and hsa-miR-34a-3p (p-value=4.325757e-5). The respective p-values can be interpreted as expressing the fact that 5p is more abundant than 3p. The list of target genes contains 30 genes for miR-34a-5p and two for miR-34a-3p (MDM2 and THBS1) that are also target genes of the miR-34a-5p. The involvement of these genes in the p53 signaling pathway is shown in (Figure 1). THBS1 By loading the list of the 30 target genes of miR-34a-5p (containing also the two targets of miR-34a-3p) in Cytoscape 3.4.0 and searching the interactions in Int Act we obtain a network composed of 2492 node and 6493 edges. After clustering this network by the cluster Maker application we obtain the result shown in (figure 2) where the cluster containing THBS1 was highlighted, this cluster is shown in (figure 3). In order to find the pathways in which the THBS1 gene is involved we use the BINGO application of Cytoscape, and the relevant GO terms that are over expressed in the corresponding cluster are given in (Table 1) Int Act is one of the largest available repositories for curated molecular interactions data, storing PPIs as well as interactions involving other molecules. It is hosted by the European Bioinformatics Institute. Int Act has evolved into a multisource curation platform and many other databases curate into IntAct and makes their data available through it.²³ According to this analysis we can deduce that the THBS1 gene is involved in regulating tumor suppression processes. But it is also clear that it contributes to some anti metastatic pathways like angiogenesis and cell adhesion.




Figure 1 The signaling pathway of p53 with the target genes of miR-34-5p/3p highlighted.




Figure 2 The result of the clustering of the initial network representing different interactions of the target genes of miR-34a-5p/3p.



Figure 3 The cluster representing THBS1 and its interactions.

Table 1 GO term enrichment for the THBS1

GO terms over expressed	p-value
Regulation of response to external stimulus	2:34 10 ⁶
Regulation of response to stress	3:09 10 ⁶
Regulation of cell adhesion	6:29 10 ⁴
Positive regulation of apoptosis	5:48 10 ³
Negative regulation of cell-matrix adhesion	3:07 10 ³
Induction of apoptosis	1:42 10 ³
Negative regulation of angiogenesis	6:48 10 ³

MDM2

MDM2 and p53 form a negative feedback loop, in which p53 induces the expression of MDM2 which mediates the degradation of p53.^{62,63} Indeed, in unstressed cells p53 is present at very low levels due to the continuous degradation by ubiquitination which is mediated by MDM2. In stressed cells, in order for p53 to accumulate in response to stress signals it has to escape the degradation by MDM2. In human cancers if the p53 is not mutated its wild-type function is inhibited by the high level of MDM2, leading to the down regulation of its tumor suppressive pathways. Thus, the inhibition of the MDM2-p53 interaction represents a promising therapeutic strategy for the treatment of cancer. According to the previous result miR-34a can contribute to the regulation of this interaction and guarantee a faster p53 response to external stimuli. Indeed, since MDM2 is a target gene of miR-34a which means that it is negatively regulated by this miRNA, we can re-present the interaction between the three molecules (p53, MDM2 and miR-34a) as a Feed Forward Loop (FFL). According to the signs of the three interactions (activation/repression) the present loop is a type 1 incoherent FFL (I1-FFL) (figure 4). The dynamics of this FFL can be described as follows: (figure 5) when p53 begins to accumulate as a response to stress signals MDM2 first rises since it is positively regulated by this gene. The level of miR-34a begins also to increase since it is a direct target gene of p53. When the expression of miR-34a reaches some threshold MDM2 begins to decrease. It has been demonstrated that the response time of the I1-FFL is smaller than the one of a single regulation system.⁶⁴ In addition to this speedup

feature the I1-FFL can produce non-monotonic response also called an amplitude filter or bi-phasic response. In this type of response, the output MDM2 first increases with the input signal that activates p53 (stress signal), but decreases when the signal is high.⁶⁵ Thus in cases where speedy responses are needed this type of regulatory loop is more advantageous than the simple one. In our case MDM2 is the principal cell antagonist of p53 that limits its anti growth function in normal cells, and the I1-FFL constitute a novel mechanism that accelerates the stabilization of p53 in response to external stimulus.




Figure 4 Type I incoherent FFL composed of p53, miR-34a and MDM2.




Figure 5 The dynamics of the I1-FFL.

Acknowledgements

None.

Conflict of interest

Author declares that there is no conflict of interest.

References

1. Lane DP. Cancer. p53, guardian of the genome. *Nature*. 1992;358(6381):15–16.

2. Levine AJ. p53, the cellular gatekeeper for growth and division. *Cell*. 1997;88(3):323–331.
3. Levine AJ, Finlay CA, Hinds PW. P53 is a tumor suppressor gene. *Cell*. 2004;116:S67–S69.
4. Vousden KH, Prives C. Blinded by the light: The growing complexity of p53. *Cell*. 2009;137(3):413–431.
5. Feng Z, Levine AJ. The regulation of energy metabolism and the IGF-1/mTOR pathways by the p53 protein. *Trends Cell Biol*. 2010;20(7):427–434.
6. Baker SJ, Fearon ER, Nigro JM, et al. Chromosome 17 deletions and p53 gene mutations in colorectal carcinomas. *Science*. 1989;244(4901):217–221.
7. Vogan K, Bernstein M, Leclerc JM, et al. Absence of p53 gene mutations in primary neuroblastomas. *Cancer Res*. 1993;53(21):5269–5273.
8. Yee KS, Vousden KH. Complicating the complexity of p53. *Carcinogenesis*. 2005;26(8):1317–1322.
9. Riley T, Sontag E, Levine A, et al. Transcriptional control of human p53-regulated genes. *Nat Rev Mol Cell Biol*. 2008;9(5):402–412.
10. Green DR, Kroemer G. Cytoplasmic functions of the tumour suppressor p53. *Nature*. 2009;458(7242):1127–1130.
11. Kubbutat MH, Jones SN, Vousden KH. Regulation of p53 stability by MDM2. *Nature*. 1997;387(6630):299–303.
12. Honda R, Tanaka H, Yasuda H. Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53. *FEBS Lett*. 1997;420(1):25–27.
13. Appella E, Anderson CW. Post-translational modifications and activation of p53 by genotoxic stresses. *Eur J Biochem*. 2001;268(10):2764–2772.
14. Malik, Yousef, Allmer Jens. *miRNomics : MicroRNA Biology and Computational Analysis*. Springer; 2014.
15. Denli AM, Tops BB, Plasterk RH, et al. Processing of primary microRNAs by the Microprocessor complex. *Nature*; 2004;432(7014):231–235.
16. Gregory RI, Yan KP, Amuthan G, et al. The Microprocessor complex mediates the genesis of microRNAs. *Nature*. 2004;432(7014):235–240.
17. Han J, Lee Y, Yeom KH, et al. The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev*. 2004;18(24):3016–3027.
18. Lund E, GÅijtinger S, Calado A, et al. Nuclear export of microRNA precursors. *Science*. 2004;303(5654):95–98.
19. Yi R, Qin Y, Macara IG, et al. mediates the nuclear export of pre-microRNAs and short hairpin rnas. *Genes Dev*. 2003;17(24):3011–3016.
20. Gregory RI, Chendrimada TP, Cooch N, et al. Human RISC couples microRNA biogenesis and posttranscriptional gene silencing. *Cell*. 2005;123(4):631–640.
21. Hu H, Yan Z, Xu Y, et al. Sequence features associated with microRNA strand selection in humans and flies. *BMC Genomics*. 2005;10:413.
22. Okamura K, Balla S, Martin R, et al. Two distinct mechanisms generate endogenous siRNAs from bidirectional transcription in drosophila melanogaster. *Nat Struct Mol Biol*. 2005;15(6):581–590.
23. Yang JS, Phillips MD, Betel D, et al. Wide spread regulatory activity of vertebrate microRNA* species. *RNA* 2011;17(2):312–326.
24. Ro S, Park C, Young D, et al. Tissue-dependent paired expression of miRNAs. *Nucleic Acids Res*. 2007;35(17):5944–5953.
25. Okamura K, Phillips MD, Tyler DM, et al. The regulatory activity of microRNA* species has substantial influence on microRNA and 3' UTR evolution. *Nat Struct Mol Biol*. 2008;15(4):354–363.
26. Chiang HR, Schoenfeld LW, Ruby JG, et al. Mammalian microRNAs: experimental evaluation of novel and previously annotated genes. *Genes Dev*. 2010;24(10):992–1009.
27. Griffiths Jones S, Hui JHL, Marco A, et al. MicroRNA evolution by arm switching. *EMBO Rep*. 2011;12(2):172–177.
28. Marco A, Macpherson JI, Ronshaugen M, et al. MicroRNAs from the same precursor have different targeting properties. *Silence*. 2012;3(1):8.
29. Packer AN, Xing Y, Harper SQ, et al. The bifunctional microRNA miR-9/miR-9* regulates REST and Co-REST and is downregulated in huntington's disease. *J Neurosci*. 2008;28(53):14341–14346.
30. Shan SW, Lee DY, Deng Z, et al. MicroRNA MiR-17 retards tissue growth and represses fibronectin expression. *Nat Cell Biol*. 2009;11(8):1031–1038.
31. Almeida MI, Nicoloso MS, Zeng L, et al. Strand-specific miR-28-5p and miR-28-3p have distinct effects in colorectal cancer cells. *Gastroenterology*. 2012;142(4):886–896.
32. Byrd AE, Aragon IV, Brewer JW. MicroRNA-30c-2* limits expression of proadaptive factor XBP1 in the unfolded protein response. *J Cell Biol*. 2012;196(6):689–698.
33. Jiang L, Huang Q, Zhang S, et al. Hsa-miR-125a-3p and hsa-miR-125a-5p are down regulated in non-small cell lung cancer and have inverse effects on invasion and migration of lung cancer cells. *BMC Cancer*. 2010;10:318.
34. Wu H, Ye C, Ramirez D, et al. Alternative processing of primary microRNA transcripts by Drosha generates 5' End variation of mature microRNA. *PLoS One*. 2009;4:e7566.
35. Zhou H, Huang X, Cui H, et al. miR-155 and its star-form partner miR-155* cooperatively regulate type I interferon production by human plasmacytoid dendritic cells. *Blood*. 2010;116(26):5885–5894.
36. Kim S, Lee UJ, Kim MN, et al. MicroRNA miR-199a* regulates the MET proto-oncogene and the downstream extracellular signal-regulated kinase 2 (ERK2). *J Biol Chem*. 2008;283(26):18158–18166.
37. Sakurai K, Furukawa C, Haraguchi T, et al. MicroRNAs miR-199a-5p and -3p target the Brm subunit of SWI/SNF to generate a double-negative feedback loop in a variety of human cancers. *Cancer Res*. 2011;71:1680–1689.
38. Kuchenbauer F, Mah SM, Heuser M, et al. Comprehensive analysis of mammalian miRNA* species and their role in myeloid cells. *Blood*. 2011;118(12):3350–3358.
39. Montag J, Hitt R, Opitz L, et al. Upregulation of miRNA hsa-miR-342-3p in ex-perimental and idiopathic prion disease. *Mol Neurodegener*. 2009;4:36.
40. Grimson A, Srivastava M, Fahey B, et al. Early origins and evolution of microRNAs and Piwi-interacting RNAs in animals. *Nature*. 2008;455:1193–1197.
41. Tsang WP, Kwok TT. The miR-18a* microRNA functions as a potential tumor suppressor by targeting on K-Ras. *Carcinogenesis*. 2009;30(6):953–959.
42. Uchino K, Takeshita F, Takahashi RU, et al. Therapeutic effects of microRNA-582-5p and -3p on the inhibition of bladder cancer progression. *Mol Ther*. 2013;21:610–619.
43. Hermeking H. The miR-34 family in cancer and apoptosis. *Cell Death Differ*. 2010;17(2):193–199.
44. Chang TC, Wentzel EA, Kent OA, et al. Transactivation of miR-34a by p53 broadly influences gene expression and promotes apoptosis. *Mol Cell*. 2007;26(5):45–752.

45. Corney DC, Flesken Nikitin A, Godwin AK, et al. MicroRNA-34b and microRNA-34c are targets of p53 and cooperate in control of cell proliferation and adhesion-independent growth. *Cancer Res.* 2007;67(18):8433–8438.
46. Raver Shapira N, Marciano E, Meiri E, et al. Transcriptional activation of miR-34a contributes to p53-mediated apoptosis. *Mol Cell.* 2007;26(5):731–743.
47. Tazawa H, Tsuchiya N, Izumiya M, et al. Tumor-suppressive miR-34a induces senescence-like growth arrest through modulation of the E2F pathway in human colon cancer cells. *Proc Natl Acad Sci USA.* 2007;104(39):15472–15477.
48. Tarasov V, Jung P, Verdoodt B, et al. Differential regulation of microRNAs by p53 revealed by massively parallel sequencing: miR-34a is a p53 target that induces apoptosis and G1-arrest. *Cell Cycle.* 2007;6(13):1586–1593.
49. Bommer GT, Gerin I, Feng Y, et al. p53-mediated activation of miRNA34 candidate tumor-suppressor genes. *Curr Biol.* 2007;17(15):1298–1307.
50. He L, He X, Lim LP, et al. A microRNA component of the p53 tumour suppressor network. *Nature.* 2007;447(7148):1130–1134.
51. Bentwich I, Avniel A, Karov Y, et al. Identification of hundreds of conserved and noncon-served human microRNAs. *Nat Genet.* 2005;37(7):766–770.
52. Lim LP, Lau NC, Garrett Engele P, et al. Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature.* 2005;433(7027):769–773.
53. Griffiths Jones S, Grocock RJ, Van Dongen S, et al. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids.* 2006;34:D140–D144.
54. Hu W, Chan CS, Wu R, et al. Negative regulation of tumor suppressor p53 by microRNA miR-504. *Mol Cell.* 2010;38(5):689–699.
55. Le MT, Teh C, Shyh Chang N, et al. MicroRNA-125b is a novel negative regulator of p53. *Genes Dev.* 2009;23(7):862–876.
56. Park SY, Lee JH, Ha M, et al. miR-29 miRNAs activate p53 by targeting p85 alpha and CDC42. *Nat Struct Mol Biol.* 2009;16(1):23–29.
57. Yamakuchi M, Ferlito M, Lowenstein CJ. miR-34a repression of SIRT1 regulates apoptosis. *Proc Natl Acad Sci USA.* 2008;105(36):13421–13426.
58. Luo J, Nikolaev AY, Imai S, et al. Negative control of p53 by Sir2alpha promotes cell survival under stress. *Cell.* 2001;107(2):137–148.
59. Vlachos, Konstantinos Zagganas, Ioannis S, et al. DIANA-miRPath v3.0: deciphering microRNA function with experimental support. *Nucleic Acids research.* 2015;43(1):460–466.
60. Shannon P, Markiel A, Ozier O, et al. Cytoscape: a soft-ware environment for integrated models of biomolecular interaction networks. *Genome Research.* 2003;13(11):2498–2504.
61. S Maere, K Heymans, M Kuiper. BiNGO: a cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks. *Bioinformatics.* 2005;21(16):3448–3449.
62. Subhasree Nag, Jiangjiang Qin, Kalkunte S, et al. The MDM2-p53 pathway revisited. *J Biomed Res.* 2013;27(4):254–271.
63. Petrenko O, Moll UM. The MDM2-p53 interaction. *Mol Cancer Res.* 2003;1(14):1001–1008.
64. U Alon S, Mangan. Structure and function of the feed-forward loop network motif. *PNAS.* 2003;100(21):11980–11985.
65. Shai Kaplan, Anat Bren, Erez Dekel, et al. The incoherent feed-forward loop can generate non-monotonic input functions for genes. *Mol Syst Biol.* 2008;4:203.