

Dendritic Cell Vaccine and its Application in Cancer Therapy

Abstract

Dendritic cells (DCs) are the most potent antigens presenting cells in the immune system. They have a high capacity to trigger antigen-specific immune responses and promote both adaptive immunity and innate immunity. In the past decade, DC vaccine has been introduced as a new therapeutic strategy in cancer patients. DC-based immunotherapy is safe and can promote antitumor immune responses and prolonged survival of cancer patients. However, the current approaches of DC vaccination are far from optimal efficacy in advanced cancers. In this paper, we present recent findings about DC vaccine, its clinical application and efficacy in various cancers, as well as improved approaches in the preparation of DC vaccine.

Keywords: Dendritic cells; Function; *Ex vivo* generation; Vaccine; Cancer treatment; Efficacy

Mini Review

Volume 1 Issue 1 - 2015

Farashi-Bonab S¹ and Khansari N^{2*}

¹Department of Immunology, Tehran University of medical Sciences, Iran

²American Medical Diagnostic Laboratory, USA

***Corresponding author:** Nemat Khansari, American Medical Diagnostic Laboratory, 1665 Garden Grove Blvd, Garden Grove CA 92843, USA, Tel: +1(949) 228-8290, Email: nkhangari928@gmail.com

Received: April 7, 2015 | **Published:** May 18, 2015

Abbreviations: DC: Dendritic Cell; APC: Antigen Presenting Cell; PRR: Pathogen Recognition Receptor; TLR: Toll Like Receptor; CLR: C-Type Lectin Receptor; MHC: Major Histocompatibility Complex; cDC: Conventional DC; pDC: Plasmacytoid DC; iDCs: Inflammatory DC; ICAM-1: Intracellular Adhesion Molecule-1; LFA-3: Lymphocyte-Function Associated Antigen-3; IL: Interleukin; NK: Natural Killer; NKT: Natural Killer T; PD-L: Programmed Death-Ligand; TRAIL: TNF-Related Apoptosis-Induced Ligand; TGF- β : Transforming Growth Factor- β ; IDO: Indoleamine 2,3-Dioxygenase; NO: Nitric Oxide; Treg: Regulatory T Cell; Breg: Regulatory B Cell; GM-CSF: Granulocyte-Macrophage Colony Stimulating Factor; TAA: Tumor Associated Antigen; TSA: Tumor Specific Antigen; Flt3: Fms Like Tyrosine Kinase 3; LPS: Lipopolysaccharide; BCG: Bacillus Calmette-Guerrin; IFN: Interferon; TNF- α : Tumor Necrosis Factor- α ; U.S. FDA: United States Food and Drug Administration; CMV: Cytomegalovirus; PAP: Prostatic Acid Phosphatase; MART-1: Melanoma Antigen Recognized By T Cell-1; NY-Eso-1: New York Esophageal Squamous Cell Carcinoma-1

Introduction

Cancer cells recruit several mechanisms to escape from immune systems [1]. Innovation of approaches to increase antitumor reactivity of effector immune cells is essential for optimal antitumor immunity. DCs, as the most potent APCs, have a major application in cancer immunotherapy by presentation of tumor specific/associated antigens, activating antitumor lymphocytes, and augmenting innate immunity. More than two decades before, DCs were generated in the culture medium from bone precursor cells [2]. Afterwards, DC-based vaccines were studied in order to use as a therapeutic or prophylactic tool in malignant disorders. At present, DC vaccines are offered as a valuable instrument in cancer treatment. In this paper, we discuss anticancer DC vaccines, their application, safety, and clinical efficacy, as well as recent advances in DC vaccine preparation.

Immunophenotype and function of DCs

DCs are a heterogeneous population of cells with a widespread

tissue distribution. They have a distinctive morphology with many veil-like projections and express PRRs, such as TLRs and CLRs, the endocytic receptor DEC-205, and FC γ receptors. DCs also express 10-100 fold higher levels of peptide-MHC complexes than other professional APCs, i.e. monocytes and B cells, on the cell surface. Indeed, DCs have a high ability in receptor-mediated endocytosis and macropinocytosis, antigen processing and presentation on the MHC class I and II molecules [3,4]. They promote induction of antigen-specific immune responses and also augment innate immunity [5,6]. DCs are also important in induction of Immunological tolerance [7]

DC subsets

Myeloid DCs and plasmacytoid DCs: DCs can be categorized into two major subsets based on their origin: myeloid or conventional DCs (cDCs) and plasmacytoid DCs (pDCs). cDCs are derived from myeloid progenitor cells in the bone marrow and are characterized by expression of CD11c. cDCs can be subdivided into monocyte-derived DCs, CD1a⁻ interstitial DCs, and CD1a⁺ Langerhans cells. pDCs differentiate from lymphoid progenitor cells in the lymphoid tissues and characterized by expression of CD123 and production of a high levels of type I interferon. During inflammatory response, inflammatory DCs (iDCs) are originated from monocytes [8].

Immature DCs and Mature DCs: DCs are classified into "immature" and "mature" subsets based on their immunophenotype and function. Immature DCs have high capacity to capture antigens and are usually found in the peripheral non-lymphoid tissues. After antigen capture, they migrate into lymphoid tissues and become matured by up-regulation of expression of MHC molecules I and II, co-stimulatory molecules such as CD40, CD80, and CD86, and adherent molecules such as ICAM-1(CD54) and LFA-3(CD58) on the cell surface as well as secrete various soluble factors including various chemokines and cytokines such as IL-12. Consequently, mature DCs can present antigens in the lymphoid tissues and prime, activate, and expand effector immune cells [5,6,9].

Immuno-stimulatory DCs and tolerogenic DCs: DCs have strong immune-stimulatory properties. They can activate CD4⁺ and CD8⁺ T cells, B cells, NK cells, NKT cells, $\gamma\delta$ T cells, and other immune cells. But, DCs have also important roles in the induction of immunological tolerance through various mechanisms including expression of the immunosuppressive molecules such as PD-L1 and -L2, CD95 (Fas), TRAIL, and galectin-1, and secretion of immunosuppressive mediators such as IL-10, TGF- β , IDO, IL-27, arginase-1, heme oxygenase-1, and NO. Both pDC and cDCs can be tolerogenic. These tolerogenic DCs suppress immune responses of different effector cells; for example, they induce T cell anergy or apoptosis and promote the development and expansion of Tregs and Bregs [1,10].

Ex vivo generation of DCs

Differentiation of DCs from DC-precursor cells: DCs exist in a low frequency in the peripheral blood, at about 0.1% of white cells. Isolation of DCs from blood or other tissues is difficult due to the low frequency of these cells. Currently, DCs are generated from precursor cells at large numbers *ex vivo* [11-13]. We and others generated DCs *ex vivo* with a high purity [14]. Various agents such as GM-CSF, Flt3 ligand, c-Kit, IL-3, and IL-4 have been used to differentiate and growth DCs *ex vivo*. Human DCs are usually generated *ex vivo* from peripheral blood CD14⁺ monocytes in the presence of GM-CSF or GM-CSF/IL-4. Also, CD34⁺ hematopoietic stem cells can be used for *ex vivo* generation of DCs. Mouse DCs are usually generated from bone marrow precursor cells.

Antigen loading of DCs: Antigen loading of DCs is an important process in *ex vivo* generation of DCs for clinical applications. TSAs/TAAAs are suitable targets for loading on DCs. Synthetic peptides have been used for antigen loading of DCs [15-17]. However, synthetic peptides are only accessible for cancers with defined TSAs/TAAAs. Tumor lysate is other approach for antigen loading of DCs. This approach has been used successfully in many studies [18-21]. Total antigen loading approaches have the potency to deliver self-antigens and immunosuppressive components, and ultimately reduce antitumor immune responses. Nevertheless, total tumor antigen vaccines contain all potential tumor specific antigens, both MHC class I- and MHC class II- restricted epitopes, therefore, these vaccines can produce multivalent CD4⁺ and CD8⁺ T cell responses. DCs pulsed with tumor lysate can expand autologous tumor reactive T cells, *in vitro*. Whole tumor lysate based vaccines also produced more clinical responses than highly specific vaccines. Recently, hypochlorous acid-oxidation has been presented as a new method for preparing tumor lysate and DCs pulsed with hypochlorous acid-oxidized tumor lysate were safe and elicited potent T cell responses against ovarian antigens [22]. DC loading of cancer stem cell lysate is appeared to be more effective than total tumor lysate [23]. DCs have also been successfully transfected with peptide- or total tumor-RNA/mRNA or DNA [24-26]. Tumor cells RNA- or mRNA-transfected DCs have induced CTLs and antitumor immunity [24,27-29]. In a recent study, total tumor mRNA-electroporated DCs were more potent than total tumor lysate-electroporated DCs in the induction of antitumor immune responses *in vitro* and in suppression of tumor growth in MC-38-carcinoembryonic antigen colon cancer-bearing mice [30].

Maturation induction in DCs: Multiple components induce maturation of DCs. TLR ligands, including microbial components such as LPS, peptidoglycan, cholera toxin, filamentous hemagglutinin, inactivated BCG; and viral double stranded RNA, cytokines such as type I interferons, IFN- γ , TNF- α , IL-1 β , TGF- β , and IL-10, PGE₂, vitamin D₃, FC γ receptors, necrotic cells, apoptotic bodies, heat shock proteins, urate crystals, T cells, NK cells, NKT cells, and $\gamma\delta$ T cells can stimulate/modulate maturation of DCs. However, IL-12 production by DCs, which have important roles in tumor immunity, is only induced by some factors. IFN- γ and IL-4 enhance IL-12 production while PGE₂ and IL-10 have inhibitory effects. IL-1 β , IL-6, TNF- α , and PGE₂ have been used as a golden standard for maturation of DCs in cancer immunotherapy. But, these agents can reduce the production of IL-12 of DCs [31,32]. Furthermore, IL-10, vitamin D₃, and the drugs dexamethasone and rapamycin induce immature DCs/tolerogenic DCs.

Route of DC vaccine administration, dose and repeats

The number of DCs, vaccination repeats, and the route of DC vaccination in patients influence the vaccine efficacy. In most studies, DCs have been injected intravenously, intraperitoneally, subcutaneously, or intradermal. Intratumor injection of DCs can also be used for the induction of antitumor immune responses. In a study by Schmidt et al. [29] intratumoral injection of tumor RNA-pulsed DCs significantly produced more protective immunity as compared with subcutaneous or intravenous injections. In our study, intratumoral injection of un-mocked DCs into large established tumor led to enhanced tumor growth (our unpublished data), indicating that the tumor microenvironment has an enormous adverse effect on DCs. It should also be noticed that antigen-loaded DCs when injected to previously vaccinated hosts may be rapidly eliminated by antigen-specific CTLs [33]. Thus, increased numbers of DC vaccinations may not lead to enhanced T cell activation/expansion. On the other hand, once antigen-specific T cells in the periphery diminish to a level that they do not eliminate DCs; DCs can again re-stimulate T cell responses, emphasizing considering appropriate DC injection intervals. The frequency of DC vaccine injections should also be taken into account. Although repeated immunizations lead to increased frequencies of memory T cells, overstimulation can result in terminal differentiation and activation induced cell death (AICD) or exhaustion of T cells [34].

Antitumor DC vaccines in animal models

In 1989, Shimizu et al. [35] reported that vaccination with APCs pulsed with tumor antigens induces protective antigen-specific antitumor immunity in mice [35]. Since the mid-1990s, tumor antigen-pulsed DCs have been studied to treat tumors in mouse tumor models. In 1994, Flamand et al. [36] observed that immunization with DCs pulsed with tumor antigen protected mice against subsequent tumor challenge in a B cell lymphoma model. Afterwards, induction of antitumor immunity was reported by vaccination with DCs pulsed with tumor peptide [15-17] and soluble protein [37] in different tumor models. DCs fused to tumor cells are able to stimulate naïve T cells and induction of protective immunity *in vivo* [38]. Tumor lysate-pulsed DCs (18-21), and tumor cell RNA- or mRNA-transfected DCs (24,27-

29) have been capable to induce CTLs and antitumor immunity in several tumor models. In a myeloma mouse model, both idiotype protein- and tumor lysate-pulsed DC vaccines induced strong antitumor immune responses and protected mice against myeloma, but tumor lysate-pulsed DC vaccine was more potent than idiotype-pulsed DC vaccine to promote antitumor immunity [39]. In a recent study, total tumor mRNA-electroporated DCs were more potent than total tumor lysate-electroporated DCs in the induction of antitumor immune responses *in vitro* and suppression of tumor growth in MC-38-carcinoma antigen colon cancer-bearing mice [30].

Application of DC vaccine in cancer patients and its efficacy

In the past decade, increased knowledge about DCs and the possibility of generation of large numbers of DCs *ex vivo*, led to use DC vaccines in cancer patients as a novel cancer therapy modality. Cancer immunotherapy with DCs represents greatest hope for patients with advanced cancers, which do not respond to the conventional cancer therapies. Vaccination with DCs pulsed with peptides or tumor lysates have induced therapeutic tumor-specific responses in some malignancies. DCs pulsed with tumor antigens have been well tolerated and autoimmunity has not been observed in vaccinated patients. In 1998, therapeutic responses were reported after vaccination with DCs pulsed with tumor lysate or a mixed of peptides in five of 16 patients with advanced melanoma; two patients with complete response and three patients with partial response [40]. Since 2000, clinical trials have been performed using DCs pulsed with tumor peptides or tumor lysate and DC-tumor cell hybrids in patients with melanoma, colorectal cancer, neuroblastoma, cutaneous T cells lymphoma, renal cell carcinoma, lung cancer, breast cancer, sarcoma, leukemia, pancreatic adenocarcinoma, cholangiocellular carcinoma, hepatocellular carcinoma, thyroid medullary carcinoma, non-Hodgkin's lymphoma, multiple myeloma, and prostate cancer. Immunological responses were induced in most vaccinated patients without obvious toxicity or autoimmunity. However, complete or partial clinical responses were rare.

Sipuleucel-T is a DC vaccine which has been approved by the U.S. FDA in 2010 to treat patients with asymptomatic, or minimally symptomatic, metastatic hormone-refractory prostate cancer. This vaccine is the only DC vaccine that approved so far by U.S. FDA to treat cancer [41]. Sipuleucel-T helps to extend patient's lives by several months. This vaccine is composed from monocyte-derived autologous DCs that pulsed with the tumor antigen PAP.

In recent years, advancing the DC vaccine preparation protocols or combinational therapy has increased the efficacy of DC vaccines. PD-L-silenced and antigen mRNA-loaded DCs had improved immunogenic potency as they superiorly boosted *ex vivo* antigen-specific CD8⁺ T cell responses from transplanted cancer patients [42]. DCs pulsed with hypochlorous acid-oxidized tumor lysate were found to be safe and two of five vaccinated ovarian cancer patients experienced durable progression-free survival of 2 years and more after DC vaccination [22]. Vaccination of 16 patients with head and neck squamous cell carcinoma with DCs loaded with the

tumor peptide p53 was associated with two-year disease free survival of 88% of vaccinated patients [43]. Injection a tetanus/diphtheria toxoid in the vaccine site one day before vaccination with DCs pulsed with CMV phosphoprotein 65 RNA resulted in improved lymph node migration of DCs and prolonged survival of patients with glioblastoma. Preconditioning the vaccine site by this toxoid also enhanced DC vaccine efficacy in a mouse tumor model [44]. Administration of tumor lysate-pulsed DCs vaccine in combination with transfer of *ex vivo*-activated T cells after curative surgery led to prolonged recurrence-free survival and overall survival of patients with invasive hepatocellular carcinoma [45]. Vaccination with autologous MART-1-pulsed DCs together with adoptive transfer of TCR transgenic T cells resulted in tumor regression in nine of 13 patients with metastatic melanoma [46]. Patients with stage III/IV head and neck squamous cell carcinoma vaccinated intranodally with apoptotic tumor cell-loaded DCs survived disease-free for more than five years [47].

Conclusion

DC vaccine is a promising approach in cancer therapy. However, vaccine-induced clinical responses were not satisfactory in patients with advanced cancers. Thus, more attempts are needed to improve the efficacy of DC vaccines in advanced cancers. DC generation protocol, route of antigen loading onto DCs, type and dose of tumor antigen(s), DC maturation status, DC migration potency, amount of cells administrated, number of injections, time of injections, and the route of vaccine administration can be optimized to improve clinical efficacy of DC vaccines as it is observed in more recent studies. Co-administration of agents that promote DCs survival and their immune stimulatory function may be beneficial. Furthermore, combinational therapy and modulation the immunosuppressive environment of the tumor, which suppresses antitumor activities of DCs, can increase DC vaccine efficacy.

References

1. Farashi-Bonab S, Khansari N (2014) Regulatory T cells in cancer patients and their roles in cancer development/progression. *MOJ Immunol* 1(4): 00024.
2. Inaba K, Inaba M, Romani N, Aya H, Deguchi M, et al. (1992) Generation of large numbers of dendritic cells from mouse bone marrow cultures supplemented with granulocyte/macrophage colony-stimulating factor. *J Exp Med* 176(6): 1693-1702.
3. Sallusto F, Cella M, Danieli C, Lanzavecchia A (1995) Dendritic cells use macropinocytosis and the mannose receptor to concentrate macromolecules in the major histocompatibility complex class II compartment: downregulation by cytokines and bacterial products. *J Exp Med* 182(2): 389-400.
4. Banchereau J, Briere F, Caux C, Davoust J, Lebecque S, et al. (2000) Immunobiology of dendritic cells. *Ann Rev Immunol* 18: 767-811.
5. Guermontprez P, Valladeau J, Zitvogel L, Thery C, Amigorena S (2002) Antigen presentation and T cell stimulation by dendritic cells. *Ann Rev Immunol* 20: 621-667.
6. Steinman RM, Hemmi H (2006) Dendritic cells: translating innate to adaptive immunity. *Curr Top Microbiol Immunol* 311: 17-58.
7. Banchereau J, Steinman RM (1998) Dendritic cells and the control of

- immunity. *Nature* 392(6673): 245-252.
8. Shortman K, Naik SH (2007) Steady-state and inflammatory dendritic-cell development. *Nat Rev Immunol* 7(1): 19-30.
 9. Purcell A, Elliott T (2008) Molecular machinations of the MHC-I peptide loading complex. *Curr Opin Immunol* 20(1): 75-81.
 10. Volchenkov R, Karlsen M, Jonsson R, Appel S (2013) Type 1 regulatory T cells and regulatory B cells induced by tolerogenic dendritic cells. *Scand J Immunol* 77(4): 246-254.
 11. Traver D, Akashi K, Manz M, Merad M, Miyamoto T, et al. (2000) Development of CD8alpha-positive dendritic cells from a common myeloid progenitor. *Science* 290(5499): 2152-2154.
 12. Manz MG, Traver D, Miyamoto T, Weissman IL, Akashi K (2001) Dendritic cell potentials of early lymphoid and myeloid progenitors. *Blood* 97(11): 3333-3341.
 13. Wu L, D'Amico A, Hochrein H, O'Keeffe M, Shortman K, et al. (2001) Development of thymic and splenic dendritic cell populations from different hemopoietic precursors. *Blood* 98(12): 3376-3382.
 14. Farashi-Bonab S, Salehi TZ, Khansari N, Hadjati J, Massoud A (2014) An optimized method for ex vivo generation of dendritic cells. *J Vet Res* 69(1): 17-23.
 15. Mayordomo JI, Zorina T, Storkus WJ, Zitvogel L, Celluzzi C, et al. (1995) Bone marrow-derived dendritic cells pulsed with synthetic tumour peptides elicit protective and therapeutic antitumour immunity. *Nat Med* 1(12): 1297-1302.
 16. Celluzzi CM, Mayordomo JI, Storkus WJ, Lotze MT, Falo LD (1996) Peptide-pulsed dendritic cells induce antigen-specific CTL-mediated protective tumor immunity. *J Exp Med* 183(1): 283-287.
 17. Porgador A, Snyder D, Gilboa E (1996) Induction of antitumor immunity using bone marrow-generated dendritic cells. *J Immunol* 156(8): 2918-2926.
 18. Cohen PJ, Cohen PA, Rosenberg SA, Katz SI, Mulé JJ (1994) Murine epidermal Langerhans cells and splenic dendritic cells present tumor-associated antigens to primed T cells. *Eur J Immunol* 24(2): 315-319.
 19. Geraghty PJ, Fields RC, Mule JJ (1996) Vaccination with tumor-pulsed splenic dendritic cells mediates immunity to a poorly immunogenic tumor. *Surg Forum* 47: 459-461.
 20. Fields RC, Shimizu K, Mule JJ (1998) Murine dendritic cells pulsed with whole tumor lysates mediate potent antitumor immune responses in vitro and in vivo. *Proc Natl Acad Sci USA* 95(16): 9482-9487.
 21. Chang AE, Redman BG, Whitfield JR, Nickloff BJ, Braun TM, et al. (2002) A phase I trial of tumor lysate-pulsed dendritic cells in the treatment of advanced cancer. *Clin Cancer Res* 8(4): 1021-1032.
 22. Chiang CL, Kandalaft LE, Tanyi J, Hagemann AR, Motz GT, et al. (2013) A dendritic cell vaccine pulsed with autologous hypochlorous acid-oxidized ovarian cancer lysate primes effective broad antitumor immunity: from bench to bedside. *Clin Cancer Res* 19(17): 4801-4815.
 23. Li Q, Lu L, Tao H, Xue C, Teitz-Tennenbaum S, et al. (2014) Generation of a novel dendritic-cell vaccine using melanoma and squamous cancer stem cells. *J Vis Exp* 83: e50561.
 24. Boczkowski D, Nair SK, Snyder D, Gilboa E (1996) Dendritic cells pulsed with RNA are potent antigen-presenting cells in vitro and in vivo. *J Exp Med* 184(2): 465-472.
 25. Condon C, Watkins SC, Celluzzi CM, Thompson K, Falo LD (1996) DNA-based immunization by in vivo transfection of dendritic cells. *Nat Med* 2(10): 1122-1128.
 26. Manickan E, Kanangat S, Rouse RJ, Yu Z, Rouse BT (1997) Enhancement of immune response to naked DNA vaccine by immunization with transfected dendritic cells. *J Leukoc Biol* 61(2): 125-132.
 27. Heiser A, Maurice MA, Yancey DR, Wu NZ, Dahm P, et al. (2001) Induction of polyclonal prostate cancer-specific CTL using dendritic cells transfected with amplified tumor RNA. *J Immunol* 166(5): 2953-2960.
 28. Milazzo C, Reichardt VL, Muller MR, Grunebach F, Brossart P (2003) Induction of myeloma-specific cytotoxic T cells using dendritic cells transfected with tumor-derived RNA. *Blood* 101(3): 977-982.
 29. Schmidt T, Ziske C, Märten A, Endres S, Tiemann K, et al. (2003) Intratumoral immunization with tumor RNA-pulsed dendritic cells confers antitumor immunity in a C57BL/6 pancreatic murine tumor model. *Cancer Res* 63(24): 8962-8967.
 30. Osada T, Nagaoka K, Takahara M, Yang XY, Liu CX, et al. (2015) Precision Cancer Immunotherapy: Optimizing Dendritic Cell-based Strategies to Induce Tumor Antigen-specific T-cell Responses Against Individual Patient Tumors. *J Immunother* 38(4): 155-164.
 31. Káliński P, Schuitemaker JH, Hilkens CM, Wierenga EA, Kapsenberg ML (1999) Final maturation of dendritic cells is associated with impaired responsiveness to IFN-gamma and to bacterial IL-12 inducers: decreased ability of mature dendritic cells to produce IL-12 during the interaction with Th cells. *J Immunol* 162(6): 3231-3236.
 32. Langenkamp A, Messi M, Lanzavecchia A, Sallusto F (2000) Kinetics of dendritic cell activation: impact on priming of TH1, TH2 and nonpolarized T cells. *Nat Immunol* 1(4): 311-316.
 33. Yang J, Huck SP, McHugh RS, Hermans IF, Ronchese F (2006) Perforin-dependent elimination of dendritic cells regulates the expansion of antigen-specific CD8⁺ T cells in vivo. *Proc Natl Acad Sci USA* 103(1): 147-152.
 34. Wherry E (2011) T cell exhaustion. *Nat Immunol* 12(6): 492-499.
 35. Shimizu J, Suda T, Yoshioka T, Kosugi A, Fujiwara H, et al. (1989) Induction of tumor-specific in vivo protective immunity by immunization with tumor antigen-pulsed antigen-presenting cells. *J Immunol* 142(3): 1053-1059.
 36. Flamand V, Sornasse T, Thielemans K, Demanet C, Bakkus M, et al. (1994) Murine dendritic cells pulsed in vitro with tumor antigen induce tumor resistance in vivo. *Eur J Immunol* 24(3): 605-610.
 37. Paglia P, Chiodoni C, Rodolfo M, Colombo MP (1996) Murine dendritic cells loaded in vitro with soluble protein prime cytotoxic T lymphocytes against tumor antigen in vivo. *J Exp Med* 183(1): 317-322.
 38. Gong J, Chen D, Kashiwaba M, Kufe D (1997) Induction of antitumor activity by immunization with fusions of dendritic and carcinoma cells. *Nat Med* 3(5): 558-561.
 39. Hong S, Li H, Qian J, Yang J, Lu Y, et al. (2012) Optimizing dendritic cell vaccine for immunotherapy in multiple myeloma: tumour lysates are more potent tumour antigens than idiotype protein to promote anti-tumour immunity. *Clin Exp Immunol* 170(2): 167-177.
 40. Nestle FO, Alijagic S, Gilliet M, Sun Y, Grabbe S, et al. (1998) Vaccination of melanoma patients with peptide- or tumor lysate-pulsed dendritic cells. *Nat Med* 4(3): 328-332.
 41. Kantoff PW, Higano CS, Shore ND, Berger ER, Small EJ, et al. (2010) Sipuleucel-T immunotherapy for castration-resistant prostate cancer.

- N Engl J Med 363(5): 411-422.
42. Hobo W, Novobrantseva TI, Fredrix H, Wong J, Milstein S, et al. (2013) Improving dendritic cell vaccine immunogenicity by silencing PD-1 ligands using siRNA-lipid nanoparticles combined with antigen mRNA electroporation. *Cancer Immunol Immunother* 62(2): 285-297.
 43. Schuler PJ, Harasymczuk M, Visus C, Deleo A, Trivedi S, et al. (2014) Phase I dendritic cell p53 peptide vaccine for head and neck cancer. *Clin Cancer Res* 20(9): 2433-2444.
 44. Mitchell DA, Batich KA, Gunn MD, Huang MN, Sanchez-Perez L, et al. (2015) Tetanus toxoid and CCL3 improve dendritic cell vaccines in mice and glioblastoma patients. *Nature* 519(7543): 366-369.
 45. Shimizu K, Kotera Y, Aruga A, Takeshita N, Katagiri S, et al. (2014) Postoperative dendritic cell vaccine plus activated T-cell transfer improves the survival of patients with invasive hepatocellular carcinoma. *Hum Vaccin Immunother* 10(4): 970-976.
 46. Chodon T, Comin-Anduix B, Chmielowski B, Koya RC, Wu Z, et al. (2014) Adoptive transfer of MART-1 T-cell receptor transgenic lymphocytes and dendritic cell vaccination in patients with metastatic melanoma. *Clin Cancer Res* 20(9): 2457-2465.
 47. Whiteside TL, Ferris RL, Szczepanski M, Tublin M, Kiss J, et al. (2015) Dendritic cell-based autologous tumor vaccines for head and neck squamous cell carcinoma: Promise Vs reality. *Head Neck*.