

# Influenza-Associated Morbidity and Mortality in Sub-Saharan Africa

## Abstract

Influenza is one of the major infectious disease threats to the world. New Global Burden of Disease estimates attribute nearly 2% of all-cause mortality during the first five years of life to influenza. In tropical developing countries, the pattern of influenza circulation may differ markedly from those in temperate, developed settings. Of influenza deaths, 99% are estimated to occur in low and middle income countries. Studies on seasonal influenza occurrence, morbidity and mortality in sub-Saharan Africa published from January 1st, 2000 to December 31st, 2015 were reviewed in order to assess the burden of seasonal influenza in the Region. The starting date of January 1st, 2000 has been selected because of the increasing number and quality of studies in later years. Relevant studies were identified using combinations of Medical Subject Headings (MeSH) and text terms as follows: ("influenza" AND "Africa") OR ("Africa" AND ("pneumonia" OR "acute respiratory infection")) OR ("influenza" AND each individual sub-Saharan African country). The study seems to show that seasonal influenza could have a greater toll on morbidity and mortality in Africa, compared with more-developed continents considering the frequency of predisposing factors such as HIV and TB and the absence of influenza vaccination programs in most of the African countries. Available data suggest both the respiratory disease and influenza-associated mortality may be increased in low-income settings such as sub-Saharan Africa.

**Keywords:** Influenza; Morbidity; Mortality; Sub-saharan Africa; Respiratory; Predisposing

## Mini Review

Volume 2 Issue 4 - 2016

**Doudou Diop<sup>1\*</sup>, Emanuele Montomoli<sup>2</sup>,  
Melvin Sanicas<sup>3</sup> and Mayassine Diongue<sup>4</sup>**

<sup>1</sup>EPLS Biomedical Research Center, Senegal

<sup>2</sup>Department of Public Health, University of Siena, Italy

<sup>3</sup>Bill and Melinda Gates Foundation, USA

<sup>4</sup>Department of Public Health, Cheikh Anta Diop University, Senegal

**\*Corresponding author:** Doudou Diop, EPLS Biomedical Research Center, Saint-Louis, Senegal, Email: [deuxdou@gmail.com](mailto:deuxdou@gmail.com)

**Received:** November 21, 2015 | **Published:** August 29, 2016

**Abbreviations:** AIDS: Acquired Immune Deficiency Syndrome; AR: Attack Rate; ARI: Acute Respiratory Infections; aOR: Adjust Odd Ratio; CFP: Case Fatality Proportion; CFR: Case Fatality Rate; CI: Confidence Interval; DNA: Deoxyribonucleic Acid; HAART: Highly Active Antiretroviral Therapy; HIV: Human Immunodeficiency Virus; ILI: Influenza-Like Illness; NICD: National Institute for Communicable Disease; PCR: Polymerase Chain Reactions; PIV: Parainfluenza Virus; PTB: Pulmonary tuberculosis; RR: Relative Risk; rRT-PCR: Reverse Real Time PCR; RSV: Respiratory Syncytial Virus; SARI: Severe Acute Respiratory Infection; TB: Tuberculosis; TIV: Trivalent Inactivated Influenza; US CDC: US Centers for Disease Control and Prevention; NAMRU: Naval Medical Research Unit; WHO: World Health Organization

## Introduction

Influenza is a highly communicable acute respiratory disease. It is one of the most prevalent vaccine-preventable diseases. The influenza virus can cause illness in individuals of all ages, results in repeated infections throughout life, and is responsible for annual worldwide epidemics of varying severity. The World Health Organization (WHO) estimates that influenza affects between 3 and 5 million individuals each year, causing between 250,000 and 0.5 million deaths [1]. New Global Burden of Disease estimates attribute nearly 2% of all-cause mortality during the first five years of life to influenza [2].

In tropical countries, the pattern of influenza circulation may differ markedly from those in temperate, developed settings.

Also, populations in low and middle income countries like many of those in sub-Saharan Africa are more vulnerable to influenza related complications because of the high prevalence of underlying medical conditions such as Human Immunodeficiency Virus (HIV) infection, tuberculosis (TB), malaria and malnutrition, and limited access to healthcare. Of influenza deaths, 99% are estimated to occur in low and middle income countries [3]. This study aims at assessing the morbidity and mortality related to seasonal influenza in sub-Saharan Africa.

## Review Methods

### Information sources

Searches of the PubMed database were completed to identify studies on seasonal influenza in sub-Saharan Africa published in English and French from January 1st, 2000 to December 31st, 2015. The starting date of January 1st, 2000 has been selected because of the increasing number and quality of studies in later years.

### Literature search

Relevant studies were identified using combinations of Medical Subject Headings (MeSH) and text terms as follows: ("influenza" AND "Africa") OR ("Africa" AND ("pneumonia" OR "acute respiratory infection")) OR ("influenza" AND each individual sub-Saharan African country). References of identified articles have been examined for additional articles, titles and abstracts

of additional articles have been screened and studies have been selected if they included some aspect of seasonal influenza in sub-Saharan Africa.

### Study selection

Studies that reported data on the occurrence of influenza in sub-Saharan Africa and on its contribution to morbidity and mortality have been included. Studies that reported results from outside sub-Saharan Africa, that only had data on avian or animal influenza, that only reported data on pandemic influenza, that duplicated data on other more comprehensive reports have been excluded.

## Results and discussion

### Results

Nine hundred and forty five articles titles and abstracts have been screened, out of which 70 reports have been retrieved and assessed for eligibility criteria, of which 28 did not meet inclusion criteria. Forty-two articles were included in this study [4–45] (Figure 1).




Figure 1: Flow chart for studies selection.

### Morbidity and co-morbidity

**Morbidity:** Influenza surveillance conducted in 15 countries in Africa from 2006 to 2010 have shown 21.7% of influenza-like illness (ILI) cases (5165/69 860) tested positive for influenza, ranging from 6.7% in Angola to 40.4% in Madagascar [4]. In Senegal, from June 1996 to December 1998, the proportion of influenza virus isolation varied from 17.5% to 40% during the peak period (July/September) of acute respiratory infections (ARI) in Dakar [5]. These data are in contrast to temperate regions where influenza occurs during winter season. Clinical and virological surveillance of acute respiratory infections among children less than 5 years old living in rural areas in Senegal showed that influenza represented 45.6% of positive viral detection in patients with ARI [6]. The recent enhancement of the surveillance network in the country has showed that co-infection and co-circulation of viruses were frequent and were responsible of ILI peaks [7]. The first months of implementation of the enhanced surveillance system confirmed that Influenza viruses were among the most frequent cause of ILI (20%). The other viruses were adenoviruses (21%), rhinoviruses (18%) and enteroviruses (15%) [7]. In Antananarivo, Madagascar,

influenza epidemiologic and virologic surveillance from 1995 to 2002 had shown that the virus had been continually spreading all year long [8]. In another study conducted from July 2008 to June 2009, Influenza A and B accounted respectively for 27.3% and 3.9% of ILI cases. Of all viral detection, influenza A was the most common virus [9]. In Mozambique, viral ARI are frequent among infants visited in Manhiça District Hospital, the influenza viruses represented 15% of viral detection in infants younger than 12 months [10]. The most frequent virus was the rhinovirus (26%) [10]. However, in Cote d'Ivoire, the most affected people were from 15 to 59 years (47%) followed by those from 0 to 5 years (34%) [11].

Recently, many West African countries have published their first data on influenza surveillance. In Togo and Niger, the highest influenza positive percentage was observed in 5 - 14 years old with respectively (30%) among ILI in Togo [12] and 25% among ILI patients and 10% among severe acute respiratory infection (SARI) patients in Niger [13]. In Burkina Faso, of the 881 patients with ILI enrolled and sampled in the sentinel surveillance system, 58 (6.6%) tested positive for influenza viruses [14]. In contrast to the findings in Togo and Niger, the most affected were patients within 0 - 5 years (41.4%), the 6 - 14 years old were the second (22.4%) [14]. In Sierra Leone also the majority (36%) of ILI and SARI cases was detected in children under the age of four [15].

Surveillance of ILI in Central Africa began only recently. The surveillance systems in Gabon and Central African Republic have provided valuable data on the circulation of influenza viruses and other respiratory viruses. In Gabon, three-quarters of the patients were children under five years old [16]. The most common viruses were adenoviruses (17.5%), followed by parainfluenza viruses (PIVs) (16.8%), influenza viruses were detected only in 11.9% of tested specimens [16]. In Central African Republic, influenza viruses were the most common viruses (8.8%) followed by RSV (3.0%) [17]. However, the study included only infants and children aged 0–15 years [17].

In Djibouti, East Africa, in response to local reports of a possible outbreak during the A(H1N1)pdm09 pandemic, enhanced surveillance for ILI was conducted at Camp Lemonnier. rRT-PCR and DNA sequencing revealed that 25 (78%) of the 32 clinical samples collected were seasonal H3N2 and only 2 (6%) were A(H1N1)pdm09 influenza [18]. In South Africa, a country with a marked influenza season, the annual detection rate amongst specimens tested at the National Institute for Communicable Diseases (NICD) during the influenza seasons ranged from 32% in 2008 to 47% in 2005 and 2007 [19].

### Severe morbidity: Influenza related-hospitalizations

Several studies have reported high frequency of severe influenza among persons with ARI. In two recent studies from South Africa (Pretoria and Cape Town), influenza was detected respectively in 3.5% [20] and 0.9% [21] of children <5 years of age hospitalized for ARI. Another study from the pediatric service of Tygerberg Children's Hospital in Western Cape, has found influenza A and B respectively in 13% and 15% of children hospitalized for ARI [22]. In a study conducted in Cape Town from June 1st, 1995 to August 31st, 1996, influenza B was isolated

in 0.6% of children <2 years of age hospitalized for ARI [23]. A severe acute institutional influenza outbreak was reported from Pretoria, in a police residential college amongst new recruits and staff members at the end of May 2003. The outbreak had an attack rate (AR) per dormitory building from 27% to 47%, with an overall AR of 34% [24].

In a recent study conducted in two long term refugee camps in Kenya, influenza associated-SARI hospitalizations were 4.8/1000 in <5 years old and 11.1/1000 in <1 year old [25]. Influenza virus was also detected in 6% of children younger than 13 years of age admitted for clinical pneumonia to a Kenyan rural hospital [26]. In a population-based study conducted in rural western Kenya, of the 2079 patients with tested swabs, infection with influenza virus was confirmed in 204 (10%). Among those tested for influenza virus, 6.8% of the children aged < 5 years and 14.0% of the patients aged ≥ 5 years were found positive. The annual rate of hospitalization (per 100 000 population) was 56.2 among patients with influenza. The rate of influenza-associated hospitalization was highest among children aged less than 5 years [27].

In Nigeria, in a 30-month study of under 5-admissions for ARI, influenza A was found in 17.3% of children admitted with severe pneumonia radiologically-confirmed [28]. In a hospital-based study in the same age group in Ghana, influenza accounted only for 1% of respiratory viruses; the most common virus was RSV (14.1%) [29]. In a clinic and hospital-based sentinel influenza surveillance in Uganda, had showed that patients aged 5–14 years had the highest influenza-positive percentage (19.6%) [30]. Influenza circulated throughout the year, but the percentage of influenza-positive specimens peaked during June–November, coinciding with the second rainy season in Uganda [30].

### **Co-morbidity: influenza-HIV, influenza-chronic lung disease, influenza-meningitis, influenza-malaria**

The association between influenza and HIV has been studied in children aged from 2 months to 5 years admitted at Chris Hani-Baragwanath Hospital in Soweto, South Africa. The study has shown that children with HIV have an 8 times higher risk of admission to hospital from influenza-associated ARI than HIV negative children [31]. Children aged ≥5 Years and adults infected with HIV experienced a 13–19 times greater SARI incidence than HIV-uninfected individuals ( $p < 0.001$ ) [32]. In another study from Kenya, HIV-infection was associated with hospitalization due to influenza (aOR=3.56; 95% CI 1.25-10.1). Chronic lung disease was also associated with hospitalization due to influenza (aOR=6.83; 95% CI 1.37-34.0) [33]. In the absence of highly active antiretroviral therapy (HAART), adults aged 25-54 years with AIDS experience a substantially elevated risk of influenza-associated death, 150 to 200-fold higher than the general population of the same age, and 2-4 folds higher than seniors age 65 and over [34].

In Burkina Faso which is a country located in the heart of the African meningitis belt, the association between respiratory infections and meningococcal carriage has been studied. The study suggested that viral respiratory tract infections, including influenza, might contribute to increased transmission and nasopharyngeal carriage of virulent meningococci [35].

Although children <5 years old in sub-Saharan Africa are vulnerable to both malaria and influenza, little is known about coinfection. A retrospective, cross-sectional study in western rural Kenya has addressed this issue. The study found that 45% (149/331) of influenza-positive patients were coinfecting with malaria, whereas only 6% (149/2408) of malaria-positive patients were coinfecting with influenza [36]. Coinfection with malaria and influenza was uncommon but associated with longer hospitalization than single infections among children 24-59 months of age [36].

### **Mortality**

A recent study of elderly South Africans estimated that the risk of influenza-associated death in that country was three to four times higher than in the United States [37]. A three-year study of influenza and respiratory syncytial virus associated adult mortality in Soweto, has shown that Influenza seasons were significantly associated with excess mortality in adults across all 3 years. Excess mortality was highest in those ≥65 years of age: 82.8/100,000 populations in the mild 1997 season and 220.9/100,000 in the severe 1998 season [38].

Some papers reported case fatality-rate during influenza outbreak or through population based studies. In 2002, an influenza outbreak in Madagascar had a case-fatality rate of 3%; most fatalities occurred in young children [39,40]. Similarly high (3.5%) case-fatality rates among children <5 years of age were observed during an influenza outbreak in the Democratic Republic of Congo in 2002 [41]. In the population based- study conducted in rural western Kenya, the case-fatality rate among admitted patients with PCR-confirmed infection with influenza virus was 2.0% [27].

In a study of mortality amongst patients with influenza-associated SARI in Soweto, South Africa, the estimated incidence of influenza-associated SARI deaths per 100,000 population was highest in children <1 year (20.1, 95%CI 12.1-31.3) and adults aged 45–64 years (10.4, 95%CI 8.4–12.9) [42]. Adjusting for age, the rate of death was 20-fold (95%CI 15.0–27.8) higher in HIV-infected individuals than HIV-uninfected individuals [42]. This study has documented that influenza cause's substantial mortality in Soweto, South Africa. An analysis of data from the same surveillance program found that the relative risk of influenza-associated hospitalization was 4–8 times higher in HIV-infected compared to HIV-uninfected individuals [42].

The role of influenza in severe acute respiratory illness deaths in Sub-Saharan Africa was studied in 8 African countries. The data were collected from 37 714 SARI cases, and 3091 (8.2%; range by country, 5.1%-25.9%) tested positive for influenza virus [43]. There were 1073 deaths (2.8%; range by country, 0.1%-5.3%) reported, among which influenza virus was detected in 57 (5.3%) [43]. Case-fatality proportion (CFP) was higher among countries with systematic death reporting than among those with sporadic reporting. The influenza-associated CFP was 1.8% (57 of 3091), compared with 2.9% (1016 of 34 623) for influenza virus-negative cases ( $P < .001$ ) [43]. Among 834 deaths (77.7%) tested for other respiratory pathogens, rhinovirus (107 [12.8%]),

adenovirus (64 [6.0%]), respiratory syncytial virus (60 [5.6%]), and *Streptococcus pneumoniae* (57 [5.3%]) were most commonly identified. Among 1073 deaths, 402 (37.5%) involved people aged 0-4 years, 462 (43.1%) involved people aged 5-49 years, and 209 (19.5%) involved people aged  $\geq 50$  years [43].

A study of the influenza-associated mortality among individuals with PTB in South Africa from 1999–2009 observed an increased risk of influenza-associated mortality in persons with PTB compared to non-tuberculosis respiratory deaths [44]. In the same period of time (1999-2009), in South Africa, the mortality associated with seasonal and pandemic influenza among pregnant and non-pregnant women of childbearing age in a high HIV-prevalence setting was conducted. The estimated mean annual seasonal influenza-associated mortality rates were 12.6 (123 deaths) and 7.3 (914 deaths) among pregnant and non-pregnant women respectively [45]. Among pregnant women, the estimated mean annual seasonal influenza-associated mortality rates were 74.9 (109 deaths) among HIV infected and 1.5 (14 deaths) among HIV-uninfected individuals [45]. Among non-pregnant women, the estimated mean annual seasonal influenza-associated mortality rate was 41.2 (824 deaths) among HIV-infected and 0.9 (90 deaths) among HIV-uninfected individuals [45]. The study suggested that pregnant women experienced an increased risk of seasonal influenza-associated mortality compared with non-pregnant women (relative risk [RR], 2.8; 95% confidence interval [CI], 1.7-3.9) [45].

## Discussion

The burden of seasonal influenza in children under 5 years of age and those from 5 to 14 years appears far greater than in the elderly [11-16]. This corroborates findings elsewhere that younger children experience a high influenza attack rate and a higher rate of influenza-related complications [46]. In the study of Brooks et al. [47] 28% of influenza positive children developed pneumonia during their illness. The average annual incidence of influenza pneumonia was greatest among children < 5 years of age (236 per 100 000) [47]. Children are not only affected by influenza themselves; they play also an important role in introducing and spreading respiratory illness into households and communities [48]. They shed influenza virus for longer periods of time as compared to adults [49,50]. Also, children characteristically exercise suboptimal hand hygiene and sneezing/coughing precautions, making them efficient vectors of influenza spread throughout the community. Focusing influenza vaccination efforts on children may therefore be an effective and practical method of reducing the burden of influenza in sub-Saharan Africa.

The burden of ARI, already very large in stable settings, increases considerably in crises. Crises due to armed conflict, forced displacement and natural disasters result in excess morbidity and mortality due to infectious diseases. Risk factors related to overcrowding due to displacement into camps or into host households with overcrowded living and sleeping quarters could increase risk of influenza transmission. Interestingly, the study from Kenya [25] provided evidence that influenza is a major contributor to pneumonia cases in refugee populations.

The high prevalence of co-morbid conditions, including HIV infection, tuberculosis, and pregnancy, may contribute to increased disease severity and mortality from influenza in Africa [51]. There are approximately 30 million persons infected with HIV worldwide, including 5 million in South Africa only [34]. Published studies suggest that in the absence of highly active antiretroviral therapy (HAART), adults with AIDS experience a substantially elevated risk of influenza-associated death [34]. Also, children with HIV have higher risk of admission to hospital from influenza-associated ARI than HIV negative children [31]. HIV remains a substantial risk factor for severe influenza-related disease in sub-Saharan Africa. These findings support the ongoing importance of influenza prevention measures in persons with HIV/AIDS. The trivalent inactivated influenza (TIV) vaccine is protective against laboratory-confirmed symptomatic influenza in persons with moderate or asymptomatic HIV infection [52]. However, further evaluation of effectiveness is warranted in severely immune-compromised HIV-infected adults and those with co-morbidities such as tuberculosis.

On the other hand the absence of influenza vaccination program in most of the African countries may increase the risk of transmission of the virus. In developed countries the vaccination of risk groups reduces the prevalence of seasonal influenza; there is also a context of herd immunity which gives protection to unvaccinated people.

When thorough documentation was available the mortality related to influenza has been found also substantial in sub-Saharan Africa. Interestingly, based on the experience in Madagascar, that island was one of the few places in sub-Saharan Africa where a large outbreak was well documented with 114,000 deaths out of a population of 3,250,000 [53]. Thus, the worldwide distribution of these reports indicates that under conditions where high mortality was clustered in time and place, the impact of influenza could easily be recognized [53].

There are now established influenza surveillance systems in several African countries with support from the World Health Organization (WHO) and different organizations such as the US Centers for Disease Control and Prevention (USCDC), Naval Medical Research Unit (NAMRU), and Institut Pasteur [54]. Although many gaps remain, much more is known about seasonal influenza in sub-Saharan Africa now than ever before. Strengthening influenza surveillance, along with conducting special studies on the burden of influenza-associated hospitalization and mortality in relation to other etiologies of pneumonia as well as risk groups for severe and complicated disease, and the economic burden of influenza are important to guide policy-makers in their allocation of scarce resources for related public health interventions.

## Conclusion

The burden of seasonal influenza in sub-Saharan Africa seems unlikely to be less than in other areas as usually thought. Seasonal influenza is a major contributor to childhood pneumonia. But, its contribution to early childhood pneumonia appears still under-appreciated in high pneumonia-endemic tropical settings such as

sub-Saharan Africa for several reasons. Also, there is good reason to believe that seasonal influenza could also have a greater toll on morbidity and mortality in Africa, compared with more-developed continents considering the frequency of predisposing factors such as HIV and TB and the absence of influenza vaccination program. When documentation was available, the mortality related to the disease has been found also substantial.

Despite the growing successes of the influenza surveillance systems in sub-Saharan Africa, they need to be strengthened and expanded in order to address some of the data gaps. The strengthening of national surveillance capacity supports the long-term goal of generating improved data for national decision-making.

### Study limitations

The authors acknowledge the following limitations of the study: Firstly, the studies reviewed did not use standardized case definitions of ILI and SARI. Secondly, data on outpatient and inpatient proportional morbidity and mortality may not be representative of population patterns due to differences in health care utilization by type of illness. Studies varied substantially on the basis of the number of years of assessment, age-groups included, number of influenza isolates obtained, and testing methods implemented.

### References

1. WHO (2015) Influenza (Seasonal). World Health Organization.
2. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, et al. (2012) Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380(9859): 2095-2128.
3. Nair H, Brooks WA, Katz M, Roca A, Berkley JA, et al. (2011) Global burden of respiratory infections due to seasonal influenza in young children: a systematic review and meta-analysis. *Lancet* 378(9807): 1917-1930.
4. Radin JM, Katz MA, Tempia S, Nzussouo TN, Davis R, et al. (2012) Influenza surveillance in 15 countries in Africa, 2006-2010. *JID In press. The Journal of Infectious Disease* 206(1): S14-S21.
5. Dosseh A, Ndiaye K, Spiegel A, Sagna M, Mathiot C (2000) Epidemiological and virological influenza survey in Dakar, Senegal: 1996-1998. *Am J Trop Med* 62(5): 639-643.
6. Niang MN, Diop ON, Sarr FD, D Goudiaby, Sompy MH, et al. (2010) Viral Etiology of Respiratory Infections in Children Under 5 Years Old Living in Tropical Rural Areas of Senegal: The EVIRA Project. *J Med Virol* 82(5): 866-872.
7. Dia N, Sarr FD, Thiam D, Sarr TF, Espie E, et al. (2014) Influenza-Like Illnesses in Senegal: Not Only Focus on Influenza Viruses. *PLoS One* 9(3): e93227.
8. Rabarijaona LP, Rakotondrarija NT, Rousset D, Soares JL, Maucière P (2003) Surveillance épidémiologique et virologique de la grippe à Antananarivo de 1995 à 2002. *Arch Inst Pasteur de Madagascar* 69(1&2): 20-26.
9. Razanajatovo NH, Richard V, Hoffmann J, Reynes JM, Razafitrimo GM, et al. (2009) Viral etiology of Influenza-Like Illnesses in Antananarivo, Madagascar, July 2008 to June 2009. *PLoS One* 6(3): e17579.
10. Gordo CC, Padrisa DN, Abacassamo F, Brena PP, Casas I, et al. (2011) Viral acute respiratory infections among infants visited in a rural hospital of southern Mozambique. *Trop Med Int Health* 16(9): 1054-1060.
11. Koffi AC, Kouakou B, Kadjo H, Elia G, Koffi SP, et al. (2007) Bilan de deux années de surveillance de la grippe à Abidjan. *Med Trop* 67(3): 259-262.
12. Maman I, Badziklou K, Landoh ED, Halatoko AW, Nzussouo TN, et al. (2014) *BMC Public Health* 14: 981.
13. Maïnassara HB, Lagare A, Tempia S, Sidiki A, Issaka B, et al. (2015) Influenza Sentinel Surveillance among Patients with Influenza-Like-Illness and Severe Acute Respiratory Illness within the Framework of the National Reference Laboratory, Niger, 2009-2013. *PLoS One* 10(7): e0133178.
14. Tarnagda Z, Yougbaré I, Ilboudo AK, Kagoné T, Sanou AM, Cissé A, et al. Sentinel surveillance of influenza in Burkina Faso: identification of circulating strains during 2010-2012. *Influenza Other Respir Viruses* 8(5): 524-529.
15. Kebede S, Conteh IN, Steffen CA, Vandemaele K, Wurie I, et al. (2013) Establishing a national influenza sentinel surveillance system in a limited resource setting, experience of Sierra Leone. *Health Res Policy Syst* 11: 22.
16. Douki LSE, Nkoghe D, Drosten C, Ngougou EB, Drexler JF, et al. (2014) Viral etiology and seasonality of influenza-like illness in Gabon, March 2010 to June 2011. *BMC Infectious Diseases* 14: 373.
17. Nakoune E, Tricou V, Manirakiza A, Komoyo F, Selekon B, et al. (2013) First introduction of pandemic influenza A/H1N1 and detection of respiratory viruses in pediatric patients in Central African Republic. *Virol J* 10: 49.
18. Cosby MT, Pimentel G, Nevin RL, Ahmed SF, Klena JD, et al. (2013) Outbreak of H3N2 Influenza at a US Military Base in Djibouti during the H1N1 Pandemic of 2009. *PLoS One* 8(12): e82089.
19. Ntshoe GM, McAnerney JM, Tempia S, Blumberg L, Moyes J, et al. (2014) Influenza Epidemiology and Vaccine Effectiveness among Patients with Influenza-Like Illness, Viral Watch Sentinel Sites, South Africa, 2005-2009. *PLoS One* 9(4): e94681.
20. Venter M, Lassaunière R, Kresfelder TL, Westerberg Y, Visser A (2011) Contribution of Common and Recently Described Respiratory Viruses to Annual Hospitalizations in Children in South Africa. *J Med Virol* 83(8): 1458-1468.
21. Smuts H (2008) Human coronavirus NL63 infections in infants hospitalized with acute respiratory tract infections in South Africa. *Influenza Other Respir Viruses* 2(4): 135-138.
22. Ijpmma F, Beekhuis D, Cotton MF, Pieper CH, Kimpen JL, et al. (2004) Human metapneumovirus infection in hospital referred South African children. *J Med Virol* 73(3): 486-493.
23. Hussey GD, Apolles P, Arendse Z, Yeates J, Robertson A, et al. (2000) Respiratory syncytial virus infection in children hospitalized with acute lower respiratory tract infection. *S Afr Med J* 90(5): 509-512.
24. Besselaar TG, Botha L, McAnerney JM, Schoub BD. Antigenic and Molecular Analysis of Influenza A(H3N2) Virus Strains Isolated From a Localised Influenza Outbreak in South Africa in 2003. *J Med Virol* 73(1): 71-78.
25. Ahmad JA, Katz MA, Auko E, Njenga KM, Weinberg M, et al. (2012) Epidemiology of respiratory viral infections in two long-term refugee camps in Kenya, 2007-2010. *BMC Infect Dis* 12: 7.

26. Berkley JA, Munywoki P, Ngama M, Kazungu S, Abwao J, et al. (2010) Viral etiology of severe pneumonia among Kenya infants and children. *JAMA* 303(20): 2051-2057.
27. Feikin DR, Ope M, Aura B, Fuller JA, Gikunju S, et al. (2012) The population-based burden of influenza-associated hospitalization in rural western Kenya, 2007-2009. *Bull World Health Organ* 90(4): 256-263A.
28. Johnson AW, Osinusi K, Aderele WI, Gbadero DA, Olaleye OD, et al. (2008) Etiologic agents and outcome determinants of community-acquired pneumonia in urban children: a hospital-based study. *J Nat Med Assoc* 100(4): 370-385.
29. Kwofie TB, Anane YA, Nkrumah B, Annan A, Nguah SB, et al. (2012) Respiratory viruses in children hospitalized for acute lower respiratory tract infection in Ghana. *Virol J* 9: 78.
30. Lutwama JJ, Bakamutumaho B, Kayiwa JT, Chiiza R, Namagambo B, et al. (2016) Clinic- and Hospital-Based Sentinel Influenza Surveillance, Uganda 2007-2010. *The Journal of Infectious Diseases* 206(S1): S87-93.
31. Madhi SA, Schoub B, Simmank K, Blackburn N, Klugman KP (2000) Increased burden of respiratory viral associated severe lower respiratory tract infections in children infected with human immunodeficiency virus type-1. *J Pediatr* 137(1): 78-84.
32. Cohen C, Walaza S, Moyes J, Groome M, Tempia S, et al. (2015) Epidemiology of Severe Acute Respiratory Illness (SARI) among Adults and Children Aged  $\geq 5$  Years in a High HIV-Prevalence Setting, 2009-2012. *PLoS One* 10(2): e0117716.
33. Ope MO, Katz MA, Aura B, Gikunju S, Njenga KM, et al. (2011) Risk factors for hospitalized seasonal influenza in rural western Kenya. *PLoS One* 6(5): e20111.
34. Cohen C, Simonsen L, Sample J, Kang JW, Miller M, et al. (2012) Influenza-related mortality among adults aged 25-54 years with AIDS in South Africa and the United States of America. *Clin Infect Dis* 55(7): 996-1003.
35. Mueller JE, Yaro S, Madec Y, Somda PK, Idohou RS, et al. (2008) Association of respiratory tract infection symptoms and air humidity with meningococcal carriage in Burkina Faso. *Trop Med Internat Health* 13(12): 1543-1552.
36. Thompson MG, Breiman RF, Hamel MJ, Desai M, Emukule G, et al. (2012) Influenza and malaria coinfection among young children in Western Kenya, 2009-2011. *J Infect Dis* 206(11): 1674-1684.
37. Cohen C, Simonsen L, Kang JW, Miller M, McAnerney J, et al. (2010) Elevated influenza-related excess mortality in South African elderly individuals, 1998-2005. *Clin Infect Dis* 51(12): 1362-1369.
38. Karstaedt AS, Hopley M, Wong M, Brown CH, Tisseau TA (2009) Influenza- and respiratory syncytial virus-associated adult mortality in Soweto. *SAMJ* 99(10): 750-754.
39. Soares JL, Ratsitorahina M, Andrianarivelo RM, Robinson R, Rousset D, et al. (2003) Epidémies d'infections respiratoires aiguës à Madagascar en 2002: de l'alerte à la confirmation. *Arch Inst Pasteur de Madagascar* 69(1&2): 12-19.
40. WHO (2002) Influenza outbreak-Madagascar. *MMWR* 45.
41. WHO (2003) Influenza outbreak in the district of Bosobolo, Democratic Republic of the Congo, November-December 2002. *Wkly Epidemiol Rec* 78: 94-96.
42. Cohen C, Moyes J, Tempia S, Groome M, Walaza S, et al. (2015) Mortality amongst Patients with Influenza-Associated Severe Acute Respiratory Illness, South Africa, 2009-2013. *PLoS One* 10(3): e0118884.
43. McMorrow ML, Wemakoy EO, Tshilobo JK, Emukule GO, Mott JA, et al. (2015) Severe Acute Respiratory Illness Deaths in Sub-Saharan Africa and the Role of Influenza: A Case Series From 8 Countries. *J Infect Dis* 212(6): 853-860.
44. Walaza S, Cohen C, Nanoo A, Cohen AL, McAnerney J, et al. (2015) Excess Mortality Associated with Influenza among Tuberculosis Deaths in South Africa, 1999-2009. *PLoS One* 10(6): e0129173.
45. Tempia S, Walaza S, Cohen AL, Mollendorf C von, Moyes J, et al. (2015) Mortality Associated With Seasonal and Pandemic Influenza Among Pregnant and Nonpregnant Women of Childbearing Age in a High-HIV-Prevalence Setting-South Africa, 1999-2009. *Clin Infect Dis* 61(7): 1063-1070.
46. Neuzil KM, ZHU Y, Griffin MR, Edwards KM, Thompson JM, et al. (2002) Burden of inter-pandemic influenza in children younger than 5 years: a 25 year prospective study. *J Infect Dis* 185(2): 147-152.
47. Brooks WA, Goswami D, Rahman M, Nahar K, Fry AM, et al. (2010) Influenza is major contributor to childhood pneumonia in a tropical developing country. *Pediatr Infect Dis J* 29(3): 216-221.
48. Halloran ME, Longini IM (2006) public Health: Community studies for vaccinating schoolchildren against influenza. *Science*. 311(5761): 615-616.
49. Tam NVJS (2006) Influenza related hospital admissions in children: evidence about the burden keeps growing but the route to policy change remains uncertain. *Arch Dis Child* 91(1): 5-7.
50. Sears SD, Clements ML, Betts RF, Maassab HF, Murphy BR, et al. (1988) Comparison of live attenuated H1N1 and H3N2 cold-adapted and avian-human influenza A reassortant viruses and inactivated virus vaccine in adults. *J Infect Dis* 158(6): 1209-1219.
51. Gessner BD, Shindo N, Briand S (2011) Seasonal influenza epidemiology in sub-Saharan Africa: a systematic review. *Lancet Infect Dis* 11(3): 223-235.
52. Madhi SA, Maskew M, Koen A, Kuwanda L, Besselaar TG, et al. (2011) TIV vaccine in African adults infected with HIV: double blind, randomized CT of efficacy, immunogenicity, and safety. *Clin Infect Dis* 52(1): 128-137.
53. Monto AS (2004) Global burden of influenza: what we know and what we need to know. *International Congress Series* 1263: 3-11.
54. Sanicas M, Forleo E, Pozzi G, Diop D (2014) A review of the surveillance systems of influenza in selected countries in the tropical region. *Pan Afr Med J* 19: 121.