Osteoporosis risk – think twice about bone density in Ethiopians African Americans

Abstract

Objective: The risk of an osteoporotic fracture is well described and readily available in developed nations where 1 in 3 women and 1 in 5 men will experience an osteoporotic fracture. The risk of an osteoporotic fracture in developing nations is less known. Considerations of skeletal and non-skeletal factors may not fully describe risk of osteoporotic fractures. The purpose of this manuscript is to highlight the risk found in Ethiopians and Ethiopian immigrants.

Discussion: In the United States, bone mass density is greatest in black women and least in Asians. Osteoporosis incidence from greatest to least in United States is as follows: Native Americans, Asians, Hispanics, Caucasians, and blacks. Data is not available in many developing nations. For example, for Ethiopians data is not available on fracture risk. Thus, Ethiopians and Ethiopian immigrants when seeking medical attention may be considered black. Yet, Ethiopians may have a high prevalence of osteoporosis.

Conclusion: Risk factors for osteoporosis must be considered when examining and treating patients. Screening for less known risk factors such as low calcium intake, prolonged lactation, smoking, and excessive alcohol consumption must be considered for primary, secondary, and tertiary prevention.

Keywords: Ethiopia, osteoporosis, bone density

Volume 4 Issue 3 - 2019

Barbara Billek-Sawhney

Department of Physical Therapy, Butler Health System, Slippery Rock University, USA

Correspondence: Barbara Billek-Sawhney, Slippery Rock University, Physical Therapy Program, Slippery Rock, PA, 16057, USA, Email barbara.billek@sru.edu

Received: April 13, 2019 | **Published:** May 17, 2019

Introduction

The population of the world in 2017 was 7.6 billion. By 2030, it is expected to be 8.6 billion.¹ Between 2017 and 2050, 50% of the world's population is expected to be in 9 countries, see Table 1.¹ With the decrease in the world's fertility rate coupled with a simultaneous increase in the aging population, there is a silver tsunami or graying of the world.¹

Table 1 Half of the world's population is expected to be in these countries between 2017 and 2050¹

India
Nigeria
Democratic Republic of the Congo
Pakistan
Ethiopia
United Republic of Tanzania
United States of America (USA)
Uganda
Indonesia

Listed in order of the expected contribution to the growth.

The percent of the population 60 years of age and older is expected to double between 2017 and 2050. By 2100, it should triple.¹ By 2050, it is expected that 35% of the European population will be 60 years and older. Conversely, at the other end of the spectrum for the older adult population growth is Africa. In Africa, the percentage of the

population over the age of 60 is expected to increase from the current 5%, to 9% by 2050, and 20% by 2100.¹

One in 4 older adults fall annually.² The consequences of falls are both physical and physiological. Physically, the consequences of falls include fractures, bruises, and other (lacerations, dislocations, sprains, pains, and hematoma). Physiologically the effects of falls, reported from 87 publications, include, death/morbidity, functional decline, inactivity, functional dependency and loss of autonomy, depression, loss of self-confidence, and a loss of self-efficacy.³ In US adults, age 50 and above, approximately 1 in 2 women and up to 1 in 4 men, will sustain an osteoporosis fracture⁴ In developing nations, the incidence of osteoporosis, frequency of falls, and their impact is not well known. Globally, the risk of having an osteoporotic fracture is 1 in 3 women and 1 in 5 men.^{4,5} Risk increases with advancing age.^{4,5} The lifetime risk of a fragility fracture is 50% in Caucasian women and 13% in Caucasian men. The overall mortality of a hip fracture is 15-30% with the majority occurring within 6-months of a fracture.⁶ Therefore, with the graying of the world, the incidence of fracture and associated mortality will increase.

Since the development of the FRAX®, the Fracture Risk Assessment Tool in 2011, more than 24.8 million individuals have calculated their individualized fracture risk. The FRAX® is a well-known web and paper-based tool to calculate the 10-year fracture risk based on individual risk factors. Models are available for 63 countries.⁷ Unfortunately, the tool is unable to consider some ethnic minorities because epidemiological information is not available. The FRAX calculation tool can be applied for Asia, Europe, Middle East and Africa, North America, Latin America, and Oceania. Within the Middle East and Africa regions, data is available for only the countries of Abu Dhabi, Iran, Jordan, Kuwait, Lebanon, Morocco, Palestine, and Tunisia.⁷

When, a person's country is not listed, the user is advised to use the country most like the user's country.⁷ unfortunately for the country of Ethiopia, it would be difficult to decide which country Ethiopians should select. Should Ethiopians consider selecting Tunisia or US black or another country? If a health care provider cares for an Ethiopian immigrant, would the health care provider be aware of the patient's potential risk for osteoporosis? Would the health care provider considered the Ethiopian patient to be black and believe the patient to have the greatest bone density? Within Ethiopia the diversity is pronounced and variable with variability in languages, norms, culture, skin color, and physis. This brief paper will stress the need to look at all risk factors associated with osteoporosis when working with individuals from Ethiopia.

Clinical considerations

Ethiopia has a reported net migration rate of 0.2 migrants/1,000 population and a population of more than 108 million as of 2018.⁸ The number of immigrants from Sub-Saharan Africa to United States (U.S.) has grown. In 1980 there were 130,000 immigrants; in 2015 this number escalated to 1.7 million.⁹ Migration to other countries has also occurred. The number of immigrants in 2015 for Ethiopia was 229,000 or 13.3% of the total Sub-Saharan (1,716,000).⁹

Although in 2004, the Surgeon General published a report on bone health and osteoporosis, addressing concerns related to ethnic minorities, racial minorities, and underserved populations, a subsequent report has not been published.¹⁰ Research by Cauley and colleagues on fracture risk in U.S. reveals fractures occur most in Caucasians and Native Americans and least in Asians, Africans, and Hispanics. A study of almost 20,000 U.S. women investigated 5 ethnic groups. Bone mass density was greatest in black women and least in Asians. Osteoporosis incidence from greatest to least was: Native Americans (11.9%), Asians (10%), Hispanics (9.8%), Caucasians (7.2%) and blacks (4.2%).¹¹ Irrespective of ethnicity, the risk of osteoporotic fractures increases as the number of risk factors increase.¹¹

Although bone mass density is greatest among African Americans in U.S.,^{11–13} health care providers should employ caution and not assume that an individual with dark pigmentation and being African American predicts lower fracture risk. In Israel the prevalence of osteoporosis and risk factors among Ethiopian immigrants in comparison to Israeli-born women relays a different story. Peled and colleagues found a high prevalence of osteoporosis (38.7%) among Ethiopian immigrants compared to an incidence of 5.2% among Israeli-born women. A strong association between osteoporosis and calcium intake during adolescent years, low body mass index, the prolonged duration of lactation among women during childbearing years, physical inactivity, and oral contraceptives was reported.¹⁴

Kahsay and colleagues researched osteoporosis in the Tigray region of Ethiopia. Participants included 130 osteoporotic individuals with a mean age of 60.9±10.1 and 266 controls with a mean age of 46.9±8.7 years. Rural residents were found to be 1.93 times more likely to develop osteoporosis than participants who resided in urban areas (adjusted odds ratio {AOR} 1.93 (95% CI, 1.11.3.36). Decreased physical inactivity had the strongest association with AOR 3.53 (95% CI, 1.98 and 6.30. Key findings revealed relationships with milk consumption and smoking. Those residing in rural areas and smoking had a greater risk of developing osteoporosis. While individuals who had milk intake >4 times/week and performing work, which was physically demanding had a lower risk of osteoporosis.¹⁵

Balk and colleagues have investigated the dietary intake of calcium, globally. The highest incidence is in Iceland with 1233 mg/day and the lowest is in Nepal at 175 mg/day. Great variability is noted in the amount with Asia having the lowest incidence, followed by Africa and South America, and the greatest calcium intake in North America and Europe.¹⁶

Risk factors for the development of osteoporosis can be found in Table 2. These are classified into modifiable and non-modifiable risk factors. Major well-known resources related to osteoporosis were used to compile the Table 2; these include: the World Health Organization, International Osteoporosis Foundation, National Osteoporosis Foundation, and FRAX®, the Fracture Risk Assessment Tool. Key risk factors identified by 3 of 4 of these resources include: age over 50 years of age, female gender, post-menopausal status in women, family history of osteoporosis or a hip fracture or fracture, low body mass index, low bone mass density, low dietary calcium intake, excessive alcohol consumption, cigarette smoking, and a sedentary lifestyle.^{4,5,7,17} The prolonged duration of lactation found in women during childbearing years described by Peled and colleagues in not listed in Table 2.¹⁴

Table 2 Risk factors associated with osteoporosis^{4,5,7,17}

Uncontrollable risk factors	Controllable risk factors
Being over the age of 50 (F,I,N)	Low dietary calcium intake (I,N,W)
Female gender (F,I,N)	
Menopause (N) (premature (I,W))	Vitamin D deficiency (I,N)
Family history of osteoporosis (N) {specific to hip fracture (F,I)} {fracture in mother (W)}	Excessive alcohol consumption (F,I,N,W) – 3 or more units/day (F)
Low body mass index / being small and thin (F,I,N,W)	
Previous fracture (F,N)	Cigarette smoking (F,I,N,W)
Height loss (N)	
Asian or Caucasian race (I)	Dietary fruit and vegetable deficiency (N)
Estrogen use (W)	Excess protein (N)
Low bone mass density (F,I,W)	Excess sodium (N)
High bone turnover (I)	Excess caffeine (N,W)
Neuromuscular disorders (I)	
Glucocorticoid long-term therapy (F,I)	Sedentary lifestyle (I,N,W)
Rheumatoid arthritis (F,I)	Losing weight (N)
Primary or secondary amenorrhea (I)	Visual impairment (I)
Secondary osteoporosis (F)	
Primary and secondary hypogonadism in men (I)	Frequent falls (I) / trauma (W)
Non-thiazide diuretic use (W)	Strength quadriceps and grip (W)
Diabetes mellitus non-insulin dependent (W)	
Gastric surgery (W)	

Key FRAX®, the fracture risk assessment tool; (f), international osteoporosis foundation; (i), national osteoporosis foundation; (n), world health organization (W)

When comparing the key risk factors identified by major organization to the research specific to Ethiopians the following risk factors were found decreased physical inactivity, milk consumption <4 times/week or low calcium intake during adolescent years, smoking, and low body mass index. The prolonged duration of lactation and use of oral contraceptives reported in the articles highlighted were not identified by key organizations.

Recommendations

Health care providers share a responsibility in primary, secondary, and tertiary prevention of osteoporosis, see Table 3. One should use

Table 3 Prevention defined and application to osteoporosis^{18,19}

Prevention	Definition	Application
Primary	Identifying risk factors and intervening before health problems occur, to reduce risk in individuals and populations	Screen all patients for fracture risk
		Screen all patients for fall risk
		Educate all patients on osteoporosis
		Make appropriate recommendations resistance, weight bearing, postural, balance, aerobic exercise, and patient education
Secondary	Identifying diseases in the early stages to prevent or slow the progression of functional decline and in individuals with an identified condition	Screen all patients for fracture risk
		Screen all patients for fall risk
		Educate all patients on osteoporosis
		Make appropriate recommendations
Tertiary	Reducing the degree of disability by slowing or stopping disease progression	Incorporate resistance, weight bearing, postural, balance, aerobic exercise, and patient education
		Screen all patients for fall risk
		Educate all patients on osteoporosis
		Make appropriate recommendations
		Acutely focus on mobilization, functional mobility, and prevention of secondary problems
		Post-acutely and when appropriate incorporate resistance, weight bearing, postural, balance, aerobic exercise, and patient education

Acknowledgements

None

Conflict of interest

The authors declare no conflicts of interest.

References

1. <https://www.un.org/development/desa/publications/world-population-prospects-the-2017-revision.html>
2. <https://www.cdc.gov/features/falls-older-adults/index.html>
3. Terroso M, Rosa N, Marques T, et al. Physical consequences of falls in the elderly: a literature review from 1995 to 2010. *Eur Rev Aging and Phys Act.* 2013;11(1):51–59.
4. <https://www.nof.org/patients/what-is-osteoporosis/>
5. <https://www.iofbonehealth.org/osteoporosis-musculoskeletal-disorders/osteoporosis/diagnosis/fracture-risk-assessment>
6. Masi L. Epidemiology of osteoporosis. *clinical cases in mineral and bone metabolism.* 2008;5(1):11–13.
7. <https://www.sheffield.ac.uk/FRAX/index.aspx>
8. <https://www.migrationpolicy.org/country-resource/ethiopia>
9. <https://www.migrationpolicy.org/article/sub-saharan-african-immigrants-united-states>
10. United States public health service, office of the surgeon general. bone health and osteoporosis: a report of the surgeon general. U.S. Dept. of health and human services public health service office of the surgeon general, USA; 2004.
11. Cauley JA, Wu L, Wampler NS, et al. Clinical risk factors for fractures in multi-ethnic women. the women's health initiative. *J Bone Miner Res.* 2007;22:1816–1826.

12. https://americanbonehealth.org/wp-content/uploads/2016/12/BONESENSEonFractureRiskinAfricanAmericans_FINAL.pdf
13. Barrett CE, Siris ES, Wehren LE, et al. Osteoporosis and fracture risk in women of different ethnic groups. *J Bone Miner Res.* 2005;20:185–194.
14. Peled R, Dahan D, Endevelt R, et al. Osteoporosis among Ethiopian immigrant women: A risk analysis. *Arch Osteoporos.* 2007;2:45–52.
15. Kahsay AB, Gashe FE, Debessai ZG, et al. Risk factors of osteoporosis among adults in Ethiopia, the case of Tigray region. A case control study. *J Public Health Epidemiology.* 2014;62(2):92–1000.
16. Balk EM, Adam GP, Langberg VN, et al. Global dietary calcium intake among adults: A systematic review. *Osteoporos Int.* 2017;28(12):3315–3324.
17. http://apps.who.int/iris/bitstream/handle/10665/42841/WHO_TRS_921.pdf?sequence=1&isAllowed=y
18. <http://guidetoptpractice.apta.org/>
19. https://www.cdc.gov/pictureofamerica/pdfs/picture_of_america_prevention.pdf