

Research Article


 Open Access


 CrossMark

Assessment of the prevalence and associated risk factors of post dural puncture headache (PDPH) after cesarean section delivery under spinal anesthesia

Abstract

Background: Post-Dural Puncture Headache (PDPH) is a common problem after a deliberate puncture of the dura-arachnoid for the purposes of diagnosis, therapy, spinal anesthesia, or unintentionally during epidural procedures. It is a clinically main complication which affects the daily life of patients with marked restriction of their physical activities. Spinal anesthesia is the frequent anesthetic procedure for obstetric patients which identified as cause for PDPH. The aim of the study was to assess the prevalence and associated risk factors of PDPH after Cesarean Section (CS) delivery under spinal anesthesia.

Method: An institution based cross sectional study design was conducted on all eligible obstetric patients who came for operation under spinal anesthesia from September, 2015 to January, 2016. The data collection method was including chart review and patient follow up for three days of post operative period.

Results: 107/251 (42.6%) patients developed PDPH. Among those patients with PDPH big needle sizes (AOR=8.6; 95% CI: 0.06-0.46) and repeated number of attempts (AOR=4.54; 95% CI: 0.52-39.14), were found to be significantly associated with the dependent variable of PDPH on the multi variate logistic regression.

Conclusion and recommendation: In this study, we showed the prevalence of PDPH was higher, 107/251 (42.6%) compared with other literatures. The study also showed that big spinal needles and repeated number of attempts were the independent associated risk factors for PDPH in Felege Hiwot Referral Hospital, Bahir Dar, Ethiopia. The higher magnitude of PDPH has to be reduced by avoiding use of big needles, and the repeated dura puncture.

Keywords: post dural puncture headache, spinal anesthesia, cesarean section, spinal needle

Volume 8 Issue 6 - 2017

 Fentahun Tarekegn,¹ Setegn Eshetie,² Adugna Aregawi,³ Kassaw Moges⁴
¹Department of Anesthesia, College of Medicine and Health Sciences, Bahir Dar University, Ethiopia

²Department of Medical Microbiology, College of Medicine and Health Sciences, University of Gondar, Ethiopia

³Department of Anesthesia, College of Medicine and Health Science, Addis Ababa University, Ethiopia

⁴Department of Anesthesia, College of Medicine and Health Science, Bahir Dar University, Ethiopia

Correspondence: Fentahun Tarekegn, Department of Anesthesia, College of Medicine and Health Sciences, Bahir Dar University, Ethiopia, Tel +251913824072, Email tarekegnfentahun@gmail.com

Received: July 07, 2017 | Published: October 24, 2017

Abbreviations: PDPH, post dural puncture headache; SA, spinal anesthesia; CS, cesarean section

Introduction

Regional anesthesia is the favored methods of cesarean section delivery due to their safety to the mother, simplicity of the technique, lesser maternal risk and satisfactory postoperative analgesic effect.¹⁻³ However, Post-dural puncture headache (PDPH) has been a problem for patients next to dural puncture which is the iatrogenic cause of patient's morbidity in modern anesthesia, as well as pain management therapy after attempted epidural and spinal blocks.² It is believed to be caused through penetration of dura matter by spinal needle with continuous cerebrospinal fluid (CSF) outflow. The concern of the new born and bond of family member may be affected by the post-operative headache.⁴

PDPH is explained as a bilateral headache which is associated with position. It is better during recumbence and worsened during upright position of the patient. The factors that can affect the incidence of PDPH includes age, gender, pregnancy, history of PDPH, shape of needle tip, size of needle, number of lumbar puncture and needle orientation to dural matter.^{4,5} The association between needle size and type with incidence of PDPH was described as 75% for 16-18G needles, 30% for 22G Quincke needles and reduced to 0.37% for 27G pencil point needles.⁶ The most favourable needle sizes for spinal

anesthesia are probably the 25G, 26G, and 27G needles.^{7,8} When we used smaller size of spinal needle, there will be decreased risk of PDPH, due to low CSF leakage through narrowed puncture of the dura.⁸ Based on the diagnostic criteria of the International Headache Society (IHS) in 2004, the Post dural puncture headache can appear up to the fifth day after the procedure and it is self limiting in a week which is defined by at least one of the following symptoms: neck stiffness, tinnitus, hypoacusia (partial loss of hearing), photophobia, and nausea are manifested.⁹

In Ethiopia Bahir Dar, Felege Hiwot Referral Hospital anaesthetists are doing spinal anesthesia by spinal needles with different sizes, but the same design for Obstetric patients. We did this cross sectional study for the purpose of knowing the magnitude of PDPH based on this difference of needle sizes. Therefore, we assessed the prevalence and associated risk factors of PDPH after cesarean section delivery under spinal anesthesia and finally, to disseminate the results of the practice to other anesthesia professionals and it will be base line information for other researchers.

Methods

Study design and patients

A Cross Sectional study design was conducted at Felege Hiwot Referral Hospital, North West Ethiopia from the time of duration

September, 2015 to January, 2016. All consecutive cesarean section patients at postoperative period were included by fulfilling the inclusion criteria of ASA status I - II patients after Cesarean Section was done upon spinal anesthesia. There were cases rejected as exclusion criteria of Uncooperative patients, Patients with impaired cognitive ability and Patients with eclampsia.

Study variables

Dependent variable is post dural puncture headache.

Independent variables are age, body mass index (BMI), and American society of Anesthesiologist (ASA), needle size, needle design, position, and number of attempts and previous history of PDPH.

The sample size was taken as total 251 patients in the time duration of September, 2015 to January, 2016.

Data collection

The entire procedures were performed at sitting position with different Anaesthetists who have greater than two years of experience. The backside of the patients was cleaned with Iodine and alcohol. Spinal anaesthesia was done using a midline approach at the L2-3 or L3-4 interspaces by using different size of spinal needles and 0.5 % isobaric bupivacaine 2.5-3.0ml was injected. The intra operative information could be collected by one of the data collector from each patient chart. Patients were interviewed by another data collector on day 1, 2, 3 and were questioned as regard to headache, location, character, and duration, associated symptoms like neck stiffness, tinnitus, hypoacusia (partial loss of hearing), photophobia, and nausea. PDPH was diagnosed as fulfilling the following criteria. These are headache develops within 3 days after dural puncture, headache that worsens within 15 minutes after sitting or standing and improves within 15 minutes after lying down, and with at least one of the following symptoms : neck stiffness, tinnitus, hypoacusia, photophobia and nausea were included.

Data analysis

Data were analyzed in SPSS version 20 by using bi-variant and multi-variant logistic regression. Odds ratio with 95% confidence interval and p-value were computed to determine the strength of the association. A p-value <0.05 was considered as statistical significant.

Ethical approval

Ethical clearance was obtained from Amhara Regional Health Bureau Research Ethics Review Committee (RERC), Bahir Dar. Written informed consent was obtained from the patient after clear explanation what they had to do throughout the study. Anyone who was not willing to participate in the study could resign at any time. Confidentiality was guaranteed with anonymous questionnaires and keeping them locked.

Results

Socio-demographic & physical characteristics of the study participants

The 251 Patients were included in this study with fulfilling the criteria. However, three patients were excluded due to refusal to take the sample. The mean age of patients participated in study was 27.24 years old with a standard deviation of 5.23 years old and 18 years old is the minimum age of patients participated in this study, where as 40

years old is the maximum age. All patients were either ASA I or ASA II (Figure 1) (Table 1).




Figure 1 Proportion of patients who developed PDPH participants underwent spinal anesthesia cesarean section delivery, in the period of September 25, 2015–January 10, 2016.

Table 1 Socio-demographic and physical characteristics of the study participants who underwent spinal anesthesia cesarean section, in the period of September 25, 2015 – January 10, 2016

Variable	Frequency: n (%)
Age in years	
18 – 30	196 (78.1%)
31 - 45	55(21.9%)
BMI	
< 18.5 (underweight)	8(3.2%)
18.5 – 24.9 (normal)	222 (88.4%)
>24.9 (over weight)	21 (8.4%)
ASA status	
ASA I	223(88.8%)
ASAII	28(11.2%)

Spinal anesthesia related parameters of the study subjects

Sixteen patients had a previous history of spinal anesthesia exposure and two of them complained a PDPH like headache after the procedure. All patients had given spinal anesthesia on sitting position. 21 G needle is the most frequently used spinal needle which is 45.4% of total patients whereas 20 G is used as 2.4%. There were 3 cases (1.2%) diagnosed as failed block which were converted to general anesthesia. None of patients developed PDPH (Table 2).

Table 2 Spinal anesthesia related parameters of the study participants who underwent cesarean section delivery in the period of September 25, 2015–January 10, 2016

Variables	Frequency: n (%)
Previous spinal anesthesia	
Yes	16(6.4%)
No	235(93.6%)
Previous history of PDPH	
Yes	6(2.4%)
No	245(97.6%)
Position of spinal anesthesia done	
Sitting	251(100%)
Lateral	0(0%)
Number of attempts	

Table Continued...

Variables	Frequency: n (%)
Single attempts	199(79.3%)
Twice attempts	42(16.7%)
>2 attempts	10(4%)
Size of spinal needle	
20 Gauge	6(2.4%)
21 Gauge	114(45.4%)
22 Gauge	77(30.7%)
23 Gauge	15(6%)
24 Gauge	21(8.4%)
25 Gauge	18(7.2%)
A successful block	
Yes	248(98.8%)
No	3(1.2%)
Associated symptoms	
Neck stiffness	92(36.7%)
Tinnitus	6(2.4%)
Hyperacusia	1(0.4%)
Photophobia	5(2%)
Nausea	46(18.3%)
None	101(40.2%)

Factors associated with PDPH

Hosmer-Lemeshow test of goodness of fit was performed to check the appropriateness of the model for analysis. Variables found to be significant at a binary logistic regression were: needle size and number of attempts. After analysis with multivariate logistic regression needle size and number of attempts were found to be significant at $p\text{-value}<0.05$ (Table 3). Size of the needle used to administer spinal anesthesia is significantly associated with the development of PDPH. Patients received spinal anesthesia using bigger spinal needles were more than eight times more likely to develop PDPH than patients who received spinal anesthesia using smaller needles. Another significant association was found between number of attempts and PDPH. Patients who received spinal anesthesia (SA) with multiple attempts were four times likely to develop PDPH than their counterpart patients who had a single attempt.

Table 3 Factors associated with PDPH of patients who underwent spinal anesthesia cesarean section delivery in the period of September 25, 2015 – January 10, 2016

Variables	PDPH		AOR(95% CI)	P – value
	Yes	No		
Spinal needles	Big needles (20 G, 21 G & 22 G)	102	101	8.6 (0.06-0.46)
	Small needles(23 G, 24 G, 25 G)	5	43	1 0
Attempt	multiple	37	70	4.54 (0.52-39.14)
	Single	15	129	1 0.015

Prevalence of PDPH

In this study PDPH was present in 107 patients (42.6%).

Discussion

Post dural puncture headache (PDPH) has been believed to be a major problem of patients after spinal anesthesia. The overall postdural puncture headache in this study was 42.6% which is comparable to Egypt study,¹⁰ but excessively higher than other studies report.^{9,11,12}

The high percentage of prevalence of PDPH in this study might be related with the most 77.8 % of participants were received spinal anesthesia using big spinal needle. Specifically, the contribution of big needle was strongly significant association for the over all of PDPH as compared with small needles. This higher PDPH percentage after spinal anesthesia by using big needles were 8.6 times more likely to develop PDPH than small needles (AOR= 8.6; 95% CI: 0.06, 0.46; $p = 0.000$). This might be linked with larger needles put down wider opening on the dura which allowed more CSF pour out than smaller hole caused by smaller needles. Our finding is in line with different studies.¹³⁻¹⁶ However, we couldn't see the associations to the outcome variable on type of design of needle, because of all were Quincke type.

The other significant association was found linking the number of attempts and the development of PDPH. The spinal anesthesia was successful at first attempts with 79.3% which is less likely to develop PDPH than those patients who have repeated attempts. In addition, patients who had an attempt of more than once are about 4.5 times at risk to develop PDPH than those patients who had a single attempt (AOR=4.54; 95% CI: 0.52, 39.14; $p=0.015$). This could be correlated with the number of attempt to increase the probability of piercing the dura matter repeatedly will increase the volume of CSF leak, thus increasing the probability of development of intracranial hypotension & PDPH. This finding is aligned with other studies.^{5,15} The proportion of repeated attempts of spinal needles related PDPH reports from a population based study in University of Basel, Switzerland (4.2 %)¹⁷ was somehow lower than our report (14.7%). However, some other studies couldn't come across significant association between the number of attempts and PDPH.^{16,18,19}

Even though different studies showed on variables of the lower BMI, younger age, and previous history of PDPH are listed as risk factors for PDPH development,^{5,14,20} our observation study did not bring into being significant association between these variables and PDPH. This might be due to the lack of sample size to compare lower to higher BMI, young to old age, and patients with versus without previous history of PDPH. There are some limitations in our study. The sample size was not sufficient for different size of needles (23G, 24 G, 25 G, and 26 G). For this reason, it was not possible to determine the proportion of PDPH in each of small needles. Our study participants were followed for three days only. However, they may develop PDPH until the seventh days of dural puncture which may underrate the overall prevalence of PDPH. We didn't also observe the severity of post dural puncture headache.

Conclusion

In conclusion, the prevalence of PDPH was higher, 107/251 (42.6 %) compared with most other studies. The study also showed that big spinal needles and repeated number of attempts were the independent associated risk factors for PDPH in Felege Hiwot Referral Hospital, Bahir Dar, Ethiopia. We recommend the higher magnitude of PDPH has to be reduced by avoiding use of big needles and the repeated dura puncture.

Availability of data and materials

No additional data are required; all information is clearly presented in the main manuscript.

Funding

No funding source.

Consent for publication

Not applicable.

Ethical approval

It was obtained from Amhara Regional Health Bureau Research Ethics Review Committee (RERC) with official permission letter to conduct the research and written informed consent was taken.

Authors' contribution

Fentahun Tarekegn conceived the study and developed the proposal, collected the data, analyzed the data and manuscript preparation. Setegn Eshetie, revised the proposal and involved in data collection, data analysis and manuscript preparation. Aregawi Aregawi, revised the proposal and involved in data collection, data analysis and manuscript preparation. Kassaw Moges also revised the proposal and involved in data collection and manuscript preparation. All authors approved the final manuscript and agreed to publication in *Journal of Anesthesia and Critical Care*.

Acknowledgments

We would like to thank Felege Hiwot Referral Hospital anesthetists for the achievable of this study.

Competing interest

We declare that have no competing interest.

References

1. Haider S, Sikander R. The Pencilpoint Lumbar Puncture Needles: Are they Worth their Price? *Ann Pak Inst Med Sci*. 1990;5(4):216–219.
2. Camann WR, Murray RS, Mushlin PS, et al. Effects of Oral Caffeine on Postdural Puncture Headache A Double-Blind, Placebo-Controlled Trial. *Anesth Analg*. 1990;70(2):181–184.
3. Bano F, Sabbar S, Zafar S, et al. Intrathecal fentanyl as adjunct to hyperbaric bupivacaine in spinal anesthesia for caesarean section. *J Coll Physicians Surg Pak*. 2006;16(2):87–90.
4. Rasooli Sousan, Moslemi Farnaz, Baybordi Ali. Post-dural Puncture Headache in the Obstetric Patient: Needle Size, Number of Dural Puncture and Timing of Ambulation. *International Journal of Women's Health and Reproduction*. 2015;3(3):163–167.
5. Lybecker H, Møller JT, May O, et al. Incidence and Prediction of Postdural Puncture Headache A Prospective Study of 1021 Spinal Anesthetics. *Anesth Analg*. 1990;70(4):389–394.
6. Gogarten W, Van Aken H. A century of regional analgesia in obstetrics. *Anesth Analg*. 2000;91(4):773–775.
7. Kang SB, Goodnough DE, Lee YK, et al. Comparison of 26-and 27-G needles for spinal anesthesia for ambulatory surgery patients. *Anesthesiology*. 1992;76(5):734–738.
8. Shaikh JM, Memon A, Memon MA, et al. Post dural puncture headache after spinal anaesthesia for caesarean section: A comparison of 25 G Quincke, 27 G Quincke and 27 G Whitacre spinal needles. *J Ayub Med Coll Abbottabad*. 2008;20(3):10–13.
9. Amorim JA, Gomes de Barros MV, Valença MM. Post-dural (post-lumbar) puncture headache: risk factors and clinical features. *Cephalgia*. 2012;32(12):916–923.
10. Hassan Mohamed Ali, Mohamed Yehya Mohamedb, Yahya Mohamed Ahmedb. Postdural puncture headache after spinal anesthesia in cesarean section: Experience in six months in 2736 patients in Kasr El aini teaching hospital – Cairo University. *Egyptian Journal of Anaesthesia*. 2014;30(4):383–386.
11. Douglas MJ, Ward ME, Campbell DC, et al. Factors involved in the incidence of post-dural puncture headache with the 25 gauge Whitacre needle for obstetric anesthesia. *Int J Obstet Anesth*. 1997;6(4):220–223.
12. Richman JM, Joe EM, Cohen SR, et al. Bevel direction and postdural puncture headache: a meta-analysis. *Neurologist*. 2006;12(4):224–228.
13. Frank RL. Lumbar puncture and post-dural puncture headaches: implications for the emergency physician. *J Emerg Med*. 2008;35(2):149–157.
14. Amorim JA, Valença MM. Postdural puncture headache is a risk factor for new postdural puncture headache. *Cephalgia*. 2008;28(1):5–8.
15. Kuczkowski KM. Post-dural puncture headache in the obstetric patient: an old problem. New solutions. *Minerva Anestesiol*. 2004;70(12):823–830.
16. Bezo D, Ashina S, Lipton R. Post-dural puncture headache: Part II—prevention, management, and prognosis. *Headache*. 2010;50(9):1482–1498.
17. Seeger MD, Kaufmann M, Staender S, et al. Repeated Dural Punctures Increase the Incidence of Postdural Puncture Headache. *Anesth Analg*. 1996;82(2):302–305.
18. Jeanjean P, Montpellier D, Carnec J, et al. Headaches after spinal anesthesia: prospective multicenter study of a young adult population. *Ann Fr Anesth Reanim*. 1996;16(4):350–353.
19. Imarengiaye C, Ekwere I. Postdural puncture headache: a cross-sectional study of incidence and severity in a new obstetric anaesthesia unit. *Afr J Med Med Sci*. 2006;35(1):47–51.
20. de Almeida SM, Shumaker SD, LeBlanc SK, et al. Incidence of Post-Dural Puncture Headache in Research Volunteers. *Headache*. 2011;51(10):1503–1510.