

**Table 4** Foraminiferal species from sediment collected in spring (October)

Species Spring (October)/ number of individuals	1	2	3	4	5	6	7	8	10	11	12	13	14	15	16	17	18	19	20	absolute frequency	relative frequency	
<i>Ammonia rolshauseni</i>	1														1	12	9			23	1.06	
<i>Ammonia tepida</i>	54	35	30	52	68	51	20	52	30	31	40	33	20	40	24	13	39	41	18	691	31.95	
<i>Buliminella elegantissima</i>	12	42	65	43	14	41	41	37	61	51	49	44	32	18	31	28	10	25	54	698	32.27	
<i>Uvigerina peregrina</i>																	1			1	0.05	
<i>Cassidulina subglobosa</i>														1		1				2	0.09	
<i>Hanzawaia boueana</i>																				0	0.00	
<i>Haynesina germanica</i>	1											1					1			3	0.14	
<i>Fursenkoina pontoni</i>									1											1	0.05	
<i>Bulimina patagonica</i>		2	2					2							1	1			1	9	0.42	
<i>Elphidium sp.</i>	22	1		3	2	4	22	6	4	9	8	18	45	58	47	37	30	11	1	328	15.16	
<i>Pseud. atlanticum</i>					5		5	1	1	4	1	5	10	4	3				34	3	76	3.51
<i>Buccella peruviana</i>	1				1															2	0.09	
<i>Bulimina sp.</i>		1																		1	0.05	
<i>Bulimina marginata</i>			2			1		1	1	1		1		1		1			1	10	0.46	
<i>Bulimina pulchella</i>					1															1	0.05	
<i>Bolivina striatula</i>	5		12	3	23	4	10	9	5	13		12	8	5	6	9	6	6	16	152	7.03	
<i>Quinqueloculina patagonica</i>	2																			2	0.09	
<i>Trochammina inflata</i>	4	1	38	5	8				1	5	5	1	4	1				3		76	3.51	

<i>Trochammina hadai</i>	1	7	7	6	14	10	1	10	4	2	3								1	1	1	1	69	3.19
<i>Gaudryina exillis</i>	1		1																	1			3	0.14
<i>Paratrochammina clossi</i>																							0	0.00
<i>Fissurina lucida</i>			1			1	2					2	1	2								3	12	0.55
<i>Oolina sp.</i>																						1	1	0.05
<i>Lagena striata</i>													1										1	0.05
<i>Polisaccamina ipohialina</i>									1														1	0.05
<b>Total</b>	10	89	15	11	13	11	10	11	10	11	108	117	121	128	114	104	97	121	98			2163	100.00	
	4		8	2	6	2	1	8	9	6														

Source: The authors, 2019.