

Synthesis, QSAR Model Study and Antimicrobial Evaluation of Esters and Thioester Derivatives of Isonicotinic Acid on the Different Strains of *Mycobacterium Tuberculosis*

Research Article

Volume 2 Issue 6 - 2016

Victor Olugbenga Fadipe^{1*}, Julián Alejandro Yunes Rojas², Eugenio Sánchez Areola³, Jocelyn Juarez Badillo³, Hassan H Abdallah⁴, and Rowland Andrew Opoku⁵

¹Department of Chemistry, University of Zululand, South Africa²Departamento de Ingeniería en Biotecnología, Tecnológico de Monterrey, Mexico³Department of Chemical Biological Sciences, Universidad de las Americas Puebla, Mexico⁴Department of Chemistry, Salahaddin University, Iraq⁵Department of Biochemistry and Microbiology, University of Zululand, South Africa

*Corresponding author: Victor O Fadipe, University of Zululand, Department of Chemistry, Private Bag X1001, KwaDlangezwa, 3886, South Africa, Tel: 035 902 6099; Fax: 035 902 6568; Email: vobfadipe@yahoo.com

Received: April 15, 2016 | Published: July 14, 2016

Abstract

Isoniazid is one of the widely used first-line drugs in the treatment of tuberculosis for many years but the mechanism of the action of the drugs is still not clear. Herein, we report the synthesis, QSAR model study and biological effects of esters and thioester derivative of isonicotinic acid (INA) on different strains of *Mycobacterium tuberculosis* in an attempt to establish the mode of action of the active form of Isoniazid (INH). The esters of INA are expected to show antibiotic activities against strains of *Mycobacterium tuberculosis* upon activation by intracellular nonspecific esterases, while thioesters is expected not to show antimicrobial effects and thus served as control. The obtained results indicated that the synthesized esters did not show antimicrobial activity. The conclusion drawn from the result is that either INA is not the active form of INH or that the esters show poor water solubility- in addition to the possibility of acquired during resistance of the strains during culture and testing - all of which could hamper mycobacterial cell up take. This suggests that more studies are needed to search for the possible active form of isoniazid.

Keywords: Oxidation; *Mycobacterium tuberculosis*; Isoniazid; Rifampicin; Pyrazinamide; Ethambutol; Streptomycin; Thioester; Micro-organism; Isonicotinic; Hydrolyzed; Prodrug

Introduction

Mycobacterium tuberculosis (*Mtb*) was discovered in 1882 by Robert Koch [1]. Tuberculosis is caused by this bacillus and it is widely spread in the world. It has been reported to be a common opportunistic infection of HIV. Isoniazid (INH) is used for the treatment of tuberculosis like other first line drugs (isoniazid, rifampicin, pyrazinamide, ethambutol or streptomycin). Isoniazid is a pro-drug that is activated by KatG enzyme, and its products inhibit the production of nicotinic acids generated in the InhA complex. KatG enzymes are limited to *Mtb* [2].

Due to the high resistance of *Mtb* in the hospitals caused by inappropriate treatment administration, it has become necessary to search for isoniazid's derivatives that are probable intermediate responsible for its biological activity. This probable intermediate which is synthesized derivatives is therefore expected to be potent enough to kill this micro-organism (*Mtb*). Ester and thioester are the active forms of isonicotinic acid (INA) that can be generated under different condition (Figure 1) [3].

The mechanism of action of isoniazid against *Mtb* has been studied from diverse approaches [4]. The *in silico* analysis suggests that the mechanism of action can either be accomplished by free radical generation [5], or possibly accomplished by redox reaction mechanism [5]. The latter assumption seems to agree

with the experimental results. Consequently, our study focuses on the mechanism of action of INH and INA. It is possible that the isonicotinic acid and its derivatives, esters and thioesters, may also be active. When these derivatives; esters or thioesters are hydrolyzed, either of the compounds can easily liberate its acid free form.

The hypothesis of the mechanism of action of Isoniazid by free radical had been reported earlier (see Figure 2 in [5]). The INH is activated by an intracellular enzyme of *Mtb*, KatG, which has a hemo group with an iron forming the complex I. The iron has +4 oxidation state (Fe⁴⁺), but the ground state of iron shows an oxidation state +3, which precedes the iron state seen on the complex (I). It interacts with hydrogen peroxide and converts it to a molecule of water and the compound I. Compound I is then bonded to an oxo group and an iron atom to make it stable (see also Figure 2 in [5]).

The generated INH is in the radical state, nitrogen at one end, which is used to form the complex II which has +3 oxidation states in iron atom. Only Fe³⁺ and water are created after this complex has gained an electron. With that, the generated cycle of one radical in the INH is closed. Since a second radical is the INH radical, it is then transformed by a radical attack on the previous nitrogen, and a second radical, that is named Nicotinyl radical, is then liberated [6-9].

The hypothesis of the mechanism of action of isoniazid by an ionic pathway also creates an intracellular isonicotinic acid which occurs only inside of the mycobacterial cell (see also Figure 2 in [5]).

The figure 2 in [5] shows an *Mtb*, since the prodrug of INH enters into the *mycobacterium*, and KatG activates the alleged prodrug, the product of the activation is the isonicotinic acid. It will be in a deprotonated form and at that stage; the pH determines its behavior. If it enters the *Mtb* cell plasma with its neutral pH and pKa value of 4.8, then its intracellular concentration will increase because the neutral compound becomes a negatively charged acyl anion. Hence, it cannot get out by passive diffusion; it is trapped inside the *Mtb* cells. Consequently, if there is the possibility to generate a substitution of nicotinic acid and to form the iso-NAD

complex, which inhibits InhA activity, letting the cell without the production of mycolic acids [4,10-14].

The findings from the literature indicated that the active form of the INH could be the a fore mentioned INA [5,15,16]. According to this hypothesis, esters and thioesters of the INA are expected to have biological activity against *Mtb* strains when they are activated by non-specific intracellular esterases. In support of this supposition, we carried out the following combined approach to gain insight into the intriguing mechanism of action of INH: (i) created a computational model, (ii) used activity data to generate QSAR models, (iii) carried out the synthesis of esters and thioesters of INA, (iv) before the determination of the antimicrobial effects of these compounds in *Mtb* strains.




Figure 1: Structural drawings. (A) Isoniazid (INH) is a first line drug in the treatment against tuberculosis. (B) Isonicotinic acid (INA) could be the possible active form of INH as a postulated prodrug generated by KatG. Both molecules were drawn using Chem Sketch [3].




Figure 2: Chemical variety of basic molecules and the metabolites which can occur either by enzymatic activities or "nonenzymatic" hydrolysis: a) INA, b) INH, c) Hydrazones, and d) Hydrazines [15].




Figure 3: The QSAR studies allowed us to interpret that INH derivatives are also precursors and their efficiency depends on conversion in INH or INA. None of the esters and thioesters of INA inhibited the growth of mycobacteria [30].

Experimental

Synthesis of ester and thioester of isonicotinic acid

The solvents used in the experiments showed technical purity (Fermont and Baker). The reagents had analytical purity (Sigma-Aldrich). The measurements of NMR of ^1H and ^{13}C were

carried out on a Varian Gemini spectrometer at 200MHz for ^1H and 50.3 MHz for ^{13}C . The solutions with deuterated chloroform contained tetramethylsilane (TMS) as an internal reference; while the infrared data was collected on a Varian-00 FT-IR (Scimitar Series) spectrometer; in addition to the ultraviolet data which was obtained from a Varian Cary-100 ultraviolet spectrometer.

General procedure for the preparation of esters of isonicotinic acid

A solution of isonicotinic acid (0.010 mol), alcohol (1-butanol, 1-hexanol, 1-octanol) (0.015 mol) and H_2SO_4 underwent reflux for 4 hours with magnetic stirring [17]. The reaction was monitored and verified to completion by TLC; it was cooled and neutralized with a saturated solution of $NaHCO_3$. After then, the product of the reaction was extracted with EtOAc (3x50 mL), the organic phase was washed with water and dried with Na_2SO_4 anhydride; the organic phase was filtered and evaporated to dryness in rotary evaporator. The product was purified again using column chromatography, using hexane initially and then increasing its polarity with chloroform as mobile phase.

General procedure for the preparation of thioesters of isonicotinic acid [8-10]

A solution of isonicotinic acid (0.010 mol), thionyl chloride (0.012 mol) and chloroform (10 mL) was heated at 70°C for 9 hrs, and the components were mixed by magnetic stirring under N_2 atmosphere, and a solution of thiol (0.015 mol) in chloroform (5mL) was added [18]. The reaction was monitored and verified to completion by TLC. The mixture was cooled and neutralized with a saturated solution of $NaHCO_3$, and was finally extracted with ethyl acetate (3 times 50 ml EtOAc). The organic phase was washed with water, followed with a solution of 10% NaOH and was dried with anhydrous Na_2SO_4 . In the next step it was filtered; subsequently, it was evaporated to dryness on the rotavapor. The product of the reaction was purified on column chromatography with silica gel, using hexane initially and then with chloroform in order to increase the polarity of the mobile phase.

Infrared spectroscopy

The infrared spectrometer is expected to provide information about two important bands: the O-C=O stretching of an ester and the S-C=O stretching of a thioester. The typical infrared signal of a C-O group of an ester is fairly different from the C-S signal of a thioester because the mass and electro-negativity of the oxygen and sulfur are quite different. Sulphur atoms are heavier in mass than oxygen atoms. Hence, an increase in mass lowers the observed frequencies. A more electronegative atom like oxygen destabilizes the resonance structure of the ester, giving the C=O a higher bond order; that is a stronger double bond character. As a general result, the ester signals are absorbed at a higher frequency. Because the sulphur is far less electronegative than oxygen, the former has a more stabilized resonance structure due to its greater ability to donate its electrons, which in general then results in lower frequencies.

The stretching vibration due to vibrational intensity of C=O bond shows greater frequencies between 1750 and 1735 cm^{-1} (less wavelength) than ketones frequencies.

The IR signals of the C-O stretching are basic 2 coupled asymmetric vibrations: C-C (=O)-O. The observed range is 1300-100 cm^{-1} while the ester saturated bands strongly appeared in a region around 1210 to 1163 cm^{-1} . The corresponding frequencies for vinylic esters and phenolic moieties were observed near 1190

to 1140 cm^{-1} . The O-C-C signals of esters coming from primary alcohols were observed around 1164 to 1031 cm^{-1} . Ester derivatives of secondary alcohols had a distinct signal at 1100 cm^{-1} . Aromatic ester derivatives of primary alcohols showed a distinct signal at 1111 cm^{-1} [19, 20].

NMR spectroscopy

The signals of 1H - NMR for alkanes oscillate generally between 0.8 and 4.0 δ , while the aromatic compounds showed displacements superior to 7.0 δ [21,22]. The ^{13}C -NMR provided the C-C scaffold structure of each compound.

Anti-biograms

The experimental data were generated from the modified proportions of dilutions for a slow growing *mycobacterium*. Using this method, the susceptible assay can be made directly and indirectly. Usually, the pure cultures of organism were made in the indirect form where it was grown in solid media, 7H10 or 7H11 Middle brook agar or Löwenstein-Jensen media [23].

Agar preparation

Water bath was heated to a temperature of 50-56°C, then into an Erlenmeyer flask was added 200mL of distilled water and the vegetal mass weighed 1.9g per 100 mL of water. Then, the water was added slowly with a magnetic stirrer to dissolve the mixture uniformly. Heating was done until the agar turned clear (approximately 20 minutes). One mL of glycerin was added and stirring continued for another 3 to 5 minutes. The flask was then removed from the heating source and covered with aluminum in autoclave for 15 minutes at 121°C.

Preparation of antimicrobial solutions

The isoniazid was used as reference substance for testing. Thus, it was used in various concentrations in order to prepare antimicrobial solutions for the experiments. Finally, 1mL of the solution was prepared, and added to the various concentration of 0.2 μ L up to 1.0 μ L of the testcompounds in ethanol.

The prepared concentrations were tested against the reference compound ethambutol. It was prepared by using 1mL of the solution along with 5 to 10 μ L preparations for each ester and thioester compounds. The last prepared solutions were used under different concentrations from 10 to 20 μ L [24].

Preparation of petri plates

- Add one OADC supplement flask (media composed of oleic acid, albumin, dextrose and catalase) per 200 ml of agar.
- Label 3 divisions on the petri plates with one plate per compound. One division will be the test control, another division will be for the higher concentration and the last one will be for the lower concentration.
- Add 100 μ L from the antimicrobial solution of high concentration and 100 μ L from the antimicrobial solution of lower concentration to the corresponding plate divisions.
- Add 5 mL of agar in each division of the plates.

Inoculation and incubation

- Take a sample from the Mtb culture with a swab.
- Suspend it in 1mL of alkaline solution and then allowing it to reach the turbidity state.
- Inoculate with a swab by simple streaking on every one of the plate divisions. Initiate tests with the control group, then followed by the lower concentration and finally, finished with the higher concentration.
- Incubate in moisture presence at 36°C for a period of 7 days and observe it till the end of the given time. If growth does not take place in the control group, then the incubation is repeated for another 7 days.

Preparation of the Ziehl-Neelsen stain

Procedure to realize this stain:

- Smear preparation.
 - Take one drop of saline solution and place it over a glass slide.
 - Take a microorganism colony and place it over the saline solution.
- Place filter paper over the smear.
- Add phenol-carbol fuchsin, let it remain for 5 minutes but heat it during this time. Do not over heat the Fuchsin preparation.
- Wash it off with water. In this step, the filter paper has to be eliminated.
- Add alcohol-acid and allow for 3 min.
- Wash it off with water.
- Add to it methylene blue and allow it mixing for 1 min.
- Wash it off with water.
- Examine under a microscope with a 100-fold magnifying objective. Mycobacteria are colored red, and the other organisms that are not from this genus are colored blue.

QSAR models

The QSAR studies were generated with the computational program Molecular Operating Environment (MOE) for 200 derivatives of INH. They were obtained from database bank at NIH [25-27]. The QSAR modeling technique and its problems are described in the literature [28].

Results and Discussion

QSAR study analysis

INH has been considered as a lead compound in the literatures [25-27]. Each of the intermediate compounds from INH has an MIC value \geq INH [15]. In the analysis of the hydrazine derivatives of INH, the first mathematical equation is related to eight

parameters as reported in Equation 1. On critical analysis, it was however discovered that isoniazid derivatives have problems with the incubation conditions and decay of derivatives would therefore have a consequence that would alter the microbiological potency measurements of the experiment [15]. In this regard, our experiment would provide a useful insight.

The QSAR study's result was an equation that related biological activity to molecules' physicochemical properties. Using multiple linear regressions, a numerical relation (see equation below) between chemical structures with physicochemical properties and its biological activity was obtained. The final equation showed a linear correlation coefficient R^2 of 0.75. ($R^2=1$, reflects a straight line, and represents a high linear correlation; $R^2<0.5$ means there is a poor linear correlation).

With the values, $n = 60$; $R^2 = 0.75$ the following was obtained:

$$\text{pMIC} = 5.41532 - (0.00416 * \text{Hydrophobic VdW surface area}) - (2.65893 * \text{AM1 imino-carbonyl charges}) + (0.16648 * \text{Kier-topological flexibility index}) - (0.93573 * \text{Molecular shape}) - (0.00751 * \text{Molecular volumen}) + (0.23376 * \text{AM1 HOMO}).$$

The model is barely logic in the chemical-microbiological interpretation. The analyzed parameters are contradictory in some cases. It should consider other possible explanations to help understand why the reported values of the 60 compounds from the QSAR studies do not correlate with the interpretation of its parameters calculated and such explanations could be as follows:

- Bacterial resistance by impure cultures.
- Environmental microbiological contamination.
- Prior disintegration of molecules in vitro extreme conditions: high temperature, humidity, slow growth, chemical nutrients.
- MIC values not reliable (measurement errors).
- In the Table 1 are shown the indexes of the 'X' descriptors on the three QSAR studies.

$$\text{Equation 1: pMIC} = -0.93551 + (0.33301 * X1) - (0.89074 * X2) - (0.41049 * X3) - (0.48897 * X4) + (0.33167 * X5) + (0.27508 * X6) + (0.11044 * X7) + (0.15883 * X8), \text{ with } n=60; R^2 = 0.596.$$

In the equation 1, two contradictions were observed for the polar - non polar area contributions. If the nonpolar character is increased in the molecule, the activity concentration (pMIC) is also expected to increase while the polar character should decrease. The nonpolarity can be quantified with X1: Log P and X2: Hydrophobic VdW surface area, with X3: AM1 imino-carbonyl charges (partial atomic charges by AM1 of "C" carbonyl and "O" atom), thus reflects polarity. The X8: Molecular acidity, which correlates with Hammett sigma or a cationic charge (positivation) atomic partial charge. From the results obtained, it is observed that the nonpolar character shows a positive sign in X1 which is in contradiction to the negative sign in x2. Similarly, the positive sign of X8 which indicates deprotonization or dissociation into ions is a contradiction to the positive sign in X4.

Table 1: Descriptors X used in the three QSAR models.

QSAR Models				
	Descriptors Used	1 st	2 nd	3 rd
X1	Octanol distribution coefficient/water Log P	*		
X2	Hydrophobic VdW Surface area	*	*	*
X3	AM1 Imino-Carbonyl Charges	*	*	*
X4	Sum of bond orders	*		
X5	Maximun electrotopological positive Variation	*		
X6	Kier topological flexibility index	*	*	*
X7	Zagreb topological index M2	*		
X8	Molecular Acidity (pKa)	*		
X9	Molecular form		*	*
X10	Molecular volume		*	*
X11	AM1 HOMO		*	*
	Number of descriptores	8	6	6

Concerning the first contradiction

The coefficient of molecular descriptor X1: Log P was positive; showed that the biological activity increases, along with increase in nonpolarity. Nonetheless, (b) the b2 coefficient of molecular descriptor X2 is negative, the biological activity also increased in order to decrease the nonpolarity. Whereas an equation cannot exist with such contradictions; if a nonpolarity character increases, then all the characters with the same type of parameters should show increase.

The second contradiction was the following

The charges tend to become a value of zero for increasing the biological activity, this means that the charges should lose positive or negative charges, as it is shown with the descriptor X3. (b) If the acidity should increase, then consequently, the biological activity should increase. This is also a sort of "positivation" as shown on the descriptor X8. It is not possible to have two parameters that imply polarity, and yet both act inversely.

A second study used the following mathematical equation using six parameters which are reported in Equation 2.

Equation 2: $pMIC = -0.09070 - (0.28233 * X2) - (0.17931 * X3) - (0.13868 * X6) - (0.45384 * X9) - (0.27005 * X10) + (0.06250 * X11)$, with $n = 60$; $R^2 = 0.784$.

In this model two contradictions were observed.

First contradiction

(a) The coefficient of X2: Hydrophobic surface area VdW

decreased as long as the biological activity increased and the polarity decreased. (b) The b3 coefficient of X3: AM1 imino-carbonyl charges decreased while the biological activity increased and the polarity decreased. It is shown that the parameters are moving in opposite direction because of their polarity index.

Second contradiction

(a) The coefficient of X3 decreased as the biological activity increased and the polarity decreased.

(b) The coefficient of X11:AM1 HOMO increased while the biological activity increased and the polarity was also increased. The parameter types are similar regarding polarity. It is expected to increase or decrease in the same direction. For an example, parameters with good correlation are X2 versus X11 where the non-polarity diminishes, besides the polarity increments.

A third mathematical model was created by Equation 3.

Equation 3: $pMIC = 5.41532 - (0.00416 * X2) - (2.65893 * X3) + (0.16648 * X6) - (0.93573 * X9) - (0.00751 * X10) + (0.23376 * X11)$, with $n = 60$; $R^2 = 0.754$.

Equation 3 of QSAR model number 3 study obtained two contradictions again in the polarity - non polarity area. In this equation, it manifested a first discrepancy: two contrary parameters cannot work in the same direction, as it is the case with hydrophobicity (X1) and polarity (X3). With regard to the second inconsistency, the charges with the quantum chemistry AM1 (X3) method correlates positively with the value of pMIC; nevertheless, the charges of imino-carbonyl (X2) correlates negatively. The two parameters under investigation with the same type of character have to act in the same direction. In this case, the biological activity constitutes the potential of antibiotic effect. The attraction between "C" carbonyl charges and "O" atoms of the parameter X2 greatly determines its activity for the trend of the metabolic breakdown of INH. Concerning the decrease in molecular volume, parameter X5, a maximum electro-topological positive variation could reflect a better reception through passive diffusion.

The linear relation in the molecules led to the particular analysis of the molecules and the way the values of reported pMIC were related. Studying the most usual procedures of *Mtb* incubation, it was found that it consists of an aqueous acidic medium as liquid phase, with glutamic acid as catalyst, high temperature, exposing during days [15]. This allows degradation of hydrazones in INH and other products. Because MIC is reported, it is not necessarily the report about the activity of its initial molecule. In most of the cases it was degraded otherwise. Only one INH concentration was reported that is not identical to the initial concentration of the microbiological experiment and the other nonregistered products [15]. However, in some other cases, it was already interpreted that the toxic molecules of hydrazine had a stronger antibacterial activity than INH [15]. The reasons were as follows: there are substances in its composition that act while INH is liberated by the hydrolysis of these molecules. These molecules are capable of acting slowly in the *mycobacterium*. PABA was identified a bacterial nutrient like in these cases. The other reason was that when the molecule is hydrolyzed, and finally INH is obtained from

the expected active concentrations of the initial molecule [15]. Hence, its MIC is only apparent and the true active concentration becomes that of INH because the initial molecules (prodrugs) are converted into INH (drug). However, INH acts as the principal molecule at the end. Other possible reason for the variations in the reported values of MIC was that it is possible during the point of determination that some strains with high bacterial resistance were introduced by impure cultures or microbial ambient contamination.

The QSAR study of this study allowed us to observe a series of errors that affected the calculus of the models and the equations of the correlations. It has been reported [28] that:

- The members of the series do not share the same molecular mechanism of action, because it should have different receptor-target although its structures are related. Do not include them in the QSAR study.
- The compounds are unstable or precursors and the structures of the active forms of these compounds are modified. The resultant structures from the reaction of instability are unknown, but occasionally are not detected. The QSAR study was sparsely utile because it was understood that the biological values are referred to other structures different from the original compounds which had been synthesized and then may have been modified by a decay process because they are not stable enough in the experiments.
- The series showed only little structural variability. The chemical space has not been exploited in a wide range by the chemists. Only a part of it was evaluated, which is called "bias by a reduced strategy of synthesis". Only those compounds which are very easily and readily synthesized exist and more complicated organic pathway was not explored for new synthesis [28].
- The series has too much variability, it is difficult to establish a rule because of the lack of similar examples of a reference compound; since the distributions of the values are not equidistant and do not exist "black holes".
- The parameters do not describe the causal form of the mechanism of molecular action against the receptor. This is caused by a bad choice of descriptors or a limited offer in the QSAR software or by an inappropriate hypothesis of work.
- There are too many descriptors, the relation between the number of descriptors and the number of compounds in the series as minimum is 1:6 or 1 to 10 etc. However, R² value diminished when the numbers of descriptors are reduced. The author had to insert many descriptors in his model to raise R² in order for the model to gain acceptability. This is called "over fitting".
- R² is interpreted like a quality indicator of the model. Descriptors explained the cause of the relation between structure and activity. Hence, accordingly R² is not a measure of causal axis. If it is drawn "X" and "Y" axes on a diagram, "X" axis is descriptor and "Y" axis is the activity.
- To find descriptors with significant probability, there must be

statistically significant differences between the descriptors in the model. Although they may not be pharmacologically significant. Consequently, if there is a statistical probability, it is not known whether the parameters are also relevant on the nature by fact. As an example: it is statistically known that there are diminishing number of stork nests and the decreasing number of baby birth in the same strip of land. Traditionally, as a joke, it has been assumed that storks bring the babies. However, this conclusion is actually not ridiculous at all. There is a common basis (a hidden covariance in statistics) for both populations to decline, the link and explanation for both trends is the destruction of natural habitat, both populations showing symptoms of stress in a highly industrialized zone, where green areas are destroyed and the humans keep busy and cannot relax [28]. Hence, before reporting any correlation, there must be knowledge about the observations, a deeper understanding of the parameters as well as their relationship to the observed data, at best a causal relation exists. Moreover, in QSAR models, linearity is a prerequisite to use MLR techniques but nonlinearity before cannot be discarded for natural processes what is in need of other ways doing statistics [28].

Generating a fourth QSAR model with 57 derivatives of hydrazones.

A QSAR study with 57 hydrazone derivatives of INH was carried out (Figure 2). The molecules were all manually selected from NIAID (National Institute of Allergy and Infectious Diseases) at http://apps1.niaid.nih.gov/struct_search/

The flow analysis of a QSAR type study was as follows: In the first place MIC values were collected to represent the biological activities of the molecules and the physicochemical parameters for the derivatives were calculated. Then multiple linear regression (MLR) was applied to correlate the molecular properties (independent variables) with the biological end points (dependent variable, activities). Then the QSAR model equations were generated [28].

The parameters were calculated for the QSAR study and the following final equation of MLR was obtained as follows:

$$\text{MIC} = -0.45 * (\text{Shape}) - 0.28 * (\text{PEOE_VSA_HYD}) - 0.27 * (\text{Vol}) - 0.18 * (\text{q_R_AM1-Mul}) + 0.14 * (\text{KierFlex}) + 0.06 * (\text{AM1_HOMO})$$

The resulting coefficient of correlation was in the upper range of values (0.7 to 0.9) which represents a reliable QSAR model, namely R²=0.8. This means that this value can be accepted, because it is close to 1, the ideal point.

Analyses of the antibiograms

The esters were prepared by esterification of the isonicotinic acid with alcohol (1-butanol, 1-hexanol and 1-octanol) that was catalyzed by H₂SO₄. The confirmation of the ester formation was made by NMR of 1H where it showed the presence of the characterized signals of the isonicotinic acid's aromatic ring and the corresponding signals of the alcohol residue. The structure was confirmed with the NMR data of the ¹³C and IR and with the literature data [29].

Thioesters were prepared by transforming isonicotinic acid with isonicotyl chloride and by also reacting it with the corresponding thiol. The obtained thioesters were identified with its spectroscopy data (UV, IR and NMR) of ^1H and ^{13}C in similar like esters (Table 2).

The prepared compounds were proved against *Mtb*. The main observations of our experiment reflected that there was no inhibition of the *Mycobacterium tuberculosis* growth. Hence, it was concluded from the obtained result of the synthesized esters that the isonicotinic acid is not the active form of the isoniazid [30].

Table 2: Spectroscopy data of the thioesters from isonicotinic acid.

Thioester	UV (EtOH) λ_{max} nm (log ϵ)	IR (CHCl ₃) cm ⁻¹	NMR- ¹ H δ ppm (J)	NMR- ¹³ C δ ppm
S-butyl pyridine-4-carbothioate	217 (4.02); 273 (3.86)	3085, 2956, 2921, 1686, 1234, 1200, 973.	8.76 (dd, 4.4, 1.6 Hz) (H-2); 7.73 (dd, 4.4, 1.6 Hz), 7.73 (dd, 4.4, 1.6 Hz), 8.76 (dd, 4.4, 1.6 Hz), 3.10 (t, 7.4 Hz), 1.68 (br quint, 7.4 Hz), 1.45 (br sext, 7.4 Hz), 0.95 (t, 7.4 Hz).	149.7, 119.5, 142.2, 119.5, 149.7, 189.2, 29.1, 31.4, 27.2, 13.8.
S-hexyl pyridine-4-carbothioate	216 (4.01); 273 (3.85)	3085, 2957, 2921, 1661, 1200, 1191, 991	8.76 (dd, 4.4, 1.6 Hz), 7.73 (dd, 4.4, 1.6 Hz), 7.73 (dd, 4.4, 1.6 Hz), 8.76 (dd, 4.4, 1.6 Hz), 3.08 (t, 7.4 Hz), 1.67 (br quint, 7.4 Hz), 1.2-1.5 m, 0.89 (br t, 6.5 Hz).	149.3, 119.1, 141.8, 119.1, 149.3, 189.2, 28.6*, 29.0*, 28.3*, 31.0, 22.3. * May be interchanged
S-octyl pyridine-4-carbothioate	216 (4.06); 273 (3.90)	3090, 2939, 2869, 1670, 1234, 1217, 956	8.75 (dd, 4.4, 1.7 Hz), 7.72 (dd, 4.4, 1.7 Hz), 7.72 (dd, 4.4, 1.7 Hz), 8.75 (dd, 4.4, 1.7 Hz), 3.07 (t, 7.0 Hz), 1.67 (br quint, 7.0 Hz), 1.2-1.5 m, 0.88 (br t, 6.6 Hz).	149.4, 119.2, 141.9, 119.2, 149.4, 189.3, 28.8*, 29.0*, 28.9*, 29.1, 29.1, 31.6, 22.5, 14.1.

Conclusion

Isoniazid (INH) is among the antibiotics of first line for the treatment of tuberculosis. INH is a prodrug activated by catalase-peroxidase (KatG) of *Mycobacterium tuberculosis* (*Mtb*). The active form is poorly characterized. There are two possible ways of INH activation. The active form interferes in the synthesis of mycolic acids which are essential to *mycobacterium*. It hinders the enoil-reductase (InhA). Our study was intended to contribute to clarify the mechanism of action of the first line drug INH. Its derivatives were designed and synthesized to be tested. These compounds are ester derivatives of INA that should have antibiotic activity in the *Mtb* strains when they are activated by intracellular unspecific esterases. Furthermore, the corresponding thioesters were used as a negative control group in our study. On the one side, the inherent risk of ester building is the decrease in water solubility. On the other side, the esters are lipophilic and permeation of lipid compartments and barriers (cell walls etc) should be possible. So, a trade-off (not too long, not too short) for the ideal chain length of esters was necessary. Microbiological tests to evaluate the pharmacologic activity were carried out with the compounds. Concerning the widely accepted mechanism, it has been proposed that KatG is responsible for the production of Isonicotinic acid's (INA) radicals that associate with NAD. Secondly, another hypothesis postulated that intracellular accumulation of INA replaces the nicotinic acid in the NAD synthesis in order to produce iso NAD. The analysis of the antibacterial effects of the INA's esters was negative. The interpretation of the negative results is as follows: either INA is

most likely not the active form or the esters could not enter the bacterial cells, due to insufficient balance between water and lipid solubility. The drug resistance phenomenon cannot be ruled out, despite the positive susceptibility (the sensitivity pattern) of the *Mtb* reference strains, but mutations characterized in KatG or InhA could confer detrimental drug resistance, all of which would affect the results.

Acknowledgement

We are much indebted to Thomas (alias Tom) Scior, at the Pharmacy Department of Facultad de Ciencias Químicas, Universidad Autónoma de Puebla, Mexico, for directing the PhD work of JAJR, manuscript reading and language corrections, and assistance in all stages of the editorial procedure.

References

- Koch R (1882) Die Aetiologie der Tuberkulose. Berliner klinische Wochenschrift pp. 428-445.
- Coll P (2003) Fármacos con Actividad frente a *Mycobacterium tuberculosis*. Enfermedades Infecciosas Microbiología Clínica 27(8): 299-308.
- ACD/Chem Sketch, version 12.01 (2010) Advanced Chemistry Development, Inc., Toronto, ON, Canada.
- Seydel JK, Tono-oka S, Schaper KJ, Bock L, Wiencke M (1976) Mode of action of isoniazid (INH). Short communication: Isolation and identification of an analog of NAD from 3H-INH treated bacterial cells. Arzneimittel forschung 26(4): 477-478.

- Scior T, Meneses MI, Garcés EJ, Domeyer D, Laufer S (2002) Antitubercular Isoniazid and Drug Resistance of *Mycobacterium tuberculosis* - A Review. *Arch Pharm Pharm Med Chem* 335(11-12): 511-525.
- Chouchane S, Lippai I, Magliozzo RS (2000) Catalase-peroxidase (*Mycobacterium tuberculosis* KatG) catalysis and isoniazid activation. *Biochemistry* 39(32): 9975-9983.
- Rozwarski DA, Grant GA, Barton DH, Jacobs WR, Sacchettini JC (1998) Modification of the NADH of the isoniazid target (InhA) from *Mycobacterium tuberculosis*. *Science* 279(5347): 98-102.
- Wengenack N, Lopes H, Kennedy M, Tavares P, Pereira A, et al. (2000) Redox potential measurements of the *Mycobacterium tuberculosis* heme protein KatG and the isoniazid-resistant enzyme KatG(S315T): insights into isoniazid activation. *Biochemistry* 39(37): 11508-11513.
- Cowan JA (1997) *Inorganic Biochemistry: An Introduction*. (2nd edn), Wiley-VCH Publications, New York, USA, pp. 456.
- Johnsson K, Schultz P (1994) Mechanistic Studies of the Oxidation of Isoniazid by the Catalase Peroxidase from *Mycobacterium tuberculosis*. *JACS* 116(16): 7425-7426.
- Banerjee A, Dubnau E, Quemard A, Balasubramanian V, Um KS, et al. (1994) InhA, a gene encoding a target for isoniazid and ethionamide in *Mycobacterium tuberculosis*. *Science* 263(5144): 227-230.
- Thiemer EK (1956) *Chemie des Isoniazids*. *J BerBorstel* 3: 192-424.
- Thiemer EK (1957) *Biochemie des Isoniazids*. *J BerBorstel* 4: 299-509.
- Seydel JK, Schaper KJ, Wempe E, Cordes HP (1976) Mode of Action and Quantitative Structure-Activity Correlation of Tuberculostatic Drugs of the Isonicotinic Acid Hydrazide Type. *J Med Chem* 19(4): 483-492.
- Scior T, Eisele GSJ (2006) Isoniazid is not a lead compound for its pyridyl ring derivatives, isonicotinoyl amides, hydrazides, and hydrazones: a critical review. *Curr Med Chem* 13(18): 2205-2219.
- Wahab HA, Choong YS, Ibrahim P, Sadikun A, Scior T (2009) Elucidating isoniazid resistance using molecular modeling. *J Chem Inf Model* 49(1): 97-107.
- Haslman E (1980) Recent developments in methods for the esterification and protection of the carboxyl group. *Tetrahedron* 36(17): 2409-2433.
- Scheithauer S, Mayer R (1979) Thio- and Dithiocarbonylic Acids and their Derivatives, *Topics in Sulfur Chemistry* 4: 1-373.
- Brown RS, Aman A (1997) Intramolecular Catalysis of Thiol Ester Hydrolysis by a Tertiary Amine and a Carboxylate. *JOC* 62(14): 4816-4820.
- Derrick LJC, Soleiman H (2000) Synthesis of Racemic Brevoxime and Related Model Compounds. *J Org Chem* 65(16): 4923-4929.
- Witter DJ, Vederas JC (1996) Putative Diels-Alder-Catalyzed Cyclization during the Biosynthesis of Lovastatin. *J Org Chem* 61(8): 2613-2623.
- Feng CL, Jing WX, Alper H (2009) Palladium-Catalyzed Ring-Opening Thiocarbonylation of Vinylcyclopropanes with Thiols and Carbon Monoxide. *J Org Chem* 74(2): 888-890.
- Wang Q, Yue J, Zhang L, Xu Y, Chen J, et al. (2007) A Newly Identified 191A/C Mutation in the Rv2629 Gene that Was Significantly Associated with Rifampin Resistance in *Mycobacterium tuberculosis*. *J Proteome Res* 6(12): 4564-4571.
- Horváti K, Bacsa B, Szabó N, Dávid S, Mező G, et al. (2012) Enhanced Cellular Uptake of a New, *in Silico* Identified Antitubercular Candidate by Peptide Conjugation. *Bioconjug Chem* 23(5): 900-907.
- National Institutes of Health [NIAID] (2004) *Anti-HIV Therapeutics Database*.
- Scior T, Eisele GSJ, Badillo JJ, Uribe CA, Yunes JA, et al. (2004) Isonicotinic Acid as Active Metabolite of antitubercular Isoniazid: Relationships between Molecular Properties and Activities Comparing anti *mycobacterial* Drugs. Symposium, Istanbul, Turkey Poster A50.
- Scior T (2005) Estudios computacionales para evaluar las actividades antituberculóticas de doscientos nuevos derivados de isoniazida. Ciclo de seminarios departamentales Facultad de química UNAM. DF México Ponencia.
- Scior T, Franco MJL, Do QT, Mayorga MK, Rojas YJA, et al. (2009) How to Recognize and Work around Pitfalls in QSAR Studies: A Critical Review. *Curr Med Chem* 16(32): 4297-4313.
- Miao C, Zong Q, Xiang Z, Wu J (2011) Synthesis and characterization of novel isonicotinate-based ionic liquids. *Advanced Materials Research* 233-235: 1266-1269.
- Rojas YJA (2011) TESIS DOCTORAL: Análisis de estudios del tipo QSAR con derivados de Isoniazida e Isoxyl; comparación de Ácido Nicotínico (AN) e Isonicotínico (INA) y estimación de una posible estructura activa vía radical al comparar Peróxido de Benzoílo, NA e INA, BUAP.