In Vitro Bacteriostatic and Bactericidal Activities of *Senna Alata*, *Ricinus Communis* and *Lannea barteri* Extracts Against Wound and Skin Disease Causing Bacteria

Abstract

Antibacterial activities of the extracts of three medicinal plants, *Senna alata*, *Ricinus communis* and *Lannea barteri* from two different extractants were studied. Bacteria strains such as *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* and *Escherichia coli* are known wound and skin disease causing bacteria. For 100g of each plant material, the percentage yields of the extracts were determined and ranged between 5.20% and 12.40%. Phytochemical screening of the crude extracts of *Senna alata*, *Ricinus communis* and *Lannea barteri* revealed the presence of some bioactive components, including tannins, anthraquinones, saponins, flavonoids and alkaloids. All plant extracts showed inhibition on test organisms and were in the range of 12.0 ± 2.83 mm to 36.0 ± 2.12 mm for all the extracts. The MIC values obtained for the entire test ranged from 3.13mg/ml to 12.50mg/ml and MBC values were found to be 200mg/ml minimum and 400mg/ml maximum. Findings support the traditional use of *Senna alata*, *Ricinus communis* and *Lannea barteri* for the treatment of skin and wound infections and form a strong basis for further exploration of these plants.

Keywords: Bacteriostatic; Bactericidal; *Senna alata*, *Ricinus communis*; *Lannea barteri*; *Staphylococcus aureus*; *Pseudomonas aeruginosa*; In vitro

Volume 3 Issue 1 - 2016

Addai Mensah Donkor, Richard Mosobil, Jennifer Suurbaar

Department of Chemistry and Biochemistry, University for Development Studies, Ghana

Correspondence: Suurbaar Jennifer, Department of Chemistry and Biochemistry, University for development studies, Ghana, Tel +233506362322,

Email jenniferexpress@yahoo.com, suurbaar@gmail.com

Received: July 22, 2016 | **Published:** September 02, 2016

Abbreviations: *S. Alata*, *Senna Alata*; *R. Communis*, *Ricinus Communis*; *L. Bartari*, *Lannea Barteri*; *K. Pneumoniae*, *Klebsiella Pneumoniae*; *E. Coli*, *Escherichia Coli*; *P. Aeruginosa*, *Pseudomonas Aeruginosa*; *S. Aureus*, *Staphylococcus Aureus*; MIC, minimum inhibitory concentration; MBC, minimum bactericidal concentration; STD, standard deviation

Introduction

Plants in several families of the plant kingdom have been used for centuries in medicine. These plants are found worldwide but they are most abundant in tropical countries.¹ These plants are sources of many chemical compounds, the most important of which are alkaloids, terpenoids, steroids phenols, glycosides and tannins. A number of modern drugs have been isolated from plants and the screening of plant extracts for antimicrobial activity has shown that higher plants represent a potential source of novel antibiotics.²

Due to some antibacterial ineffectiveness, there is an increase in the population of patients that visit medical centers in developing countries. The problem of antibiotic resistance has led to the revival of interest in herbal products as sources of novel compounds to suppress the ever increasing problems of re-emergence of diseases. A large group of microbial pathogens are associated with various skin infections. The *streptococci* and *staphylococci* cause wound infections. These microbes cause opportunistic skin infections in immunosuppressed patients.

The *E. coli* are part of physiological intestinal flora, however outside the intestine they may cause wound infection and sepsis.³ *Pseudomonas aeruginosa* is a prevalent burn patient pathogen; this microbe is able to infect different parts of human body.⁴

Senna alata belongs to the family *Caesalpinaceae*. The shrub stand 3-4 meter tall with leaves up to 50-80cm long; the inflorescence looks like a yellow candle. The plant has fungicidal properties for treating ringworm and other infections of the skin.⁵ Abatan,⁶ reported that leaf juice and decoction of *senna alata* are used in the treatment of ringworm and other skin diseases.⁶ *Ricinus communis* is from the Family *Euphorbiaceae*, popularly known as castor plant. The plant is wide spread throughout tropical regions as ornamental plant. The antimicrobial activities of *Ricinus communis* were good against dermatophytic and pathogenic bacterial strains *Staphylococcus aureus*, *Pseudomonas aeruginosa* as well as *Klebsiella pneumoniae*, *Escherichia coli*.⁷ The *Ricinus communis* possess wound healing activity due to the active constituent of castor oil which produces antioxidant activity and inhibits lipid peroxidation. A study conducted by Prasad et al.⁸ reported the potential of Castor oil in wound healing activity by reducing the scar area and also the epithelization time in excision wound model.⁸ *Lannea barteri* belongs to the family *Anacardiaceae* and they are used as medicinal plants in most African countries for curing some infectious diseases.⁹ In this study, we validated the antibacterial activity of the extracts of *Senna alata*, *Ricinus communis* and *Lannea barteri* from aqueous and ethanolic solvent.

Materials and methods

Plant material

Plant materials of *Senna alata* (leaves), *Ricinus communis* (leaves) and *Lannea barteri* (bark) were collected from different areas in Navrongo, Upper East Region, Ghana. The plants were authenticated by Dr. Isaac Sackey in the Department of Applied Biology, University for Development Studies, Ghana.

Table 1 Phytochemical profile of plant extract

Pythochemicals								
Plants/ Solvents	Tannins	Saponins	Poluronoids	Reducing	Sugars	Terpenoids	Flavonoids	Alkaloids
<i>Senna alata</i>								
Aqueous	+	+	+	+	+	+	+	+
Ethanollic	+	+	-	+	+	+	+	+
<i>Lannea barteri</i>								
Aqueous	+	+	+	+	+	+	+	+
Ethanollic	+	+	+	+	+	+	+	+
<i>Ricinis communis</i>								
Aqueous	+	+	+	+	+	+	+	-
Ethanollic	+	+	+	-	-	+	+	-

+: Presence; -: not Detected

Table 2 MIC and MBC of the plant extract *Senna alata* on the bacterial strains

Senna Alata				
Test organisms	Aqueous extract (mg/ml)		Ethanollic extract (mg/ml)	
	MIC	MBC	MIC	MBC
<i>E. coli</i>	3.13	200	6.25	200
<i>S. aureus</i>	12.5	300	12.5	300
<i>P. aeruginosa</i>	6.25	200	6.25	200
<i>K. pneumoniae</i>	6.25	200	12.5	300

Table 3 MIC and MBC of the plant extract, *Lannea barteri*, on the bacterial strains

Lannea Barteri				
Test organisms	Aqueous extract (mg/ml)		Ethanollic extract mg/ml)	
	MIC	MBC	MIC	MBC
<i>E. coli</i>	6.25	300	6.25	UD
<i>S. aureus</i>	12.5	200	25	400
<i>P. aeruginosa</i>	6.25	200	6.25	200
<i>K. pneumoniae</i>	12.5	200	6.25	200

*UD – undetected

Table 4 MIC and MBC of the plant extract *Ricinis communis* on the bacterial strains

Ricinis communis				
Test organisms	Aqueous extract (mg/ml)		Ethanollic extract (mg/ml)	
	MIC	MBC	MIC	MBC
<i>E. coli</i>	6.25	UD	6.25	UD
<i>S. aureus</i>	3.13	300	25	400
<i>P. aeruginosa</i>	3.13	200	6.25	200
<i>K. pneumoniae</i>	12.5	400	6.25	200

8UD – Undetected

Preparation of plant crude extracts

Plants materials from *Senna alata* (leaves), *Ricinus communis* (leaves) and *Lannea barteri* (bark) were air dried at room temperature for 2 weeks. The respective leaves were ground into uniform powdered using a blender and the back was powdered using pestle and mortar.

The ethanol and water extracts were prepared by socking 100g each of the powdered plant materials in 1L of solvent at room temperature for 48h. The extracts were filtered separately through Whatman filter paper No 42. The extracts were concentrated using a rotary evaporator and further warmed in a water bath to obtain semi-solid and solid products.

Phytochemical screening

Phytochemical screening for all plant extracts was done for the presence of tannins, saponins, terpenoids, puluronoids, reducing

sugars, flavanoids, alkaloids and anthraquinones. The presence of alkaloids was detected by following a method described by Fernand et al.¹⁰ The sample was basified with ammonium hydroxide, extracted with chloroform and subsequently warmed in a water bath to remove traces of chloroform. The residue was dispersed in 2N HCl and a few drops of Mayer's reagent were added. Presence of yellowish cloudy precipitate indicated a positive test for alkaloids. Saponins were tested by adding 4ml of water to the sample and shaking vigorously for 10-15mins. Persistence of foamy layer indicated saponins were present.¹⁰ Tannins were also detected by adding 4.0ml of distilled water to 2.0ml of sample. A few drops of 10% ferric chloride solution were then added and a blue-black color was observed indicating the presence of tannins.¹¹ A few drops of acetone were added to a portion of the sample resulting in a precipitate sticking along the walls of the test tube reminiscent of the presence of polyuronoids.¹⁰ The presence of reducing sugars was verified by adding equal volumes of Fehling's

solution A and B to the sample and heating in a water bath for 15mins. Brick-red precipitate indicated the presence of reducing sugars.¹⁰ Flavonoids were detected using Shibata's reaction test as the extracts were evaporated to dryness and the residue was dissolved using 50 % methanol in a test tube. A piece of magnesium was dropped into the mixture, followed by 3 drops of concentrated HCl with subsequent appearance of orange color indicative of the presence of flavonoids.¹⁰

Antibacterial activity of the plant extracts

Preparation of plant extracts concentrations: The crude extracts obtained from the respective solvents were prepared using dimethylsulfoxide (DMSO). Stock solutions, 400 mg/ml concentration was prepared by adding 0.4g of each extract in 1ml of DMSO in sterile containers. Serial dilutions were carried out to obtain concentrations of 200mg/ml, 100mg/ml, 50mg/ml, 25mg/ml, 12.5mg/ml, 6.25mg/ml, 3.125mg/ml and 1.5625mg/ml of extracts concentration.

Test organism: Wound and skin disease causing bacteria were taken into consideration and four of these bacterial were considered. Selected bacterial were *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*. Clinical isolates of the bacterial were obtained from the Microbiology Department of the Tamale Teaching Hospital in the Northern Region of Ghana, in March 2016. Bacterial isolates were stored at temperatures between 2°C and 8°C in nutrient broth until further analysis.

Agar well diffusion assay: The agar well diffusion method was employed, where Molten Mueller Hinton agar (OXOID, Basingstoke, Hampshire, England) at 40°C was poured aseptically into sterile petri dishes (about 6mm thick) and allowed to solidify. Test organisms in normal saline were spread on the solidified agar. Wells, 6mm, were created and filled with 1.0ml of the various concentrations of the plant extracts. DMSO was used as negative control while Amoxicillin 250mg/ml was used as a positive control. Test were carried out in duplicates and plates were allowed to stand for 30mins at room temperature and finally incubated at 37°C for 24h. The antibacterial activity was evaluated by measuring the diameter of zones of inhibition in millimeters (mm).

Inoculum preparation for minimum inhibitory concentration (MIC) and minimum bactericidal concentrations (MBC): Inocula were obtained from an overnight agar culture of the test organism. Inoculum for the MIC and MBC test was prepared by taking at least three to five well isolated colonies of the same morphology from agar plate culture. The top of each colony was touched with a sterile loop and the group was transferred into a tube containing 5ml of normal saline and then vortexed. The broth culture was incubated at 37°C until it achieved the turbidity of 0.5 McFarland standard (1.0×10^6 cfu) for 6 hours.

Preparation of McFarland standard (Turbidity standard MIC and MBC for inoculum): Barium sulphate turbidity standard equivalent to a 0.5 McFarland was used to standardize the inoculum density for susceptibility test.¹² To make the turbidity standard 0.5 McFarland, Barium chloride solution, 200ml, 175% w/v, was added to 99.5ml 1% sulphuric acid solution, mixed well and stored in the dark at room temperature.

Determination of minimum bactericidal concentration (MBC): The tube diffusion method was employed for the determination of MBCs.¹³ Minimum bactericidal concentrations of the extract was prepared by using sterilized transparent bottles for the different organisms cultured. Normal saline solution, 2ml, containing the test

organisms was each dispensed into the sterilized transparent bottles. A 1ml extract of different concentrations was dropped into sterile transparent bottles containing the test organisms. The bottles were corked and incubated aerobically at 37°C for 24h. Sterile nutrient agar plates were inoculated with sample from each of the transparent bottles. The plates were further incubated for 24h at 37°C and subsequently observed for bacterial growth. The lowest concentration that killed 100% of the inoculum bacteria, thus, no growth on plate was recorded as the MBC.

Determination of minimum inhibitory concentration (MIC): Minimum inhibitory concentration (MIC) of the extracts was carried out using the agar well diffusion method. Variable concentrations of the extract were prepared by serial dilution to obtain concentrations of 400mg/ml, 200mg/ml, 100mg/ml, 50 mg/ml, 25mg/ml, 12.5mg/ml, 6.25mg/ml, 3.125mg/ml 1.5625mg/ml. Nutrient broth, 2ml, containing the test organisms was each dispensed into sterilized test tubes. A 1ml extract at varying concentrations was added to the test tubes containing the test organisms. The tubes were corked and incubated aerobically at 37°C for 24 hours. The tubes were observed after incubation to determine the MIC.

Statistical Analysis: Means and standard error of the mean were calculated for the zones of inhibition measured for the two sets of experiments in each case. These means were statistically compared using the one-way ANOVA to determine if they were significantly different at $P < 0.05$.

Results and discussion

The increasing use of medicinal plants by a sizable proportion of people in the industrial world has been traced to the extraction and development of several drugs from these plants as well as from traditionally used herbal remedies. Three medicinal plants were used in this study and their antibacterial activities were evaluated. The percentage yields were determined from 100g of each plant material and were found to be between 5.20 and 12.40%. *Lannea barteri* gave the highest yield; both aqueous and ethanolic extracts gave 12.40 and 12.10% respectively. *Senna alata* gave the second highest yield of 7.40 and 7.20% in aqueous and ethanolic respectively. The yields for aqueous and ethanolic extracts for *Recinnus cominus* were 6 and 5.20% respectively.

Phytochemical screening on *Senna alata*, *Ricinus communis* leaves extracts and *Lannea barteri* bark extract of the three extractants are indicated in Table 1. Tannins, saponins and alkaloids were present in all the plant extracts of their respective extractants. Polyrunoids, reducing sugars, terpenoids and flavonoids were found in all the plant extracts except *Recinis communis* which indicated absence of anthroquinones. These bioactive compounds have potentially significant application against bacteria, including those that cause wounds and skin diseases.¹⁴ Tannins are widely used in traditional medicine in treating wounds and to arrest bleeding.¹⁵ Some of these bioactive compounds which are synthesized as secondary metabolites as the plant grows, also serve to protect the plant against microbial attacks and predation by animals.¹⁶

Antibacterial activities of crude extracts of *Senna alata*, *Recinnus communis* and *Lannea bartari* were evaluated by measuring the diameters of zones of growth inhibition on bacteria. All test organisms used in this current research were susceptible to all the extracts though to varying degrees. The susceptibility of bacteria to plant extracts, on the basis of inhibition zone diameters varied according to strains and species as well as the extractants. All the plant extracts of both solvents

gave inhibitory zones between $17.5 \pm 0.71\text{mm}$ and $36.0 \pm 2.12\text{mm}$ against *P. aeruginosa*. Both the ethanolic and aqueous extracts of *Senna alata* exhibited similar potency against the test organisms (Figure 1). The degree of inhibition of the different concentrations of the extracts on *P. aeruginosa* were higher than that of Amoxicillin, the positive control which gave between $15.0 \pm 0.71\text{ mm}$ and $16.5 \pm 2.12\text{ mm}$ zones of inhibition on the test organism. Regarding *L. barteri*, the aqueous extract gave higher zones and hence higher potency than the ethanolic extracts at all concentrations used (Figure 2). Similar trends were also exhibited by the plant *R. communis*, in which aqueous extracts showed higher potency with higher zones of inhibition than the ethanolic extracts (Figure 3). All the bacteria used in this study were gram-negative bacteria except *S. aureus* which was gram positive. Gram negative bacteria are known to be resistant to the action of most antimicrobial agents including plant-based extracts.¹⁷ Nikaido et al.¹⁸ reported that gram-negative bacteria have an outer phospholipid membrane with the structural lipopolysaccharide components, which makes their cell wall impermeable to antimicrobial agents.¹⁸ Although *P. aeruginosa* is a gram negative bacterium, it was susceptible to the plant extracts in spite of permeability difference. Hence all the plant extracts used in this work would be considered as having broad spectrum inhibitory activity.

Figure 1 Representative antibacterial activity of ethanolic (e) and aqueous (a) crude extracts of *Senna alata* against *E. coli*, *S. aureus*, *P. aeruginosa* and *K. pneumoniae*. The data shown represent the average of three wells treated on the same day. The experiment was repeated 2 times and day-to-day variation was found to be within one fold of the presented data.

Figure 2 Representative antibacterial activity of ethanolic (e) and aqueous (a) crude extracts of *Lanneabarberi* against *E. coli*, *S. aureus*, *P. aeruginosa* and *K. pneumoniae*. The data shown represent the average of three wells treated on the same day. The experiment was repeated 2 times and day-to-day variation was found to be within one fold of the presented data.

Figure 3 Representative antibacterial activity of ethanolic (e) and aqueous (a) crude extracts of *Ricinus communis* against *E. coli*, *S. aureus*, *P. aeruginosa* and *K. pneumoniae*. The data shown represent the average of three wells treated on the same day. The experiment was repeated 2 times and day-to-day variation was found to be within one fold of the presented data.

Antibacterial activities of *R. communis* and *S. alata* extracts against *P. aeruginosa* have been reported.¹⁹ The degree of inhibition of most of the concentrations of the extracts against *S. aureus* was also higher than the inhibition of Amoxicillin against *S. aureus*. This was consistent with other studies,^{5,19,20} which indicated that the bacterium was highly susceptible to the plant extracts. For the other test organisms, although there was inhibition, the amoxicillin exhibited higher activity than both aqueous and ethanolic extracts from all plant extracts applied in the current research. This could be attributed to the fact that unlike conventional antibiotics and other pharmaceutical products which are usually prepared from synthetic materials by means of reproducible manufacturing techniques and procedures, herbal medicinal products are prepared from materials of plant origin, which may be subjected to contamination, degradation or deterioration. The storage of extracts may require special condition of humidity, temperature and/or protection from light. Moreover, the plant extracts might contain little of the active ingredients.

Extracts with low activity against the organisms gave high MIC and MBC values, while the highly active ones gave low MIC and MBC values. The MIC and MBC techniques were used to evaluate the efficacies of antimicrobial agents and in this study, the MIC and MBC values tend to support the results obtained in the antibacterial screening (Tables 2–4). The MIC values obtained for the entire test ranged from 3.13mg/ml to 12.50mg/ml and MBC values from 200mg/ml as lowest and 400mg/ml being highest. MBC was undetected for *E. coli* with *Recinnus communis* extracts; *P. aeruginosa* however indicated low MIC and MBC (Table 3) (Table 4). All the extracts showed inhibitory activities against skin and wound disease-causing pathogens used as demonstrated by similar works.^{5,14,19–23} The efficacy of the plant extracts provide a scientific basis and thus validates their traditional use in the treatment of skin and wound diseases associated with these bacteria.

Conclusion

The antimicrobial results showed that the plants used in this study are highly bactericidal for broad spectrum activities. It also suggested that the plant extracts could be a potent raw material for pharmaceutical applications.

Acknowledgments

We recognize the support by the research team of the Microbiology and Immunology Department, Navrongo Health Research Centre, Navrongo, UER, Ghana.

We also recognize the support Staff of the Faculty of Applied Sciences, University for Development Studies. Navrongo Campus, UER, Ghana.

Conflicts of interest

The authors declare there is no conflict of interests.

Funding

None.

References

1. Calixto JB. Twenty-five years of research on medicinal plants in Latin America: A personal view. *J Ethnopharmacol*. 2005;100(1-2):131-134.
2. Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev*. 1999;12(4):564-582.
3. Madigan MT, Martinko JM, Dunlap PV, et al. Brock Biology of microorganisms. 12th edn. International Microbiology. 2008;11:65-73.
4. Chanda S, Baravalia Y. Novel leads from herbal drugs for infectious skin diseases. *Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology*. 2010;1:451-456.
5. Owoyale JA, Olatunji GA, Oguntoye S. Antifungal and antibacterial activities of an alcoholic extract of *Senna alata* leaves. *Journal of Applied Sciences and Environmental Management*. 2006;9(3):105-107.
6. Abatan M. A note on the anti-inflammatory action of plants of some *Cassia species*. *Fitoterapia*. 1990;61(4):336-338.
7. Jena J, Gupta AK. *Ricinus communis* Linn: a phytopharmacological review. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2012;4(4):25-29.
8. Prasad MR, Shete R. Pharmacological investigation on the wound healing effects of castor oil in rats. *International Journal of Universal Pharmacy and Life sciences*. 2011;1(1):21-28.
9. Mahesh B, Satish S. Antimicrobial activity of some important medicinal plant against plant and human pathogens. *World journal of agricultural sciences*. 2008;4(5):839-843.
10. Fernand WN, Adama H, Jeanne FM, et al. Phytochemical composition, antioxidant and xanthine oxidase inhibitory activities of *Amaranthus cruentus* L. and *Amaranthus hybridus* L. Extracts. *OALIB Journal*. 1982. 67 p.
11. Sofowora E. Medicinal plants and Medicine in Africa. 2nd edn. USA: John Wiley and Sons; 1993.
12. Jean BP, Franklin RC, Patricia AB, et al. Performance standards for antimicrobial disk susceptibility tests: approved standard M2-A7. *National Committee for Clinical Laboratory Standards*. NCCLS 1997;35(1):1-15.
13. Lin J, Opoku AR, Geheeb KM, et al. Preliminary screening of some traditional Zulu medicinal plants for anti-inflammatory and antimicrobial activities. *J Ethnopharmacol*. 1999;68(1):267-274.
14. El-Mahmood A, Doughari J. Phytochemical screening and antibacterial evaluation of the leaf and root extracts of *Cassia alata* Linn. *Afr J Pharm Pharmacol*. 2008;2(7):124-129.
15. Begum D, Nath SC. Ethnobotanical review of medicinal plants used for skin diseases and related problems in Northeastern India. *Journal of herbs, spices & medicinal plants*. 2000;7(3):55-93.
16. El-Mahmood AM, Doughari JH, Chanji F. *In vitro* antibacterial activities of crude extracts of *Nauclea latifolia* and *Daniella oliveri*. *Scientific Research and Essay*. 2008;3(3):102-105.
17. Ncube NS, Afolayan AJ, Okoh AI. Assessment techniques of antimicrobial properties of natural compounds of plant origin: current methods and future trends. *African journal of biotechnology*. 2008;7(12):1797-1806.
18. Nikaido H, Vaara M. Molecular basis of bacterial outer membrane permeability. *Microbiol Mol Biol Rev*. 1985;67(4):593-656.
19. Obumselu FO, Onwukeme VO, Eze R. Phytochemical and antibacterial analysis of the leaf extracts of *Ricinus communis*. *Journal of Basic Physical Research*. 2011;2(2).
20. Adegoke Sa, Agada fd, Ogundipe lo. Antibacterial activity of methanol and ethanol leaf extracts of *Antidesma venosum* and *Lannea barteri*. *African Journal of Microbiology Research*. 2013;7(27):3442-3447.
21. Benjamin T, Lamikanra A. Investigation of *Cassia alata*, a plant used in Nigeria in the treatment of skin diseases. *Quarterly Journal of Crude Drug Research*. 1981;19(2-3):93-96.
22. Parekh J, Jadeja D, Chanda S. Efficacy of aqueous and methanol extracts of some medicinal plants for potential antibacterial activity. *Turk J Biol*. 2006;29(4):203-210.
23. Dave H, Ledwani L. A review on anthraquinones isolated from *Cassia species* and their applications. *Indian Journal of Natural Products and Resources*. 2012;3(3):291-319.