Stability evaluation and degradation kinetics of rutin in *Ficus pumila* leaves formulated with oil extracted from *Moringa oleifera* seeds

Abstract

Ficus pumila leaf and the seed oil of *Moringa oleifera* contain significant amount of vitamins and vital nutrients which mitigate vulnerable diseases. Essentially, the goal is to screen the stability of rutin, a valuable antioxidant and validate the mechanism of its kinetic degradation in *Ficus pumila* leaf with and without oil extracted from *M. Oleifera* seeds. Experiments were planned according to standard methods and practices and rutin degradation in both raw leaf extract and the extract formulated with the seed oil were investigated at varying temperatures (25-100°C) and at different time intervals. The results showed that reaction kinetics through heat treatments of the rutin were well characterized by first-order kinetics. The activation energy for the rutin degradation in the raw leaf extract and the extract treated with *M. oleifera* seed oil were 2.1 and 6.9 Kcal/mol respectively. The shelf life of rutin in the formulated extract at 25°C was approximately ten times higher than that of the raw extract. The results indicate that *M. oleifera* seed oil incorporated in food preparation containing rutin enhances rutin enrichment and stability.

Keywords: baobab, seed oil, antioxidant enrichment, kinetics, activation energy, shelf life, *moringa oleifera*, *ficus pumila*

Volume 3 Issue 2 - 2016

Addai Mensah Donkor, Yussif Nashirudeen, Jennifer Suurbaar

Department of Chemistry and Biochemistry, University for Development Studies, Ghana

Correspondence: Addai-Mensah Donkor, University for development studies, Ghana, Tel +233509125157, Email addamenshdonkor@gmail.com, dr.donkor@uds.edu.gh

Received: September 16, 2016 | **Published:** September 29, 2016

Introduction

Oxidative stress is the condition ensuing from the disproportion in the level of pro-oxidant and antioxidant. Oxidative stress is activated by free radicals, which seek stability by way of electron pairing with biological macromolecules such as protein and DNA in human cells in good physical shape and cause protein and DNA impairment along with lipid peroxidation. These changes result in cancer, atherosclerosis, cardiovascular ailments, ageing and inflammatory syndromes. Epidemiological studies have sturdily advocated that intake of certain plant materials may ease the risk of chronic illnesses related to oxidative stress on account of their antioxidant activity and promote broad health benefits.^{1,2} Leafy vegetables apart from being a source of minerals similarly encompass antioxidant vitamins and pigments and are also recognized for their therapeutic importance. Although extensive research on the antioxidant activity of medicinal plants is published yearly, only a few plants have been comprehensively studied for their antioxidant stability, kinetics and enrichment.

Natural products, expressly of plant origin are excellent source of finding antioxidants, since they contain a much higher molecular diversity to those derived from synthetic product. For decades, numerous plants have been used in the treatment of a number of diseases, for which *Ficus pumila* is no exception. Rutin in the leaves of *F. Pumila* has been found to exhibit antioxidant properties and thus has potential against cancer, cardiovascular disorders, neuro-inflammatory ailments, neuropsychiatric syndromes, oxidative stress related diseases and parasitic infections.³

The human body has inimitable enzymatic antioxidants such as superoxide dismutase, catalase and glutathione peroxidase that safeguard the cells from oxidative damage. However, the imbalance of pro-oxidant and endogenous antioxidant in the body leads to deleterious effect. Therefore, the human body requires "reinforcement", which is antioxidant from food and plant source to shield against oxidative stress. Natural antioxidants from plants are widely studied currently not just as preferred food preservatives⁴ but

also their potential in preventing oxidative damages in the body.⁵ Epidemiological studies exhibited that antioxidants from plants such as polyphenols were effective in the protection against cancer and cardiovascular diseases.⁶

The characterization of the oil from the seeds of *Moringa oleifera* has shown that the oil could be exploited successfully as a source of edible oil for human consumption. It is found to contain high monounsaturated to saturated fatty acids ratio, and might be an acceptable substitute for highly monounsaturated oils in diets especially in the developing world. The leaves of *Ficus pumila* has been traditionally consumed either as a beverage or used as an invaluable medicinal herb to treat diabetes, dizziness, high blood pressure, and neuralgia.^{7,8} Research has shown that some of the compounds in the plant have powerful antioxidant and positive hormonal effects⁹ and few side effects have been associated with using them as food, condiment, spice as well as tea. Both the leaves and fruits of *F. pumila* have been used in Chinese traditional herbal medicine since ancient times for treating bleeding, swelling, hemorrhoids, intestinal disorders and impotence and the bioactive compounds were found to be flavonoids.

The antioxidant activity of substances present in several plant species has been extensively studied which reflects their fundamental role in the fortification of skin tissue against the harmful action of reactive oxygen species. Given the significance of effective and long-lasting protection against ultraviolet radiation, we studied the influence of heat treatment over time on rutin, one of the important antioxidants, in both raw extract of *Ficus pumil* and the extract formulated with oil extracted from *Moringa oleifera* seeds. Several chemical *in vitro* methods may be used to evaluate rutin content in plants, among which the ammonium molybdate complexation method stands out, despite its un-specificity, as the most cited and described method in the literature. In this work the extent and kinetics of degradation at variable temperatures were evaluated at different times up to 3 hours from the baseline reading.

This study aimed at investigating the enrichment and stability of rutin in the leaves extract of *F. pumila* treated with oil from *M. oleifera* seeds to ascertain if the oil has some level of rutin and its ability to protect the rutin in the leaves from degradation at high temperature. Specifically, to find out the level of rutin content in the leave extract of *F. Pumila* -formulated with oil from *M. oleifera* seeds and the kinetics of degradation of rutin at variable temperatures.

Materials and Methods

Preparation of plant material

The leaves of *F. pumila* L. were collected from the campus of Kwame Nkrumah University of Science and Technology, Ghana. The plant was identified and authenticated by a Botanist, Dr. Isaac Sackey at the Department of Applied Biology and University for Development Studies Navrongo Campus. Mature pods of *Moringa oleifera* were collected to obtain approximately 400g of seeds from St. John Bosco College of Education, Navrongo, in the Upper East Region of Ghana. The seeds were removed from the pods, sun-dried and stored at 25°C until further use. Commercial rutin, 25grams, and 12.5cm size of Fluted Filter Paper were purchased from Benburto Chemical Enterprises Ltd, Accra, Ghana.

Preparation of *F. pumila* leaf extracts

The leaves were washed and air dried at room temperature (25°C) for three days. The dried leaves were pulverized into powdered form. The powdered sample, 80g was weighed and macerated in 80% methanol at room temperature for 24hours. The extract was filtered and the residue re-extracted twice under the same conditions with a total volume of 300ml 80% methanol. The combined extract was kept in a refrigerator at about 10°C until further analyses.

Extraction of *M. Oleifera* seed oil

The seed kernels were pulverized into fine powder using mortar and pestle and the powdered seed kernel, 40g, was used for the extraction process applying a Soxhlet extractor. Mixture of chloroform/methanol (50:50,v/v), 200ml, was measured into separate 250ml round bottom flask and the Soxhlet with a thimble containing the seed powder and a condenser were assembled. The solvent mixtures were refluxed for 4hours and the mixture was concentrated using rotary evaporator and oven dried for 1hour to obtain light yellowish oil, yield of 11.48g.

Rutin assay

Analysis of the rutin content in the leaf extracts was carried out employing the method described elsewhere.¹⁰ The mixtures of the extracts were heated at varied temperatures (25°C, 40°C, 60°C, 80°C and 100°C), at time interval of 30 minutes up to 150 minutes. The absorbances of the samples were determined using UV-Vis spectrophotometer at (λ_{max} = 360nm). Standard rutin calibration curve was constructed with distilled water as blank. Based on the measured absorbances of the extracts, the concentration of rutin in each sample was determined (mg/ml) using the calibration curve and the content of rutin in the extract was expressed in terms of rutin equivalent (mg/g dry weight of leaf), equation 1. All samples were analyzed in triplicates.

$$R = \frac{A_{sample} * C * 100}{A_0 * W * 0.5} \quad (1)$$

R(mg/g) -plant extract in rutin equivalents; A_{sample}- Absorbance

of sample; A_o- Absorbance of standard; C- concentration of standard rutin solution (mg/ml); W- weight of plant sample (g).

Formulation of *Ficus pumila* leaf extract with *Moringa oleifera* seed oil.

Moringa oleifera seed oil, 0.5ml was added to 0.5ml of *Ficus pumila* leaf extract. The mixture was stirred continuously at room temperature for 5minutes. 2ml distilled water and 5ml of 5% Ammonium molybdate solution were added and transferred into 100ml and diluted with distilled water. A mixture of 0.5ml *Moringa Oleifera* seed oil and distilled water was used as blank. The absorbance was determined using UV-Vis spectrophotometer at (λ_{max} = 360nm). The formulation process was continued with samples heated at various temperatures ranging from 25, 40, 60, 80 and 100°C accordingly, at 0, 30, 60, 90, 120 and 150 minutes.

Kinetic studies of rutin in *F. Pumila* extract at variable temperatures.

Degradation constant analysis

The observed degradation rate constants, kobs, were calculated from the slopes of semi- logarithmic plots of the drug fraction remaining versus time in accordance with Equation 2.

$$\begin{aligned} \ln [C] - \ln [C]_0 &= -kt \\ \ln [C] &= \ln [C]_0 - kt \\ C &= C_0 e^{-kt} \\ \log C &= \log C_0 - \frac{kt}{2.3030} \end{aligned} \quad (2)$$

A plot of ln[C] vs. time *t* gives a straight line with a slope of $-k$, where C₀ was the initial concentration and C is the remaining concentration of rutin at time, *t*. Zero-order kinetics was also investigated and the degradation rate constants were calculated from the slopes of concentration plots of the rutin fraction remaining versus

$$Ct - C_0 = -kt \quad (3)$$

time in accord with Equation 3.

A plot of [Ct] vs. time *t* gives a straight line with a slope of $-k$, where C₀ is the initial concentration and Ct was the remaining concentration of rutin at time, *t*. Arrhenius noted that the rate of degradation with respect to temperature, *k*(*T*) for many reactions fit the equation below:

$$\begin{aligned} k(T) &= Ae^{-\frac{E_a}{RT}} \\ \ln k &= \ln A - \frac{E_a}{RT} \end{aligned} \quad (4)$$

Where A and k are constants characteristic of the reaction and R is the gas constant.

E_a is the energy of activation and A is the pre-exponential factor or the Arrhenius factor. The units of A are the same as those of k. E_a is usually expressed in kcal/mol or kJ/mol. In most cases, A is considered to be temperature-independent.¹¹ This will also be applicable in this current study. If the Arrhenius equation is obeyed, a plot of lnk versus 1/*T* is a straight line with slope $-\frac{E_a}{2.303R}$ and intercept lnA. This allowed E_a and A to be evaluated.

Result and discussion

The study was to evaluate the kinetics of degradation of rutin in both raw leaf extract of *F. pumila* and the extract treated with *M. oleifera* seed oil at varying temperatures. From the data collected at different temperatures, reduction in rutin concentration was monitored by formation of complex with aluminum molybdate. The results of the kinetic experiments showed reduction in rutin concentration in both samples, although the rate of reduction was slow at all temperatures for the formulated extract compared with the raw extract (Figure 1).

Figure 1 Percent of rutin content in dry weight of *Ficus pumila* raw leaf extract and extract formulated with *Moringa oleifera* seed oil at variable temperature.

Figure 2 shows the degradation profile of rutin in *F. pumila* leaf at varying temperatures for the raw extract and for the graph depicted in Figure 3 is the profile for the formulated extract with *M. oleifera* seed oil. As projected, higher temperature promoted high rutin degradation in the raw extracts.

Figure 2 First order degradation kinetics of rutin in raw extract at variable temperatures, $T = 25-100^{\circ}\text{C}$.

In this study, preliminary rutin concentration was 8.47mg/g found in the raw leaf extract while 8.96mg/g was estimated in the formulated extract with the seed oil. It was detected that there was no significant change in rutin content at 25°C for both the raw and the formulated extracts for the treatment period of two hours. The level of rutin in the formulated extract gradually reduced from 6.16mg/g at 40°C to 4.2mg/g at 60°C and further from 3.7mg/g at 80°C and 2.7mg/g at 100°C , during the treatment time. A parallel trend with more rapid decline was witnessed for the raw extract. The rutin content reduced from 5.7mg/g to 0.85mg/g at temperatures ($40-100^{\circ}\text{C}$) during the two hour treatment retro.

The percent decline of the original concentration of rutin in the raw extract at the two hour storage conditions was from 6.1% to 10.5% to 22.3% to 33.1% at 40°C . At maximum temperature of 100°C , and storage time of 2 hours, the percent reduction was drastic, from 57.1% to 72.8% to 77.3% to 83.8 and 97.4 respectively (Figure 4). The

gradual degradation of the formulated sample is apparent in the percent reduction distribution (Figure 5). At 40°C , the percent reduction was 47% at treatment time of one hour. When the temperature was raised to 100°C for an hour, the percent reduction was about 69% relative lower than the raw extract which recorded about 91% reduction, which supports the antioxidant capacity of the *M. oleifera* seed oil. Research by Donkor et al.,¹² reported low temperature treatment of edible plants as relevant in order to protect some vital nutritional components against degradation.

Figure 3 Plots of first order degradation kinetics of rutin formulated with *Moringa oleifera* seed oil at variable temperatures, $T = 25-100^{\circ}\text{C}$.

Figure 4 Plots of percent rutin reduction in raw *ficus pumila* leaf extract at variable temperatures $T_5 = 25^{\circ}\text{C}$, $T_4 = 40^{\circ}\text{C}$, $T_3 = 60^{\circ}\text{C}$, $T_2 = 80^{\circ}\text{C}$, $T_1 = 100^{\circ}\text{C}$.

Figure 5 Plots of percent reduction of rutin in *ficus pumila* leaf exact formulated with *Moringa oleifera* oil from the seeds at variable temperatures $T_5 = 25^{\circ}\text{C}$, $T_4 = 40^{\circ}\text{C}$, $T_3 = 60^{\circ}\text{C}$, $T_2 = 80^{\circ}\text{C}$, $T_1 = 100^{\circ}\text{C}$.

The reaction rate constant k , was determined for each temperature from the slope of the line obtained by least squares regression analysis. Summary of $\ln[C]$ data at variable temperature is given in Table 1&2. Similarly, summary of rate constant for degradation of rutin at variable temperature is given in Table 3&4. Summary of concentration remaining and rate constant for degradation of rutin at variable temperatures is given in Table 5.

The reaction rate constant for the formulated sample is indicated in Table 6. The k values showed that, the degradation kinetics was

first order, demonstrated by the alteration in k values at different temperatures. Even though, the curves for both first and zero order plots were similar for the raw and the formulated samples, the first order rate constants k varied significantly at different temperatures, indicating first order kinetics for the studies. The curves were steeper specifically at 100°C than at the lower temperatures for both the raw and the formulated samples, demonstrating higher degradation.

Figure 6 Plots of $\ln(k)$ against inverse of absolute temperature for first order degradation kinetics of rutin at variable temperatures (25 -100°C) (□) raw fruit pulp extract (●) leaf extract formulated with the seed oil.

Table 1 Rutin concentration (mg), at variable temperatures for raw leaf extract from first order kinetics.

Time (min)	25°C		40°C		60°C		80°C		100°C	
	Conc. (mg)	Ln C	Conc. (mg)	Ln C	Conc. (mg)	Ln C	Conc. (mg)	Ln C	Conc. (mg)	Ln C
30	8.47	2.14	7.95	2.07	5.02	1.61	4.53	1.51	3.63	1.29
60	8.47	2.14	7.58	2.03	4.61	1.53	4.34	1.47	2.3	0.83
90	8.48	2.14	6.58	1.88	4.51	1.51	3.74	1.32	1.92	0.65
120	8.47	2.14	6.06	1.8	4.24	1.44	3.29	1.19	1.37	0.31
150	8.48	2.14	5.67	1.74	3.49	1.25	2.77	1.02	0.85	-0.2

*Conc., concentration; *LnC, Log_eC

Table 2 Rutin concentration (mg), at variable temperatures for formulated extract with Moringa seed oil from first order kinetics

Time (min)	25°C		40°C		60°C		80°C		100°C	
	Conc. (mg)	Ln C	Conc. (mg)	Ln C	Conc. (mg)	Ln C	Conc. (mg)	Ln C	Conc. (mg)	Ln C
30	8.96	2.19	8.17	2.1	5.94	1.78	5.19	1.65	5.07	1.62
60	8.96	2.19	7.88	2.06	5.48	1.7	5.12	1.63	4.74	1.56
90	8.94	2.19	6.83	1.92	5.39	1.68	4.71	1.55	4.16	1.43
120	8.96	2.19	6.61	1.89	4.31	1.46	4.14	1.42	3.71	1.31
150	8.96	2.19	6.16	1.82	4.2	1.44	3.86	1.35	2.71	1

*Conc., concentration; *LnC-, Log_eC

Table 3 Temperature (K), inverse temperature, $1/T$ (K^{-1}), Rate Constant, k (mins^{-1}) and $\ln k$ for raw leaf extract from first order kinetics

Temperature (K)	$1/T$ (K^{-1})	k (min^{-1})	$\ln k$
298	3.356×10^{-3}	-8E-06	***
313	3.195×10^{-3}	0.003	-5.809
333	3.003×10^{-3}	0.0027	-5.915
353	2.833×10^{-3}	0.0042	-5.473
373	2.681×10^{-3}	0.0114	-4.474

Table 4 Temperature (K), inverse temperature, $1/T$ (K^{-1}), Rate Constant, k (mins^{-1}) and $\ln k$ for formulated extract with Moringa seed oil from first order kinetics

Temperature (K)	$1/T$ (K^{-1})	k (mins^{-1})	$\ln k$
298	3.356×10^{-3}	-2.00E-17	***
313	3.195×10^{-3}	0.0025	-5.991
333	3.003×10^{-3}	0.0031	-5.776
353	2.833×10^{-3}	0.0027	-5.915
373	2.681×10^{-3}	0.005	-5.298

Table 5 Data for rutin amount (mg), at variable temperatures for the raw leaf extract and the extract formulated with Moringa seed oil from zero order kinetics

Time (min)	25°C		40°C		60°C		80°C		100°C	
	Concentration (mg)		Concentration (mg)		Concentration (mg)		Concentration (mg)		Concentration (mg)	
	Raw	Formulated	Raw	Formulated	Raw	Formulated	Raw	Formulated	Raw	Formulated
30	8.47	8.96	7.95	8.17	5.02	5.94	4.53	5.19	3.63	5.07
60	8.47	8.96	7.58	7.88	4.61	5.48	4.34	5.12	2.3	4.74
90	8.48	8.94	6.58	6.83	4.51	5.39	3.74	4.71	1.92	4.16
120	8.47	8.96	6.06	6.61	4.24	4.31	3.29	4.14	1.37	3.71
150	8.48	8.96	5.67	6.16	3.49	4.2	2.77	3.86	0.85	2.71

Table 6 Data for inverse of temperature, $1/T$ and rate constant, k for Arrhenius plots of raw leaf extract and the extract formulated with Moringa seed oil from first and zero order kinetics

Tempt. (K)	$1/T (K^{-1}) \times 10^{-3}$	k (mins ⁻¹) Zero Order		k (mins ⁻¹) First Order		Ln k Zero Order		Ln k First Order	
		Raw	Formulated	Raw	Formulated	Raw	Formulated	Raw	Formulated
298	3.355	-0.007	0	-8.00E-06	-2.00E-17	***	***	***	***
313	3.195	0.02	0.018	0.003	0.0025	-3.897	-4.04	-5.809	-5.991
333	3.003	0.011	0.016	0.0027	0.0031	-4.474	-4.167	-5.915	-5.776
353	2.833	0.015	0.012	0.0042	0.0027	-4.186	-4.415	-5.473	-5.915
373	2.681	0.022	0.019	0.011	0.005	-3.835	-3.953	-4.474	-5.298

*Tempt, temperature, *Raw, raw extract

Conclusion

The application of extracted oil from *M. oleifera* seeds to the leaf extract increased the total rutin content and protected it from degradation at higher temperatures. It seems reasonable to consider the *M. oleifera* seed oil and the leaves of *F. pumila* as a remarkable food source for rutin supplement. Preservation of rutin in leaves of *F. pumila* by heat treatments, such as boiling for producing soup or high-temperature short-time treatments for producing clarified juice is a good alternative for enhancing the broad-spectrum of public's food quality intake.

Acknowledgements

We credit the Department of Applied Chemistry and Biochemistry as well as the Head of the Microbiology Laboratory of the Department of Applied Biology, Faculty of Applied Sciences and University for Development Studies, Navrongo.

Conflicts of Interest

It is hereby declared that the authors have no competing financial interests whatsoever in relation to the work described here. It is purely for academic and intellectual purposes.

Funding

None.

References

- Schafer FQ, Buettner GR. Redox environment of the cell as viewed through the redox state of the glutathione disulfide/glutathione couple. *Free Radic Biol Med.* 2001;30(11):1191–1212.
- Rahman K. Studies on free radicals, antioxidants, and co-factors. *Clin Interv Aging.* 2007;2(2):219–236.
- Kaur J. Pharmacognostical and Preliminary Phytochemical Studies on the Leaf Extract of *Ficus Pumila* Linn. *Journal of Pharmacognosy and Phytochemistry.* 2012;1(4):1–7.
- Pokorny JNY, Gordon MH. Antioxidants in food: practical applications. CRC press, USA; 2001.
- Wiseman H. Dietary influences on membrane function: importance in protection against oxidative damage and disease. *The Journal of Nutritional Biochemistry.* 1996;7(1):2–15.
- Arts IC, Hollman PC. Polyphenols and disease risk in epidemiologic studies. *Am J Clin Nutr.* 2005;81(1):317S–325S.
- Mitsuhashi HMO, Nunome S, Terabayashi S, et al. Illustrated medicinal plants of the world in colour. Hokuryukan, Tokyo, Japan; 1988. 493 p.
- Leong CNA, Tako M, Hanashiro I, et al. Antioxidant flavonoid glycosides from the leaves of *Ficus pumila* L. *Food Chemistry.* 2008;109(2):415–420.
- Nakatani N. Natural antioxidants from spices. in ACS symposium series. 1992;507:72–86.
- Atanassova M, Bagdassarian V (x) Rutin content in plant products. *Journal of the University of Chemical Technology and Metallurgy.* 1992;44(2):201–203.
- Starink M. A new method for the derivation of activation energies from experiments performed at constant heating rate. *Thermochimica Acta.* 1996;288(1):97–104.
- Donkor AM, Tei M, Suurbaar J, et al. Stability Evaluation and Degradation Kinetics of Ascorbic Acid in Baobab Fruit Pulp Formulated with the Seed Oil. *British Biotechnology Journal.* 2014;4(5):566.