

# *In vitro* evaluation of Anticancer Activity of Methanolic Extract of *Annona reticulata* Linn. (Ramphal) Leaves on Different Human Cancer Cell Lines

## Abstract

Cancer is one of leading cause of death worldwide and sustained focus on development of novel anticancer agents from medicinal plants. Traditional medicinal systems are based on plant as a source of medicine for the treatment of chronic debilitating and life threatening disease and disorders. *Annona reticulata* Linn. is widely used in the traditional medicine for the treatment of various disease conditions. Methanolic leaves extract of *Annona reticulata* Linn. Was investigated for anticancer potential using sulforhodamine B (SRB) cytotoxicity assay against colon cancer (HCT15), Human lung cancer (Hop65) and Human hepatoma (HEPG2) cell lines. Adriamycin was used as a standard to compare the results. The extract exhibited anticancer effect against all the cell lines. The median growth inhibition (GI50) concentration for extract was <10 µg/ml against Human lung cancer and hepatoma cell line, showing moderate anticancer efficacy of methanolic extract against these cell lines. In conclusion, methanolic leaves extract of *Annona reticulata* Linn. Possess moderate anticancer activities *in vitro* which could be attributed to the presence of anticancerous phytochemicals like acetogenins.

**Keywords:** Medicinal plant; *Annona reticulata* Linn.; Anticancer activity; Sulforhodamine B (SRB) assay

## Research Article

Volume 3 Issue 8 - 2016

**Ashok P Gingine<sup>1\*</sup>, Mandge SV<sup>2</sup>, Prasad G Jamkhande<sup>1</sup>**

<sup>1</sup>SRTM University, India

<sup>2</sup>Department of Botany, India

**\*Corresponding author:** Ashok P Gingine, Assistant Professor, School of Education, Sami Ramanand Teerth Marathwada University, Dnyanteerth, Vishnupuri, Nanded 431 606, Maharashtra, India, Tel: +919822268485; Fax: +91-2462-229242; E-mail: gingine@rediffmail.com

**Received:** October 31, 2016 | **Published:** December 30, 2016

**Abbreviations:** LC: Lethal Concentration; TGI: Total Growth Inhibition; HCT: Human Colon Cancer; GI: Growth Inhibition; SRB: Sulphorhodamine B

## Introduction

Cancer is deadly disease and a major cause of mortality. The incidences of cancer are continuously increasing worldwide [1]. According to WHO, the incidences of cancer will reach to 15 million until 2020. Abnormal uncontrolled growth of cells leads to the development of cancer [2]. From the ancient time, plants and plant based products are the first choice as medicine. WHO specified that till date near about 20,000 plant species studied for medicinal purposes [3]. The phytochemicals obtained from plant such as alkaloids, terpenoids, flavonoids, glycosides and phenolics exhibits meticulous physiological effects. These effects mainly include desirable beneficial effects [4]. The plant based medicines were found to be safer than synthetic chemotherapeutic agents. Toxicity, manifestation of deleterious side-effects, and a narrow margin of error are the major limitations in the use of chemotherapeutic agents [1]. Development of new anticancer agent having minimum or no side effects is the need of time.

*Annona reticulata* Linn. Commonly known as custard apple is a traditionally important plant used for the treatment of various ailments. Traditionally plant is used for the treatment of dysentery, fever and as insecticide [5]. Different plant parts are effectively employed as folk medicine in the rural area of India. The plant is rich in several phytochemicals such as dopamine,

salsolinol, coclaurine, sesquiterpenes and acetogenin [6]. It also contains minerals such as Ca, P, K, Mg, Na, Cl, S, Mn, Zn, Fe, Cu, Se, Co, Ni and Cr [5]. In account with the medicinal properties of *Annona reticulata* Linn. was targeted for the investigation of anticancer property.

## Materials and Methods

### Chemicals

All the solvents and reagents were of analytical grade purity and obtained from Rankem (India).

**Collection and authentication of plant material:** The leaves of *Annona reticulata* Linn. was identified and authenticated by Dr. Gachande B.D., Botanist, Associate Professor of Botany department, N. E. S. Science College, Nanded, India.

**Extract preparation:** Collected *Annona reticulata* Linn. leaves were converted to small pieces and dried under the shade at room temperature. Powder of dried leaves (200g) was prepared using grinder. Methanolic extract was obtained by Soxhlet extractor using 1 L methanol for 8 h at 64 °C and sample was concentrated with the help of rotary evaporator. The 21 % yield of methanol extract was obtained.

**Preliminary phytochemical investigation:** Methanolic extract leaves of *Annona reticulata* Linn. was investigated for phytochemicals using standard procedures [7].

**Anticancer activity using SRB assay:** Sulphorhodamine B (SRB) assay was employed for screening of anticancer activity of methanolic leaves extract of *Annona reticulata* Linn using Human colon cancer cell line (HCT15), Human lung cancer cell line (Hop65) and Human hepatoma cell line (HEPG2). Cell lines were cultured in medium RPMI 1640 containing 10% fetal bovine serum, 2mM L-glutamine and inoculated into 96 well microtiter plates in 100µL at plating densities. Cell were inoculated and microtiter plates were incubated at 37°C, 5 % CO<sub>2</sub>, 95% air and 100% relative humidity for 24h prior to addition of extract. Initially extract was solubilized in dimethyl sulfoxide at 100mg/ml and diluted to 1mg/ml using water and stored frozen prior to use. During extract addition, an aliquote of frozen concentrate (1mg/ml) was thawed and diluted to 100µg/ml, 200µg/ml, 400µg/ml and 800µg/ml with complete medium containing test article. Aliquots of extract (10µl) was mixed to appropriate microtiter wells containing 90µl of medium and final extract concentrations of 10µg/ml, 20µg/ml, 40µg/ml, 80µg/ml were obtained. The plates with extract concentrations were incubated at standard conditions for 48 hours and assay was terminated by the addition of cold TCA. 50µl of cold 30 % (w/v) TCA (final concentration, 10% TCA) was added to fix the cells in situ and incubated for 60 minutes at 4°C. Supernatant fluid was discarded; plates were washed five times with tap water and air dried. In each well sulforhodamine B (SRB) solution (50µl) at 0.4 % (w/v) in 1 % acetic acid was added and it was incubated for 20 minutes at room temperature. One staining is completed, unbound dye was recovered and the residual dye was removed by washing five times with 1 % acetic acid. The plates were air dried and bound stain was subsequently eluted with 10 mM trizma base and absorbance was observed at a reference wavelength of 540nm with 690nm. The percentage growth was calculated on a plate-by-plate basis for extract wells relative to control wells. Adriamycin (Doxorubicin), standard anticancer drug was used as positive control. The percentage growth was calculated at each of the extract concentration levels by using the six absorbance measurements [time zero (Tz), control growth (C), and test growth in the presence of extract at the four concentration levels (Ti)]. The percent growth was calculated using following formula.

Percent growth = (average absorbance of the extract well/ average absorbance of the control wells) X 100

Percentage growth inhibition, growth inhibition, total growth inhibition (TGI) and LC50 was calculated. Percentage growth inhibition was calculated as  $[(Ti - Tz) / C - Tz] \times 100$  for concentration for which  $Ti \geq Tz$  ( $Ti - Tz$ ) positive or zero and  $[(Ti - Tz) / Tz] \times 100$  for concentration for which  $Ti < Tz$  ( $Ti - Tz$ ) negative. Dose response parameters for each test article were calculated. GI50; Growth inhibition of 50% was calculated from  $[(Ti - Tz) / (C - Tz)] \times 100 = 50$ . The drug concentration resulting in total growth inhibition (TGI) was calculated from  $Ti = Tz$ . The LC50, concentration which inhibits 50% of protein measured from beginning was calculated from  $[(Ti - Tz) / Tz] \times 100 = -50$ . GI50 value of  $\leq 20\mu\text{g/ml}$  is considered to demonstrate activity.

Above three parameters were calculated only when the level of activity was observed. The values were expressed as greater or less than maximum or minimum concentration tested when the effect was not reached or exceeded [8,9].

## Results and Discussion

### Phytochemical screening

Preliminary phytochemical screening of methanolic extract of *Annona reticulata* Linn. Leaves showed the presence of major phytochemicals such as alkaloid, amino acids, flavonoid, glycosides, phenolic compound, proteins, steroids and triterpenoid.

### Anticancer activity

Anticancer efficacy of methanol extract of *Annona reticulata* Linn. Leaves was screened using Human colon cancer cell line (HCT15), human lung cancer cell line (Hop65) and human hepatoma cell line (HEPG2) at 10, 20, 40 and 80 (µg/ml) concentrations (Table 1 & 2).

**Table 1:** % Control growth of cell lines in presence of methanolic extract of *Annona reticulata* Linn. leaves and standard adriamycin.

% Control Growth				
Concentrations (µg/ml)	10	20	40	80
<b>Human colon cancer cell line (HCT15)</b>				
Methanolic extract				
Experiment 1	76.5	69.8	54.1	21.5
Experiment 2	62.2	60.7	52.9	25.8
Experiment 3	65.5	62.8	46.4	14.8
Average values	68.1	64.5	51.1	20.7
<b>Adriamycin (ADR)</b>				
Experiment 1	-11.9	-21.6	-25.3	-39.8
Experiment 2	-28.1	-28.5	-36.1	-47.4
Experiment 3	-46.6	-60	-62.8	-63.7
Average values	-28.9	-36.7	-41.4	-50.3
<b>Human lung cancer cell line (Hop65)</b>				
Methanolic extract				
Experiment 1	46.4	45.2	44.9	44.2
Experiment 2	40	39.4	37.9	37.5
Experiment 3	38.8	36.6	35.8	32.2
Average values	41.7	40.4	39.5	38
<b>Adriamycin (ADR)</b>				
Experiment 1	13.6	4.9	3.5	0.6
Experiment 2	7.6	6.5	1.4	-7
Experiment 3	6.3	4.3	-4.1	-17.7
Average values	9.1	5.2	0.2	-8
<b>Human hepatoma cell line (HEPG2)</b>				
Methanolic extract				
Experiment 1	49.1	41.3	35.6	32
Experiment 2	35.4	30.4	30.4	24.7
Experiment 3	41	26.5	15.1	12.4
Average values	41.8	32.7	27	23
<b>Adriamycin (ADR)</b>				
Experiment 1	-10.8	-18.8	-43.3	-51.9
Experiment 2	-19.9	-22.1	-43.1	-53.1
Experiment 3	-20.4	-25.4	-51.8	-56.4
Average values	-17	-22.1	-46.1	-53.8

**Table 2:** lethal concentration value LC50 (µg/ml), total growth inhibition (TGI) and median growth inhibition (GI50) for tested methanolic extract of *Annona reticulata* Linn. and adriamycin.

Cell line	Name of Drug	Lc50	Tgi	Gi50
HCT15	Extract	>80	>80	39.1
	Adriamycin	61.9	21.8	<10
HOP62	Extract	>80	>80	<10
	Adriamycin	>80	54.5	<10
HEPG2	Extract	>80	>80	<10
	Adriamycin	61.2	25.4	<10

Extract inhibited percent control growth of all the cell lines in dose dependent manner. The effect is more in Human colon cancer cell line (HCT15) and Human hepatoma cell line (HEPG2) at 80 µg/ml. While, the effect is least in Human lung cancer cell line (Hop65). The lethal concentration value (LC50) and total growth inhibition (TGI) for all the cell lines were >80µg/ml. The median growth inhibition (GI50) concentration for extract was <10µg/ml against Human lung cancer and hepatoma cell line, showing anticancer efficacy of methanolic extract. However, median growth inhibition (GI50) concentration against Human colon cancer cell line is 39.1µg/ml which indicated no anticancer effect of extract.

Several previous studies showed that plant extract contains abundant number of phytochemicals which possess anticancer properties and might be responsible for anticancer effect of these plants extract [10]. Plants of Annonaceous family are rich in acetogenins, an anticancer property phytochemical [6]. The presence of acetogenin may be attributed to anticancer effect of methanolic extract of *Annona reticulata* Linn.

### Conclusion

The present study was carried out to explore anticancer potential of methanolic extract of *Annona reticulata* Linn. Maximum anticancer activity was observed against Human lung cancer (Hop65) and hepatoma (HEPG2) cell line which

might be correlated to its acetogenins and other anticancerous phytochemicals. In future, work will be needed to isolate bioactive constituents of fresh *Annona reticulata* Linn. Leaves extract to locate potential anticancer phytochemical.

### References

1. Akindele AJ, Wani Z, Mahajan G, Sharma S, Aigbe FR, et al. (2014) Anticancer activity of *Aristolochia ringens* Vahl. (Aristolochiaceae). *J Tradit Complement Med* 5(1): 35-41.
2. Safarzadeh E, Shotorbani SS, Baradaran B (2014) Herbal Medicine as Inducers of Apoptosis in Cancer Treatment. *Adv Pharm Bull* 4(1): 421-427.
3. Ashraf A, Sarfraz RA, Rashid MA, Shahid M (2015) Antioxidant, antimicrobial, antitumor, and cytotoxic activities of an important medicinal plant (*Euphorbia royleana*) from Pakistan. *Journal of Food and Drug Analysis* 23(1): 109-115.
4. Andriani Y, Ramli NM, Syamsumir DF, Kassim MNI, Jaafar J, et al. (2015) Phytochemical analysis, antioxidant, antibacterial and cytotoxicity properties of keys and cores part of *Pandanus tectorius* fruits. *Arabian Journal of Chemistry*.
5. Jamkhande PG, Wattamwar AS (2015) *Annona reticulata* Linn. (Bullock's heart): Plant profile, phytochemistry and pharmacological properties. *J Tradit Complement Med* 5(3): 144-152.
6. Jamkhande PG, Wattamwar AS, Kankudte AD, Tidke PS, Kalaskar MG (2015) Assessment of *Annona reticulata* Linn. Leaves fractions for *in vitro* antioxidative effect and antimicrobial potential against standard human pathogenic strains. *Alexandria Journal of Medicine* 52(1): 19-25.
7. Khandelwal KR (2006) *Practical Pharmacognocny techniques & Experiments*. (6<sup>th</sup> edn) Nirali Prakashan Publisher, Pune, pp. 320.
8. Vichai V, Kirtikara K (2006) Sulforhodamine B colorimetric assay for cytotoxicity screening. *Nat Protoc* 1(3): 1112 - 1116
9. Skehn P, Storeng R, Scudiero A, Monks J, McMohan D, et al. (1990) New colorimetric cytotoxicity assay for anticancer drug screening. *J Natl Cancer Inst* 82(13): 1107-1112.
10. Gaidhani SN, Singh A, Kumari S, Levekar GS, Juvekar AS, et al. (2013) Evaluation of some extracts for standardization and anticancer activity. *Indian Journal of Tradional Knowledge* 12(4): 682-687.