Chemical speciation of binary complexes of Pb(II), Hg(II) and Cd(II) with 2-(1,5-Dimethyl-4-hexenyl)-3-hydroxy-5-methyl-1,4-benzoquinone (Perezzone), a Sesquiterpene

Abstract

Binary complexes of Pb(II), Hg(II) and Cd(II) with 2-(1,5-Dimethyl-4-hexenyl)-3-hydroxy-5-methyl-1,4-benzoquinone are reported. Their chemical composition and geometries are supported by elemental analysis and infrared spectroscopy studies. The lead metal content was estimated by flame atomic absorption method. The mercury metal content was estimated by preparing the metallic hydrides using the "cold vapor" atomic absorption technique. Finally, the cadmium metal content was estimated by the calcination, formation of the respective oxide and determined by flame atomic absorption spectrometry method.

Keywords: natural product, toxic metals, chelation therapy, heavy metal toxicity

Volume 5 Issue 4 - 2017

Soriano García M,¹ Erasmo Flores Valverde,² María Rita Valladares Rodríguez²

¹Departamento de Química de Biomacromoléculas, Universidad Nacional Autónoma de México, México

²Departamento de Ciencias Básicas, Universidad Autónoma Metropolitana, México

Correspondence: Soriano-García M, Departamento de Química de Biomacromoléculas, Instituto de Química, Universidad Nacional Autónoma de México, UNAM, Delegación Coyoacán, Ciudad de México 04510, Tel 5255-5622 4569, Fax 5255 5616 2217, Email Soriano@unam.mx

Received: June 27, 2017 | Published: July 21, 2017

Introduction

As a global problem, the potential health effects of metallic hazards should be a matter of public health concern, especially if the emissions of toxic metals into the environment continue at the current rate.¹ Lead has been known since ancient times. Although lead makes up only about 0.0013% of the earth's crust, it is not considered to be a rare element since it is easily mined and refined. Lead affects every organ of the body, especially the bones and teeth, the kidneys, the nervous, cardiovascular, immune and reproductive systems.² Lead and other heavy metals create reactive radicals which damage cell structures including DNA and membranes.³ Mercury is the 80th element of the Periodic table of elements. Mercury exists in three forms: Hg⁰ (zero valent), Hg₂I (Hg I) (monovalent) and HgII (divalent). Our daily intake is less than 0.01 milligrams (about 0.3 grams in a lifetime), and this we can cope with easily. However, in much higher doses it is toxic such as methylmercury is particularly dangerous. Mercury affects the immune system, alters genetics and enzyme systems, and damages the nervous systems, and the senses of touch, taste and vision.⁴ Cadmium enters the environment through volcanic activity and forest fires.⁵ Cadmium affects different kinds of organisms, ranging from microbes to humans. Human exposure to cadmium mainly occurs through cigarette smoking, but exposure can also occur through contaminated food, water or air.⁶ Cadmium is a known carcinogen to mammals.⁷ Cadmium accumulates in plants, where it is detoxified by binding to phytochelatin,⁸ a family of thiol-rich peptides. Perezzone was first reported in 1852 by Leopoldo Río de la Loza⁹ and the roots of *Acourtia cuernavacana* is used in laxative drinks, as a diuretic, analgesic, as a hypoglycemic.¹⁰ The structure of perezzone was determined by using NMR spectroscopic methods.¹¹ and its crystal structure was reported in 1986.¹² The main objective of this paper is to provoke and stimulate debate on the health effects of long-term, low-level exposure of human populations to toxic metals. Herein we report on the synthesis and spectroscopic characterization

of Pb(II), Hg(II) and Cd(II) complexes of perezzone in the solid state, in order to understand the molecular mechanism involve in the elimination of toxic metals in animals.

Materials and methods

Materials

All chemicals were available commercially and the solvents were purified as conventional methods before use.¹³ Perezzone was isolated from dry roots of *Acourtia cuernavacana* spp, and extracted three times by using maceration with hexane at room temperature. As a result orange crystals were obtained. Purity was checked by single spot test.

Preparation of of binary complexes of Pb(II), Hg(II) and Cd(II) with Perezzone

Compound Pb-Perezzone: The solid lead-perezzone complex was prepared by using a glass beaker of 250mL containing 25mL of 1mmol of alcoholic perezzone solution with 25mL of 1mmol aqueous lead acetate. The addition of lead acetate solution was slowly with continuous stirring at room temperature for 1h. The presence of an excess of perezzone in the flask is separated by a decanting process. A dark brown powder was formed in the beaker. The precipitated complexes were separated by filtration and the excess metal acetate was removed by three times washing with 10mL of deionized water. The complex was finally dried at room temperature. The powder was collected using a plastic spatula and the Pb-perezzone complex is ready for the spectroscopic analysis.

Compound Hg-perezzone: The solid mercury-perezzone complex was prepared by using a glass beaker of 150mL containing 25mL of 1mmol of 1mmol aqueous solution of mercury acetate and adding 10mL of 1mmol alcoholic perezzone solution. The addition of mercury acetate solution was slowly with continuous stirring at room temperature until

the brick red was formed in the glass beaker. The product is rested for 12 hours. The precipitated complexes were separated by filtration and the excess metal acetate was removed by three times washing with 10mL of deionized water. The complex was finally dried at room temperature. The powder was collected using a plastic spatula and the Hg-perezona complex is ready for the spectroscopic analysis.

Compound Cd-perezona: The solid cadmium-perezona complex was prepared by using a glass beaker of 150mL containing 25mL of 1mmol aqueous cadmium acetate. Then, adding 15mL 1mmol of alcoholic perezona solution. The addition was slowly and continuous stirring with the help of a solid glass stirrer and the beaker glass surface during 5 minutes, a dark green precipitated was formed. The excess of perezona in the beaker is separated by a decanting process. The precipitated complexes were separated by using a vacuum filtration process and the excess metal acetate was removed by three times washing with 10mL of deionized water. The complex was finally dried at room temperature. The powder was collected using a plastic spatula and the Cd-perezona complex is ready for the spectroscopic analysis.

Results

Keeping in view the applications of perezona complexes, we describe the synthesis, their chemical composition and geometries are supported by elemental analysis and infrared spectroscopy of binary complexes of Pb(II), Hg(II) and Cd(II) with perezona. The bonding of the metals (Pb, Hg and Cd) to the ligand is studied by comparing the IR spectrum of the ligand with those of the complexes. In the

ligand spectrum, bands at 3304.6, 1626.8 and 1283.3 cm⁻¹ are due to intramolecular hydrogen bonded of hydroxyl group at position 3 and the carbonyl oxygen at position 4, the $\nu\text{C}=\text{O}$ and $\nu\text{C}-\text{O}$, respectively. The spectra of the complexes in comparison show shifts in the above mentioned bands due to coordination to Pb and Cd ions. The band due to presence of an intramolecular hydrogen bond involving the hydroxyl group disappears in the spectra of the complexes as a result of the coordination to the metal ion through the oxygen atom of the hydroxyl and abstraction of protons.¹⁴ In the complex of Pb(II), Hg(II) and Cd(II) bands at 3391.0, 2922.9 and 3303 cm⁻¹ respectively are assigned in the stretching modes of coordinated water molecules. The bands at 608.6, 595.8 and 713.9 cm⁻¹ are due to $\nu\text{Pb}-\text{O}$, $\nu\text{Hg}-\text{O}$ and $\nu\text{Cd}-\text{O}$ stretching modes. The color, elemental analysis and IR analysis results are given in Table 1 suggest a molecular formula $[\text{M}(\text{C}_{15}\text{H}_{19}\text{O}_3)_2 \cdot 2\text{H}_2\text{O}]$ where M = Pb(II), Hg(II) and Cd(II).

Discussion

In the present work, the first three new binary complexes of Pb(II), Hg(II) and Cd(II) with Perezona were synthesized, and their color, elemental and IR spectral analysis of these complexes were determined. The experimental data suggest a molecular formula $[\text{M}(\text{C}_{15}\text{H}_{19}\text{O}_3)_2 \cdot 2\text{H}_2\text{O}]$ where M=Pb(II), Hg(II) and Cd(II). The proposed geometries of the Pb, Hg and Cd complexes contain two ligand molecules and two water molecules arranged around a central metal atom defining the vertices of a distorted octahedron.

Table 1 Colour, Elemental Analysis, and IR spectral analysis of the complexes

Complex	Color	Elemental analysis (Calc./Found ^a)			IR Spectral bands (cm ⁻¹)		
		C	H	M	$\nu(\text{C}=\text{O})$	$\nu(\text{C}-\text{O})$	$\nu(\text{M}-\text{O})$
Pb(C ₁₅ H ₁₉ O ₃) ₂	Dark brown	59.0/59.1	6.6/6.5	29.5/30.4	1550.3	1282.4	608.6
Hg(C ₁₅ H ₁₉ O ₃) ₂	Brick red	59.1/60.4	7.0/6.9	28.9/30.4	1553.2	1281.9	595.8
Cd(C ₁₅ H ₁₉ O ₃) ₂	Green	60.5/60.1	8.2/8.2	18.5/21.0	1549.9	1282.1	713.9

Each value corresponds to an average of 5 determinations.

Conclusion

This work brings some useful experimental evidence regarding the molecular interaction between three toxic metal ions (Pb, Hg and Cd) with a Mexican natural product which may provide an incentive to understanding the mechanism of elimination of toxic metals present in the outdoors air pollution in the cities.

Acknowledgements

This work has been carried during my sabbatical leave (M. Soriano-García) at the área de Química, Departamento de Ciencias Básicas, División de Ciencias Básicas e Ingeniería de la Universidad Autónoma Metropolitana-Unidad Azcapotzalco, México. This work was supported by the Cátedra “Leopoldo Río de la Loza Guillén” del Departamento de Ciencias Básicas. All the authors contributed to the biochemical, spectroscopic studies and final editing of the manuscript. We thank the technical assistance from Rocío Patiño (IR) and María de la Paz Orta Pérez (EA) of the Instituto de Química, Universidad Nacional Autónoma de México, UNAM.

Conflicts of interest

Do not exists any financial or conflict of interest.

References

1. Nriagu JO. A silent epidemic of environmental metal poisoning? *Environ Pollut.* 1988;50(1-2):139–161.

2. Freitag CM. The genetics of autistic disorders and its clinical relevance: a review of the literature. *Mol Psychiatry.* 2007;12(1):2–22.
3. Carocci A, Catalano A, Lauria G, et al. Lead Toxicity, Antioxidant Defense and Environment. *Rev Environ Contam Toxicol.* 2016;238:45–67.
4. Grandjean P. Methylmercury toxicity and functional programming. *Reprod Toxicol.* 2007;23(3):414–420.
5. Filipi M, Fatur T, Vudrag M. Molecular Mechanisms of Cadmium Induced Mutagenicity. *Hum Exp Toxicol.* 2006;25(2):67–77.
6. Järup L. Hazards of heavy metal contamination. *Br Med Bull.* 2003;68(1):167–182.
7. Bertin G, Averbeck D. Cadmium: cellular effects, modifications of biomolecules, modulation of DNA repair and genotoxic consequences (a review). *Biochimie.* 2006;88(11):1549–1559.
8. Cobbett CS, Goldsbrough PB (Mechanisms of metal resistance: Phytochelatins and metallothioneins. In: Raskin I, Ensley BD, editors. *Phytoremediation of Toxic Metals Using Plants to Clean up the Environment.* New York, USA. John Wiley and Sons Inc; 1999:247–269.
9. Noriega JM. Speech to the Medical Academy of México. *Escritos de Leopoldo Río de la Loza.* Imprenta de Ignacio Escalante, México. 1911.
10. Alarcón-Aguilar FJ, Román-Ramos R, Jiménez-Estrada M, et al. Effects of three Mexican medicinal plants (Asteraceae) on blood glucose levels in healthy mice and rabbits. *J Ethnopharmacol.* 1997;55(3):171–177.

11. Wagner ER, Moss RD, Brooker RM, et al. A correction of the structure of perezzone. *Tetrahedron Lett.* 1965;47(6):4233–4239.
12. Soriano-García M, Toscano RA, Flores-Valverde E, et al. Structure of 2-(1,5-Dimethyl-4-hexenyl)-3-hydroxy-5-methyl-1,4-benzoquinone (Perezzone), a Sesquiterpene. *Acta Cryst.* 1986;42(3):327–329.
13. Armarego WLF, Perrin DD. Purification of Laboratory Chemicals. 6th edn. Butterworth Heinemann: Oxford, USA. 2009:138–159.
14. Singh O, Kumar A, Singh K. Polymeric complexes of 2,5-Dihydroxy-1,4-benzoquinon. 2012.