Iron and selenium speciation in enriched adzuki bean sprouts after fractionation procedures by graphite furnace atomic absorption spectrometry

Abstract

Selenium (Se) and iron (Fe) are essential trace elements for human health, and their content in foods can vary significantly depending on the food characteristics, cultivation conditions (fertilizers, climate, and water), and geographical origin. The consumption of enriched edible sprouts is an effective strategy to combat nutritional deficiencies of Fe and Se. Notably, the daily Se intake range between essential and toxic levels is very narrow, inorganic Se species are more toxic than the organic forms and soluble Fe species are more bioaccessible than other species. Thus, a rapid method for determination of Fe species (soluble and inorganic) and a sensitive and selective micelle-mediated speciation and preconcentration method with elemental determination by GF AAS was developed to evaluate the effect of Fe and Se enrichment of adzuki bean sprouts, respectively, as well as the evaluation of the enrichment in the concentration of Fe and Se species that are potentially absorbed by the human organism. Fractionation procedures (extraction) were used for the separation of Fe species. The results indicated that the enrichment with Fe(III)-EDTA favoured the translocation to the aerial part of the plant and increased 75% total Fe concentration in the stem. In the proposed methods for Fe (soluble and inorganic) speciation, it was found that the Fe enrichment increased the concentration of soluble Fe (50%) and inorganic Fe (145%) in the edible part of the plant. Therefore, the Fe enrichment procedure applied is a good alternative to help reduce Fe deficiencies and the recommended daily intake. Extraction procedures were used for Se speciation, the method was based on CPE using nonionic surfactant Triton X-114 in the presence of dithizone as a chelating agent, which reacts selectively with Se(IV). The proposed method was successfully applied to the speciation and determination of organic (hydrophobic and hydrophilic) and inorganic (Se(IV) and Se(VI)) species of Se in enriched adzuki bean sprouts. The obtained results show that hydrophilic organic Se species were predominant ($96 \pm 2 \mu\text{g g}^{-1}$), and the enrichment procedure proved to be an appropriate alternative to reduce Se deficiency, due to the presence of Se bioaccessible species.

Keywords: speciation, selenium, iron, enriched adzuki bean sprout

Volume 7 Issue 2 - 2018

Aline Pereira de Oliveira,¹ Cassiana Seimi Nomura,² Sidnei Gonçalves da Silva,³ Juliana Naozuka¹

¹Departamento de Química, Universidade Federal de São Paulo, Brazil

²Instituto de Química, Universidade de São Paulo, Brazil

³Centro de Ciências Exatas e Tecnologia, Universidade Federal de Uberlândia, Brazil

Correspondence: Juliana Naozuka, Departamento de Química, Universidade Federal de São Paulo, Diadema, São Paulo, Brazil, Email jnaozuka@gmail.com, j.naozuka@unifesp.br

Received: February 01, 2018 | **Published:** March 12, 2018

Abbreviations: GF AAS, graphite furnace atomic absorption spectrometry; CPE, cloud point extraction; ICP OES, inductively coupled plasma optical emission spectrometry; LOD, limit of detection; LOQ, limit of quantification

Introduction

The production of edible sprouts is a fast-growing segment of the food industry worldwide. Although several kinds of seeds may be eaten as sprouts, adzuki (*Vigna angularis*) and mung (*Vigna radiata*) beans are preferred and commonly used for sprout production.¹ Studies have shown that grain germination increases the content of essential nutrients, such as minerals, vitamins, and proteins, and reduces the antinutritional factors, such as trypsin inhibitors, phenolic compounds, and nitrate,² enhancing the quality and digestibility of the grains. Moreover, previous studies have clearly demonstrated that it is possible to enrich sprouts with essential elements to combat nutritional deficiencies, which is particularly important for populations in developing countries.³⁻⁶

The consumption of Se-enriched sprouts is a strategy to prevent the deficiency of this element in the human body, which is associated with various disorders including increased cholesterol levels, cardiomyopathy, muscular dystrophy, dysfunction of brain and liver, and Keshan disease, a disease affecting millions of people worldwide, particularly in China.^{7,8} The chemical form of dietary Se is important because it affects its bioavailability, toxicity, and chemoprotective activity. In nature, Se exists in the form of inorganic species, such as selenite [Se(IV)], selenate [Se(VI)], and selenides (e.g., HgSe), or as organic species.^{9,10} In general, the inorganic forms are more toxic^{11,12} and less bioavailable to humans than the organic species.¹³ On the other hand, the consumption of Fe-enriched can increase iron consumption, which helps cellular homeostasis, oxygen transport and energy metabolism, as well as a cofactor of enzymes for mitochondrial respiratory chain and nitrogen fixation.^{14,15} It is important to point out that legumes have low concentrations of Fe species potentially absorbed by the human organism due to interactions of inorganic Fe-species, mainly bivalent cations, with phytic acid, tannins and other anti-nutrients, not being able to supply nutritional deficiencies.^{8,14,15}

The success of elemental enrichment of sprouts depends on chemical species of the elements added to the culture medium, which may be absorbed, translocated, accumulated and converted in others chemical species potentially bioavailable.^{16,17} In this way, it is necessary to determine and identify the species formed after enrichment, establishing rapid analytical methods. Chemical speciation studies in the edible part of the enriched plant are important and may estimate the bioavailability, toxicity and elemental essentiality,¹⁸ promoting knowledge about the conscious use of these foods enriched with essential elements, such as Fe and Se. Therefore, the objective of this study was proposed a method to determine species of Fe (water soluble, acid soluble and inorganic) and Se (Se(IV), Se(VI), and organic (hydrophobic and hydrophilic)) in enriched adzuki bean sprouts, combining GF AAS with derivatization reactions or CPE, for Fe and Se, respectively.

Material and methods

Instrumentation

Total Fe and Se quantification was done by ICP OES, using an equipment (model iCAP 6300 Duo, Thermo Fisher Scientific, England) with axial and radial viewed plasma and a charge-injection device detector, which allows measurements from 166.25 to 847.00nm. An echelle polychromator purged with argon, a radiofrequency source of 27.12MHz, a Meinhard nebulizer combined with a cyclonic spray chamber was used. The instrumental parameters for ICP OES are given in Table 1.

Table 1 Instrumental conditions for Fe and Se determination by ICP OES

Power	1250 W
Plasma gas flow	15 L min ⁻¹
Intermediate gas flow	1.0 L min ⁻¹
Nebulizer gas flow	0.45 L min ⁻¹
Sample introduction	1.5 L min ⁻¹
Analytical wavelength – axial view	
(I) Atomic line	Se(I) = 196.0nm
(II) Ionic line	Fe(II) = 259,8nm

A ZEE nit 60 model atomic absorption spectrometer (AnalytikjenaAG, Jena, Germany) equipped with a transversely heated graphite atomizer, pyrolytically coated graphite tube, and a transversal Zeeman-effect background corrector was used for Fe and Se determination, after fractionation procedures and in phases obtained after CPE, respectively. All measurements were made using a hollow cathode lamp, and were based on integrated absorbance values. The optimized operating conditions of the spectrometer and the heating program are shown in Table 2. Argon 99.998% (v v⁻¹) (Air Liquide Brazil, São Paulo, Brazil) was used as a protective and purge gas. The digestion of samples for total Fe and Se determination was carried out in a closed-vessel microwave system (Speedwave-4, Berghof, Germany), equipped with 16 fluoropolymer vessels and a ceramic vessel jacket.

Table 2 Instrumental parameters and heating program for Fe^a and Se^b determination by GFAAS

Step	T (°C)	Ramp (s)	Hold (s)	Ar flow rate (mL min ⁻¹)
Drying I	100	10	15	250
Drying II	130	10	20	250
Pyrolysis	1200 ^a , 900 ^b	100	20	250
Atomization	2300 ^a , 2100 ^b	2300 ^a , 2100 ^b	5	0
Cleaning	2500	1200	2	250

A thermostatic water bath (model Q226M2, Quimis, Brazil) at 95°C and 55°C was used for iron fractionation procedures with heating and selenium speciation with CPE experiments, respectively. An ultrasonic bath (model 75D Acquasonic, VWR Scientific Instruments, USA) was used to evaluate the effectiveness of total Se extraction for speciation studies. Phases separation was accelerated using a centrifuge (model Q222TM, Quimis, Brazil). For preliminary sample preparation, the stems (edible part) of the adzuki beans were separated and dried using a freeze dryer (Thermo Fisher Scientific, England). Homogenization of the culture medium was performed using an orbital shaker (model 0225M, Quimis, Brazil).

Reagents and samples

Adzuki beans (500g, a single brand) were purchased from the local market of Sao Paulo, Brazil. According to the manufacturer, the grains were produced in Brazil. All solutions were prepared from analytical reagent grade chemicals, using high-purity deionized water obtained from a Milli-Q water purification system (Millipore, USA) as the solvent. Analytical grade 65% (v v⁻¹) HNO₃ distilled in a quartz sub-boiling still (Marconi, Brazil) and 30% (v v⁻¹) H₂O₂ (Merck, Germany) were used for sample digestion.

For Fe speciation were prepared two different precipitant reagents:

- 25g trichloroacetic acid (Merck, Germany) and 25mL hydrochloride acid (Synth, Brazil) and
- 25g trichloroacetic acid (Merck, Germany), 25g hydroxylamine hydrochloride (Merck, Germany), and 25mL hydrochloride acid (Synth, Brazil). A final volume of 250mL was completed by deionized water.

Titrisol standard solution of 1000mg L⁻¹ of Fe (FeCl₃) (Merck, Germany) was used to prepare the reference analytical solutions in 0.1% (v v⁻¹) HNO₃ (10 to 80µg L⁻¹ and 1.25 to 80.0mg L⁻¹ for Fe determination by GF AAS and ICP OES, respectively).

For Se speciation, an acetone solution of dithizone (Merck, USA) was prepared. A 10% (v v⁻¹) ethanol solution (Merck, USA) containing 0.1mol L⁻¹ HNO₃ was used to dilute the surfactant-rich phase, and Na₂SeO₃ (Merck, Germany), Na₂SeO₄ (Sigma Aldrich, USA), selenomethionine (Sigma Aldrich, USA), NaOH (Merck, USA), and

HCl (Merck, USA) were used for optimization of the CPE procedure. Titrisol standard solution of 1000 mg L⁻¹ of Se (SeO₂) was used to prepare the reference analytical solutions, by dilution with 0.1% (v v⁻¹) HNO₃ for total Se determination by ICP OES, and with 0.1% (v v⁻¹) HNO₃ and 5% (v v⁻¹) Triton X-114 or 2mmol L⁻¹ dithizone and 5% (v v⁻¹) Triton X-114 for Se determination in the phases obtained after CPE by GF AAS. For Se determination by GF AAS, solutions of 20g L⁻¹ Pd(NO₃)₂ (Merck) and 10g L⁻¹ Mg(NO₃)₂ (Merck) were used to prepare the chemical modifier. Seed sterilization was performed using NaClO (Sigma Aldrich, USA) and a solution of Na₂SeO₄ (Sigma Aldrich, USA) or (NaFe(III)-EDTA (Sigma Aldrich, USA)) were used for the enrichment procedure.

Cultivation of sprouts and preliminary sample preparation

Cultivation of adzuki bean sprouts was carried out in triplicate for each treatment. The adzuki bean seeds were first cleaned with deionized water. After sterilization for 5 min in 12% (v v⁻¹) NaClO,² the seeds (*ca.* 5g) were cleaned again and soaked in deionized water (*ca.* 200mL) at room temperature for 24h, to promote the breaking of seed dormancy.¹⁹ Next, the seeds were placed on an absorbent cotton (*ca.* 3g) in a glass container (previously decontaminated with 10% v v⁻¹ HNO₃). The seeds were cultivated for 13 days in a portable greenhouse at 70% relative humidity, in the dark, and at a temperature of about 27 and 18°C during the day and night, respectively.

The germination of adzuki bean seeds was carried out with 20mL of deionized water (control group), 20mL of a 150mg L⁻¹ Fe solution (3mg of Fe for Fe(III)-EDTA) or 20mL of a 5mg L⁻¹ Se solution (250µg of Se for Na₂SeO₄) in the growing medium. During germination, 13 days, all sprouts were soaked once a day with 20mL of deionized water. The adzuki bean sprouts were collected and washed with deionized water, and the stem (edible part) was separated from the other plant parts. They were dried by lyophilization (*ca.* 2 days) and ground using a pestle and mortar (decontaminated) after freezing the samples in liquid nitrogen.

Procedure for total Fe and Se determination

The method characteristics for determination of total Fe and Se in enriched adzuki beans sprouts by ICP OES have been published by our group.^{16,17} For total Fe and Se determination, the adzuki beans sprouts were subjected to acid digestion in a closed-vessel microwave system. Sample masses ranging from 150 to 250mg were digested using a diluted oxidant mixture (2mL HNO₃ + 1mL H₂O₂ + 3mL H₂O).²⁰ The heating program consisted of three steps (temperature/°C; ramp/min; hold/min): (1) (140; 5; 1); (2) (180, 4, 5); and (3) (220, 4, 10). The system was cooled by forced ventilation for 20min. Digestion was performed in duplicates for each sample, and the final volume of 10mL was completed with deionized water.

Fe speciation procedure

For the speciation of soluble and inorganic Fe, the adapted procedure proposed by Oliveira et al.¹⁵ was used. Initially it was added 15mL of deionized water to 400mg of stems (control and Fe-enriched groups). These mixtures were agitated for 5min at room temperature and their supernatants were separated by centrifugation

at 4000rpm for 10min. The supernatants were analyzed by GF AAS, obtaining the concentration of Fe in the aqueous extract (Fe soluble and Fe associated with macromolecules). Subsequently, it was added 2.5mL of a precipitant reagent 1 to 5mL of the cloudy supernatant obtained after extraction with water, promoting the precipitation of macromolecules.²¹ The mixture was heated in a thermostatic bath at 95°C, centrifuged at 4000rpm for 10min and the supernatant was analyzed by GF AAS, obtaining the concentration of soluble Fe not associated with macromolecules.²²

Another procedure was to add 2.5mL of the precipitant reagent 2 to 5mL of the aqueous extract. The mixture was heated in a thermostatic bath at 95°C, centrifuged at 4000rpm for 10min and the supernatant was analyzed by GF AAS. This procedure was applied to precipitate macromolecules and promote the reduction of Fe (III) to Fe (II).^{22,23} The determination of Fe in the obtained supernatants was performed by GF AAS and the heating program according to Table 2. The volume of 10µL of the samples or analytical solutions was injected into the graphite tube and no chemical modifier was used in the Fe determination. The calibration of the instrument was done with reference analytical solutions of concentrations ranging from 10 to 80µg L⁻¹ in 0.1 % (v v⁻¹) HNO₃. After calibration, the supernatants obtained were previously diluted 20 times with deionized water before being analyzed. To evaluate possible matrix interference in elemental determination, the addition and recovery test was used. For this purpose, an analytical solution of 20µg L⁻¹ of Fe was added in the presence of the precipitant reagents 1 and 2, without and with addition of hydroxylamine, respectively.

Se speciation procedure: initial extraction

Two extractants were investigated for Se species extraction: deionized water and 1.0mol L⁻¹ NaOH. The dried and ground Se-enriched sprouts (250µg) were subjected to initial extraction with 20mL of the extractants. With the aim to extract all Se species, the extraction was tested in a thermostatic water bath at 80°C and in an ultrasonic bath, each for 30 min. The supernatant (aqueous extract) was isolated by centrifugation at 4000rpm for 15min.

Se speciation procedure: CPE

A volume of 10mL of the aqueous extract (see “Initial extraction”) was mixed with 1mL of 1% (v v⁻¹) nonionic surfactant Triton X-114, and the mixture was heated in a thermostatic water bath at 55°C for 20min. Next, the obtained surfactant-rich phase was separated by cooling in an ice-water bath for 20min followed by centrifugation at 4000rpm for 15min. In this phase, the hydrophobic organic Se species were determined.

The surfactant-poor phase was subjected to a second CPE, using 1mL of 1% (v v⁻¹) Triton X-114 and 0.2mL of 2mmol L⁻¹ dithizone. In this second extraction, the pH of the medium was evaluated, ranged from 1 (adjusted by 0.1mol L⁻¹ HCl) to 9 (adjusted by 0.1mol L⁻¹ NaOH), using solutions containing 500µg L⁻¹ of Se from Se(IV), Se(VI), and Se-methionine (Se-Met). The mixture was kept in a thermostatic water bath at 55°C for 20min. Formation and separation of the surfactant-rich and -poor phases was achieved by cooling in an ice-water bath for 20min followed by centrifugation at 4000rpm

for 15min. After optimization of the CPE procedure, Se(IV) was determined in the surfactant-rich phase.

A second aliquot of 5mL of the aqueous extract was subjected to CPE using only 1mL of 1% (v v⁻¹) nonionic surfactant Triton X-114. After phase separation, the surfactant-poor phase containing hydrophilic organic and inorganic Se species was heated to 55°C for 5 min in the presence of HCl. The HCl concentration was evaluated, ranged from 1 to 6mol L⁻¹, using a solution of Se(VI) (500µg L⁻¹). The HCl-treated surfactant-poor phase was diluted to 10mL with deionized water and subjected to CPE using 1mL of 1% (v v⁻¹) Triton X-114 and 0.2mL of 2 mmol L⁻¹ dithizone.²⁴ Inorganic Se species were determined in the surfactant-rich phase.

Se determination was carried out by GF AAS in the different phases obtained after CPE. Prior to analysis, the surfactant-rich phases were diluted with a 10% (v v⁻¹) ethanol solution containing 0.1mol L⁻¹ HNO₃. Moreover, 10µL of chemical modifier (5µg Pd+3µg Mg) was co-injected with 10µL of extract/phase or analytical solution. The thermal behavior of Se in the absence and presence of Triton and dithizone was studied. The pyrolysis temperature was optimized, ranging from 400 to 1400°C with increments of 100°C at an atomization temperature of 2300°C. At the optimum pyrolysis temperature, the atomization temperature was studied from 1600 to 2300°C with increments of 100°C. The chemical modifier was always co-injected during the analysis. Instrument calibration was performed using analytical solutions with concentrations ranging from 20 to 180µg L⁻¹ in 0.1% (v v⁻¹) HNO₃ or 1% (v v⁻¹) Triton X-114 and 2mmol L⁻¹ dithizone. Chemical interferences during Se determination by GF AAS were evaluated by addition and recovery tests, adding a solution

of 40µg L⁻¹ Se (Se(IV), Se(VI), or Se-Met) in the surfactant phases.

Results and discussion

Analytical methods characteristics

For Fe and Se total determination by ICP OES, the analytical characteristics and evaluation of the chemical interferences and accuracy are shown in the previous worked of the research group.^{16,17} The Fe and Se species determination was conducted by GF AAS. The characteristic parameters of the analytical calibration curves (linear range, correlation coefficient (R²), LOD and LOQ) are shown in Table 3, as well as the results of recovery of the addition and recovery test that was used to verify chemical interferences in the Fe and Se determination by GF AAS. The LOD was calculated from the standard deviation of 10 measurements of the analytical blank sample ($3 \times \sigma_{\text{blank}}$, where σ is the standard deviation), and the LOQ was calculated as $3.3 \times \text{LOD}$.²⁵ The values were obtained in µg g⁻¹. For Fe, the LOD and LOQ values were obtained considering a 400mg sample and a final volume 10mL and 15mL for aqueous extract and precipitant reagents, respectively. For Se, was considered a sample mass of 200 mg and a final volume of 10mL for total Se determination, and a sample mass of 100 mg and a final volume of 2mL for Se speciation. The recoveries ranged between 82 (Se(VI)) and 115 (iron soluble in water) %, indicating the absence of chemical interferences during Fe and Se determination in the presence of the various reagents used for fractionation and CPE, respectively. According to the US Food & Drug Administration for elemental determination by spectrometric techniques, the recovery tolerance for food samples ranges from 80 to 120%.²⁶

Table 3 Characteristics method for Fe and Se determination by GF AAS

			Linear range (µg L ⁻¹)	R ²	LOD (µg g ⁻¹)	LOQ (µg g ⁻¹)	Recovery (%)
Se	Inorganic selenium	Se(IV)	20 – 180	0.9994	0.26	0.79	96
		Se(VI)			0.13	0.39	82
	Organic selenium	Hydrophilic	20 – 180	0.9962	4.29	12.88	98 ^a
		Hydrophobic			0.25	0.75	ND ^b
Fe	Water				0.036	0.118	0.115
	Precipitant reagent 1		10–80	0.9983	0.032	0.105	0.108
	Precipitant reagent 2				0.027	0.090	0.99

^aaddition of 80µg L⁻¹ Se-Met; ^bND = not determined.

Fe speciation in enriched sprouts

The Fe determination in the aqueous extract and after addition of different precipitant reagents was performed by GF AAS. The soluble Fe concentration was obtained in the supernatant from the extraction with water, Fe not associated with macromolecules (including inorganic Fe) in the supernatant of the extraction with the precipitant reagent 1, and the concentration of Fe species not associated to macromolecules and which are not inorganic Fe species, such as the association of Fe to soluble fibers, in the supernatant of extraction with precipitant reagent 2. The Fe concentrations in the supernatants of the aqueous extract and the different precipitant reagents with and without addition of hydroxylamine in the samples of bean sprouts (control and Fe-enriched group), are presented in Table 4.

It has been found that soluble Fe (aqueous extract) is a fraction of total Fe in both culture conditions (control and Fe-enriched group). The enrichment procedure increased the Fe concentration in the aqueous extract, but when compared to the total concentration a decrease in the soluble Fe after enrichment (49% (control) and 42% (Fe-enriched)) was observed. It is possible that the decrease in the concentration of water-soluble proteins (albumins) after enrichment with Fe(III)-EDTA, is one of the causes for the decrease in the soluble Fe percentage.

After addition of the precipitant reagents 1 and 2, it was possible to quantify soluble non-macromolecular and inorganic Fe, respectively. The Fe concentration in the supernatant after addition of precipitant reagent 1 reduced 80 % (control) and 82 % (enriched) when compared

to the Fe concentration in the aqueous extract. The trichloroacetic acid added to the precipitant reagent was used as a precipitant of proteins and peptides, and its efficiency in the macromolecules precipitation has been reported by Carreira et al.²⁷ Therefore, it is possible to deduce that in the stem of the Fe-enriched sprouts and control group there is a significant amount of Fe associated with macromolecules. In addition, it was also verified that the enrichment did not significantly alter the percentage of total Fe associated with macromolecules.

Table 4 Concentrations of iron in the supernatants and the effect of hydroxylamine

	Concentration of Fe ($\mu\text{g g}^{-1}$) $\pm$ standard deviation (n = 3)	
	Control	Fe(III)-EDTA
Total	89 $\pm$ 1	156 $\pm$ 1
Aqueous extract	44 $\pm$ 2	66 $\pm$ 1
Precipitant reagent 1 (without hydroxylamine, pH <1)	8.6 $\pm$ 0.8	12 $\pm$ 1
Precipitant reagent 2 (with hydroxylamine, pH=5.5)	5.5 $\pm$ 0.1	4.0 $\pm$ 0.7
Inorganic Fe ^a	3.1 $\pm$ 0.8	7.6 $\pm$ 0.8

^aSubtraction of the Fe concentration in the extract with hydroxylamine of the extract without hydroxylamine.

In the absence of hydroxylamine, it is possible to obtain the concentration of Fe associated with macromolecules, whereas when adding hydroxylamine, the total inorganic Fe concentration was determined. Comparing the Fe concentration after the addition of the precipitant reagents 1 and 2, it was verified that the Fe concentration decreased 36 and 65% in the control and enriched groups, respectively.

According to Quinteros et al.,²² Kosse et al.²³ the addition of hydroxylamine to the precipitant reagent promotes the reduction of Fe(III) to Fe(II) ions and studies have shown that Fe(II) ions interact with macromolecules that are precipitated with the addition of trichloroacetic acid and other matrix components, such as antinutritional components.^{8,15} The polyphenols, such as tannins, may react with bivalent metal ions, such as Fe(II) through its carboxylic and hydroxyl groups forming insoluble complexes and reducing the absorption of non-heme Fe.^{14,15,17} Furthermore, another antinutrient present in legumes, the phytic acid has high potential chelating. This compound exhibits 12 ionizable protons and can interact with metal ions. Phytate, abundant in legumes, is one of the main inhibitors of non-heme iron absorption, binds mainly to bivalent metal ions and forms insoluble salts, altering the solubility, functionality, digestibility and absorption of Fe.²⁸ In acid or neutral media, such as the precipitant reagent 2 (pH=5.5), the 6 phosphate groups of the phytic acid molecule expose its 12 negative charges, favoring the direct or indirect complexation of this molecule with bivalent cations of Fe(II), starch, proteins, and enzymes.²⁹ In addition, the interaction of Fe(II) ion with phytate molecules is more effective at lower pH, on the other hand, the phytate-Fe(III) complex is more stable when the ligand is deprotonated due to the high Fe(III) (central atom) charge which produces a more effective interaction with phytate and, consequently, more protons are replaced in the ligand at higher pH.³⁰

Therefore, due to significant decrease in the Fe concentration of extracts without and with hydroxylamine of the control and Fe-

enriched group it is possible to infer that the hydroxylamine reacts only with Fe(III) ions reducing them to Fe(II), which is associated with antinutrients, such as polyphenols and phytates, forming products that are not extracted in water or precipitate together the macromolecules when adding trichloroacetic acid. Thus, it is possible to quantify total inorganic Fe in the stem of the azuki bean sprouts, subtracting the Fe concentration in the extract with hydroxylamine of the extract without hydroxylamine (Table 4). It is important to point out that the concentration of Fe in the extract with hydroxylamine is due to other soluble species of Fe, such as Fe associated with compounds of low molecular weight and that are not precipitated by the action of trichloroacetic acid.

The enrichment increased the concentration of inorganic Fe. Non-heme Fe can be found in foods of plant origin such as inorganic Fe(II) and/or Fe(III). The absorption of these iron species by the human organism is less than 5%. So, although bean sprouts cannot be considered as a plant source rich in non-heme iron, Fe(III)-EDTA enrichment may assist in increasing of Fe species available for absorption, which was able to increase in 75, 50 and 145% the concentration of total, soluble and inorganic Fe, respectively.

Se speciation in enriched sprouts

Optimization of the total Se extraction: Se species separation by extraction must be considered carefully in order to maintain the integrity of the species. To effectively release analytes from the matrix without alteration, two extractants, namely, deionized water and 1.0mol L⁻¹ NaOH, and different conditions, namely, heating in a thermostatic water bath and ultrasonic treatment, were studied, and the results are shown in Table 5.

Statistical tests were used to compare the total Se concentration obtained after acid digestion in a microwave oven by extraction using different extractants (deionized water or NaOH solution) or systems (water bath or ultrasound). The Student's *t*-test (95% confidence) and *p*-values (<0.05) did not show significant differences between the obtained results. A CPE procedure using deionized water and a thermostatic water bath was selected for Se speciation.

Table 5 Evaluation of the total selenium extraction method

Extractant	Extraction method	Concentration ($\mu\text{g g}^{-1}$) $\pm$ standard deviation (n=3)	
		After acid digestion	After extraction procedure
H ₂ O	Thermostatic water-bath	105 $\pm$ 2	93 $\pm$ 4
	Ultrasound-assisted	105 $\pm$ 2	92 $\pm$ 8
NaOH	Thermostatic water-bath	105 $\pm$ 2	97 $\pm$ 5
	Ultrasound-assisted	105 $\pm$ 2	94 $\pm$ 6

For total protein extractions, NaOH has been widely used, but it is not protein-specific.³¹ As the proposed method aims at speciation of organic and inorganic Se species, the use of NaOH extractant can be harsh mainly for inorganic Se species. Moreover, previous studies showed that Se(IV) can be determined by co-precipitation, after precipitation of metal hydroxides, such as iron(III), lanthanum, or magnesium hydroxides.¹² Thus, Se(IV) determination would not be possible by CPE with NaOH.

Preservation of Se species was also the reason for choosing a thermostatic water bath over ultrasonic treatment. The latter is known to cause cavitation, a physical process by which tiny gas bubbles are generated. Upon implosion, these bubbles behave as microreactors with temperatures and pressures reaching about 5000°C and 100atm, respectively. The power of ultrasonic waves has raised concerns for applications in chemical speciation, because it can result in the formation of radical species.³² In the determination of Se-methylselenocysteine and Se-methionine in Se-enriched plants (*Allium sativum* and *Brassica juncea*), the effect of the ultrasonic probe on the extraction using different extractants (namely, HCl, protease, and ammonium acetate buffer) was evaluated, and a lower amount of Se was extracted with HCl and ammonium acetate buffer when an ultrasonic probe was used. In the case of Se speciation, the application of an ultrasonic probe provided negligible species degradation.³³

Evaluation of pyrolysis and atomization temperatures

The heating program for Se determination by GF AAS was optimized in the presence of the reagents used for CPE. Pyrolysis and atomization temperature curves were obtained using analytical solutions of Se in Triton X-114 and dithizone, co-injected with the chemical modifier (Pd+Mg), and are shown in Figure 1. It was observed that Se species were thermally stable up to 900°C. At a pyrolysis temperature below 400°C, absorption could not be measured because of the high background signals, mainly due to the presence of surfactant and complexing agent. The optimum atomization temperature was 2100°C, because at lower temperatures double analytic signals were observed.

Figure 1 Pyrolysis (●) and atomization (▲) temperatures curves. (40µg L⁻¹) Se(IV), chemical modifier, Triton X-114 and dithizone.

Upon addition of the Pd/Mg mixture, a high pyrolysis temperature (900°C) was achieved, due to the formation of a non-volatile, solid solution between Se and Pd, which allowed the elimination/minimization of the matrix and surfactant and complexing agent. The chemical modifier is essential for Se determination by GF AAS because, in its absence and at temperatures above 130°C, Se volatilizes in the form of oxides (SeO and SeO₂).³⁴

Optimization of the Se speciation procedure

CPE is based on the formation of turbidity and the subsequent separation of hydrophilic and hydrophobic components.³⁵ Aqueous solutions of nonionic surfactant materials become cloudy when their temperature reaches the cloud point, and the analyte collapses with a small volume by a surface-active material.³⁶ This leads to a large preconcentration factor due to a very small volume of micellar phase binding the analyte that was dispersed in the original matrix.³⁷ Thus, CPE is not only a preconcentration process but also a separation

method, and has been applied for speciation of Se in trace levels.³⁸

The solution pH plays an important role in the formation of metal-chelate complexes, their chemical stability, and subsequent extraction. Only Se(IV) reacts with dithizone to form a hydrophobic complex, which is then trapped by surfactant micelles and separated from the aqueous phase.³⁹ The pH effect on the extraction of different Se species (Se(VI), Se(IV), and Se-methionine) was investigated, and the results are shown in Figure 2. The extraction efficiency (%) was calculated from the initial Se concentration with respect to that after CPE. Se(IV) extraction was higher at pH 2, which was used in subsequent studies. At pH values above 3, a significant decrease in the Se(VI) analytical signal was observed, due to dithizone instability under these conditions.²⁴ For the other Se species (namely, Se(VI) and Se-methionine), as expected, complex formation did not occur, and the CPE efficiency was zero, confirming the selective reactivity of dithizone toward Se(IV).

Figure 2 Effect of pH on complexation of Se(IV) (▲), Se(VI) (●) and SeMet (■) with dithizone and cloud point extraction efficiency.

Dithizone and Triton X-114 concentrations as well as equilibration temperature and incubation time are important parameters for CPE, which should be optimized to achieve complete reaction, easy phase separation, and efficient preconcentration.³⁷ However, several studies have been reported on the application of CPE using Triton X-114 and complexing agents such as dithizone for total Se determination and fractionation (organic and inorganic species).^{24,34,36-39} Hence, in this work, these parameters were not investigated, and the conditions established by Sun et al.²⁴ were applied.

The effect of HCl concentration on the reduction of Se(VI) to Se(IV) was examined, as shown in Figure 3. The HCl solution was added to the surfactant-poor phase obtained after the first CPE using only Triton X-114, and the mixture was heated, subjected to a second CPE in the presence of dithizone, and analyzed by GF AAS. The HCl can serve as a reducing agent for Se(VI) reduction, by the following chemical reaction: $\text{HSeO}_4^- + 3\text{H}^+ + 2\text{Cl}^- \rightleftharpoons \text{H}_2\text{SeO}_3 + \text{H}_2\text{O} + \text{Cl}_2$. Other reducing agents have also been proposed in the literature, such as TiCl₃, KI, thiourea, HBr, and a mixture of HCl and HBr.⁴⁰ Using only HCl, efficient Se(VI) reduction occurred at high HCl concentrations,⁴⁴ as shown in Figure 3, and the highest extraction efficiency was observed using a 6.0mol L⁻¹ HCl solution. However, 4.0mol L⁻¹ was chosen as the optimal HCl concentration, because at this value the extraction efficiencies were statistically indistinguishable (Student's *t*-test and 95% confidence).

After optimization of the conditions, the hydrophobic organic Se species were determined in the surfactant-rich phase after CPE in the absence of dithizone. The surfactant-poor phase containing hydrophilic organic and inorganic Se species was subjected to a

second CPE in the presence of dithizone for Se(VI) determination. For inorganic Se species determination, the surfactant-poor phase was first subjected to Se(VI) reduction by addition of HCl and heating, followed by CPE in the presence of dithizone. Se(VI) species were determined by subtracting the Se(IV) concentration from the total concentration of inorganic Se species. Moreover, the hydrophilic organic Se species were quantified by subtracting the Se concentration in the surfactant-poor phase obtained after CPE without dithizone from the total concentration of inorganic Se species.

Figure 3 Effect of HCl concentration on reduction of Se(VI) to Se(IV). CPE conditions: 1% v v⁻¹ Triton X-114 and 2mmol L⁻¹ dithizone.

Analysis of the enriched sprouts

Se enrichment of adzuki bean sprouts is an inexpensive strategy to improve the quality and nutritional value of food, and could also help combat nutritional deficiencies. The success of the enrichment is related not only to the increase in the total Se concentration, but also to the formation of bioaccessible and bioavailable species. According to the literature, organic Se species are more bioavailable and less toxic than the inorganic species,^{11–13} and Se(IV) is more toxic than Se(VI).⁴⁰ For the cultivation of Se-enriched sprouts, Se(VI) was used, which is absorbed by plant roots through the sulfate carriers and easily distributed to the aerial parts of the plant. In plant metabolism, Se(VI) is initially reduced by the catalysis of ATP sulfurylase to Se(IV), which is assimilated into organic compounds of Se.⁴¹ Thus, enriched adzuki bean sprouts can be safely used as dietary supplements, and can be considered as an appropriate alternative to reduce Se deficiency. The results demonstrating the applicability of the procedure to Se speciation in enriched adzuki bean sprouts are summarized in Table 6.

Table 6 Selenium speciation results in enriched adzuki beans sprouts

		Se concentration (µg g ⁻¹) ± standard deviation (n=3)
Total		105 ± 2
Inorganic species	Se(IV)	1.5 ± 0.1
	Se(VI)	4.2 ± 0.2
Organic species	Hydrophobic species	1.5 ± 0.1
	Hydrophilic species	96.2 ± 2.3

An estimated safe and adequate daily dietary intake of Se has been established as 10–200 µg per day, depending on age.¹¹ Considering the maximum value (200 µg per day) and total Se concentration (105 µg g⁻¹), only 1.9 g (dry mass) of enriched adzuki bean sprouts should be consumed per day. The sum of Se species concentrations corresponded to 99% of the total Se concentration. Chemical

speciation studies demonstrated that organic, mainly hydrophilic, Se species were the major constituents, corresponding to 93% of the total Se concentration. The inorganic Se species corresponded to 5.4% of the total Se concentration, with a percentage of Se(IV) (1.4%) lower than that of Se(VI) (4.0%). Because Se organic species are more bioavailable than inorganic Se species, a consumption of approximately 2.0 g of enriched adzuki bean sprouts corresponds to the maximum safe and adequate daily dietary intake of Se (200 µg per day).

Conclusion

The method applied for the Fe speciation in adzuki bean sprouts allowed to evaluate the enrichment efficiency of these agricultural products through a rapid procedure that uses the complexation of Fe(II) with antinutritional components present naturally in the plant matrix, which allowed to determine the concentrations of soluble and inorganic Fe. In the bean sprouts, it was verified that the enrichment increased the total, soluble, and inorganic Fe species concentrations, species that are potentially absorbed by the human organism, indicating the effectiveness of the proposed enrichment procedure.

An analytical strategy based on the combination of CPE as a preconcentration and separation method with GF AAS allowed the determination of organic (hydrophilic and hydrophobic) and inorganic (Se(IV) and Se(VI)) Se species in enriched adzuki bean sprouts. The developed methodology is simple and safe compared to other extraction procedures because it does not require exposure to organic compounds and includes a sustainable method of elemental preconcentration and speciation. The methodology enabled the identification and quantification of Se forms in the sprouts, showing that organic, mainly hydrophilic, Se species were predominant. Thus, because organic Se species are more bioavailable than inorganic Se species, enriched adzuki bean sprouts can be safely used as dietary supplements, and can be considered as an appropriate alternative to reduce Se deficiency.

Acknowledgements

Aline Pereira de Oliveira thanks the Fundação de Amparo à Pesquisa do Estado de São Paulo/FAPESP (2015/01128-6) for the fellowship provided. Juliana Naozuka thanks the FAPESP for financial support (2015/15510-0).

Compliance with ethics requirements

The author declares that they have no conflict of interest.

This article does not contain any studies with human or animal subjects.

References

- McGILL JA. Michigan: Japan and azuki beans. *Michigan Dry Bean Digest*. 1995;19:4–7.
- Sangronis E, Machado CJ. Influence of germination on the nutritional quality of *Phaseolus vulgaris* and *Cajanus cajan*. *LWT Food Sci Technol*. 2007;40(1):116–120.
- Yuan L, Wu L, Yang C, Lv Q. Effects of iron and zinc foliar applications on rice plants and their grain accumulation and grain nutritional quality. *J Sci Food Agr*. 2013;93(2):254–261.

4. Lintschinger J, Fuchs N, Moser J, et al. Selenium-enriched sprouts. A raw material for fortified cereal-based diets. *J Agric Food Chem*. 2000;48(11):5362–5368.
5. Wei Y, Shohag MJ, Ying F, et al. Effect of ferrous sulfate fortification in germinated brown rice on seed iron concentration and bioavailability. *Food Chem*. 2013;138(2–3):1952–1958.
6. Davidsson L, Ziegler E, Zeder C, et al. Sodium iron EDTA [NaFe (III) EDTA] as a food fortificant: erythrocyte incorporation of iron and apparent absorption of zinc, copper, calcium, and magnesium from a complementary food based on wheat and soy in healthy infants. *Am J Clin Nutr*. 2005;81(1):104–109.
7. Alfthan G, Xu GL, Tan WH, et al. Selenium supplementation of children in a selenium-deficient area in China. *Biol Trace Elem Res*. 2000;73(2):113–125.
8. Bravo L. Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutr Rev*. 1998;56:317–333.
9. Foster LH, Sumar S. Selenium in health and disease: a review. *Crit Rev Food Sci Nutr*. 1197;37(3):211–228.
10. Papp LV, Lu J, Holmgren A, et al. From selenium to selenoproteins: synthesis, identity, and their role in human health. *Antioxid Redox Signal*. 2007;9(7):775–806.
11. Chen S, Zhu S, Lu D. Solidified floating organic drop microextraction for speciation of selenium and its distribution in selenium-rich tea leaves and tea infusion by electrothermal vapourisation inductively coupled plasma mass spectrometry. *Food Chemistry*. 2015;169:156–161.
12. Tuzen M, Saygi KO, Soylak M. Separation and speciation of selenium in food and water samples by the combination of magnesium hydroxide coprecipitation–graphite furnace atomic absorption spectrometric determination. *Talanta*. 2007;71(1):424–429.
13. Rayman MP, Infante HG, Sargent M. Food-chain selenium and human health: spotlight on speciation. *Br J Nutr*. 2008;100(2):238–253.
14. Lynch SR. Interaction of iron with other nutrients. *Nutr Rev*. 1997;55(4):102–110.
15. Oliveira AP, Naozuka J. Chemical speciation of iron in different varieties of beans (*Phaseolus vulgaris* L.): cooking effects. *J Braz Chem Soc*. 2015;26(10):2144–2149.
16. Oliveira AP, Naozuka J. Effects of iron enrichment of adzuki bean (*vigna angularis*) sprouts on elemental translocation, concentrations of proteins, distribution of Fe–metalloproteins, and Fe bioaccessibility. *J Braz Chem Soc*. 2017;28(10).
17. Oliveira AP, Nomura CS, Naozuka J. Evaluation of selenium enrichment of adzuki bean (*Vigna angularis*) sprouts: Translocation, bioaccessibility and Se–protein speciation. *Microchemical Journal*. 2017;134:19–26.
18. Templeton DM, Ariese F, Cornelis R, et al. Guidelines for terms related to chemical speciation and fractionation of elements. Definitions, structural aspects, and methodological approaches (IUPAC Recommendations 2000). *Pure and Applied Chemistry*. 2000;72(8):1453–1470.
19. Machado ALDL, Barcelos MDP, Teixeira AHR, et al. Avaliação de componentes químicos em brotos de Fabaceae para o consumo humano. *Cienc Tecnol Aliment*. 2009;33(4):1071–1078.
20. Naozuka J, Vieira EC, Nascimento AN, et al. Elemental analysis of nuts and seeds by axially viewed ICP OES. *Food Chemistry*. 2011;124(4):1667–1672.
21. Carreira RL, Ornellas CBD, Morais HA, et al. Efeito da precipitação pelo ácido tricloroacético (TCA) e da ultrafiltração sobre o perfil peptídico de hidrolisados de caseína. *Ciênc Agrotec*. 2003;7(2):414–421.
22. Quinteros A, Farré R, Lagarda M. Optimization of iron speciation (soluble, ferrous and ferric) in beans, chickpeas and lentils. *Food Chemistry*. 2001;75(3):365–370.
23. Kosse JS, Yeung AC, Gil A, et al. A rapid method for iron determination in fortified foods. *Food Chemistry*. 2001;75(3):371–376.
24. Sun M, Liu G, Wu Q (2013) Speciation of organic and inorganic selenium in selenium-enriched rice by graphite furnace atomic absorption spectrometry after cloud point extraction. *Food Chem*. 2013;141(1):66–71.
25. Silva JCJ, Cadore S, Nobrega JA, et al. Dilute-and-shoot procedure for the determination of mineral constituents in vinegar samples by axially viewed inductively coupled plasma optical emission spectrometry (ICP OES). *Food Addit Contam*. 2007;24(2):130–139.
26. <https://www.fda.gov/food/foodscienceresearch/laboratorymethods/ucm2006954.htm>
27. Carreira RL, Ornellas CBD, Morais HA, et al. Efeito da precipitação pelo ácido tricloroacético (TCA) e da ultrafiltração sobre o perfil peptídico de hidrolisados de caseína. *Ciênc Agrotec*. 2003;7(2):414–421.
28. Hallberg L, Rossander L, Skander AB. Phytates and the inhibitory effect of bran on iron absorption in man. *American J Clin Nutr*. 1987;45(5):988–996.
29. Pallauf J, Rimbach G. Nutritional significance of phytic acid and phytase. *Arch Anim Nutr*. 1997;50(4):301–319.
30. Quirrenbach HR, Kanumfre F, Rosso ND, et al. Behaviour of phytic acid in the presence of iron(II) and iron(III). *Ciênc Tecnol Aliment*. 2009;9(1):24–32.
31. Chunhieng T, Pétritis K, Elfakir C, et al. Study of selenium distribution on the protein fractions of the Brazil nut, *Bertholletia excels*. *J Agr Food Chem*. 2004;52(13):4318–4322.
32. Santos HM, Capelo JL. Trends in ultrasonic-based equipment for analytical sample treatment. *Talanta*. 2007;73(5):795–802.
33. Montes-Bayón M, Molet MJD, González EB, et al. Evaluation of different sample extraction strategies for selenium determination in selenium-enriched plants (*Allium sativum* and *Brassica juncea*) and Se speciation by HPLC–ICP–MS. *Talanta*. 2006;68(4):1287–1293.
34. Correia PRM, Oliveira E, Oliveira PV. Simultaneous determination of manganese and selenium in serum by electrothermal atomic absorption spectrometry. *Talanta*. 2002;57(3):527–535.
35. Sounderajan S, Kumar GK, Udas AC. Cloud point extraction and electrothermal atomic absorption spectrometry of Se (IV)—3, 3'-diaminobenzidine for the estimation of trace amounts of Se (IV) and Se (VI) in environmental water samples and total selenium in animal and fish tissue samples. *J Hazard Mater*. 2010;75(1–3):666–672.
36. Güler N, Maden M, Bakirdere S, et al. Speciation of selenium in vitamin tablets using spectrofluorometry following cloud point extraction. *Food Chemistry*. 2011;129(4):1793–1799.
37. Escaleira LA, Santelli RE, Oliveira EP, et al. Preconcentration procedure for determining trace amounts of Ni, Cd, Pb and Cu in high-salinity waters after cloud-point extraction. *Int J Environ Anal Chem*. 2009;89(7):515–527.
38. Ghambariana M, Yaminia Y, Saleha A, et al. A 16 orthogonal array design for the optimization of cloud point extraction for selenium

- determination in environmental and biological samples by tungsten-modified tube electrothermal atomic absorption spectrometry. *Talanta*. 2009;78(3):970–976.
39. Soruraddin MH, Heydari R, Puladvand M, et al. A new spectrophotometric method for determination of selenium in cosmetic and pharmaceutical preparations after preconcentration with cloud point extraction. *Int J Anal Chem*. 2011;1–8.
40. Gallignani M, Valero M, Brunetto MR, et al. Sequential determination of Se(IV) and Se(VI) by flow injection–hydride generation–atomic absorption spectrometry with HCl:HBr microwave aided pre-reduction of Se(VI) to Se(IV). *Talanta*. 2000;52(6):1015–1024.
41. Pilon-Smits E, Quinn C. *Selenium metabolism in plants*. Berlin: Springer; 2010.