

# New nano-molecule kurumi - $C_{13}H_{20}BeLi_2SeSi$ / $C_{13}H_{19}BeLi_2SeSi$ , and raman spectroscopy using *ab initio*, hartree-fock method in the base set CC-pVTZ and 6-311G\*\* (3df, 3pd)

## Abstract

The work characterizes the Raman spectrum of the new nano-molecule  $C_{13}H_{20}BeLi_2SeSi$  /  $C_{13}H_{19}BeLi_2SeSi$ , nano-molecule Kurumi. Calculations obtained in the methods Restrict Hartree-Fock of the first principles (*ab initio*), on the set of basis used indicate that the simulated molecule  $C_{13}H_{20}BeLi_2SeSi$  /  $C_{13}H_{19}BeLi_2SeSi$  features the structure polar-apolar-polar predominant. The set of basis used that have are correlation-consistent polarized Triple-zeta (CC-pVTZ) and Pople's basis sets six gaussian functions in the shell, three double zeta Gaussian functions, Slater type orbitals with polarization function (6-311G\*\* (3df, 3pd)). In the CC-pVTZ base set, the charge density in relation to 6-311G\*\* (3df, 3pd) is 50% lower. The length of the molecule  $C_{13}H_{20}BeLi_2SeSi$  is of 15.799Å. The Raman spectrum was calculated indicating the characteristic of the nano-molecule and their frequency ( $cm^{-1}$ ) were obtained in the set of bases used. The highest for Raman scattering activities peaks are in the frequency 3,348 $cm^{-1}$  with 7.107609729 $\text{\AA}^4/\text{amu}$  and 2,163 $cm^{-1}$  with 8.902805583 $\text{\AA}^4/\text{amu}$ , for CC-pVTZ and 6-311G\*\* (3df, 3pd), respectively. As the bio-inorganic molecule  $C_{13}H_{20}BeLi_2SeSi$  is the basis for a new creation of a biomembrane, later calculations that challenge the current concepts of biomembrane should advance to such a purpose. The new nano-molecule Kurumi is well characterize computationally. As its scientific name 3-lithio-3-(6-{3-selena-8-beryllatricyclo[3.2.1.0.2,4]oct-6-en-2-yl}hexyl)-1-sila-2-lithacyclopropane.

**Keywords:** biomembrane, CC-pVTZ, hartree-fock method, nano-molecule Kurumi, raman spectroscopy, 6-311G\*\* (3df, 3pd)

Volume 8 Issue 1 - 2019

Ricardo Gobato,<sup>1</sup> Marcia Regina Risso Gobato,<sup>2</sup> Alireza Heidari,<sup>3</sup> Abhijit Mitra<sup>4</sup>

<sup>1</sup>Laboratory of Biophysics and Molecular Modeling Genesis, State Secretariat of Education of Parana, Brazil

<sup>2</sup>Green Land Landscaping and Gardening, Seedling Growth Laboratory, Brazil

<sup>3</sup>Faculty of Chemistry, California South University, USA

<sup>4</sup>Department of Marine Science, University of Calcutta, India

**Correspondence:** Ricardo Gobato, Laboratory of Biophysics and Molecular Modeling Genesis, State Secretariat of Education of Parana, Bela Vista do Paraíso, Parana, 86130-000, Brazil, Email ricardogato@seed.pr.gov.br

**Received:** January 05, 2019 | **Published:** January 21, 2019

## Introduction

The structure of the  $C_{13}H_{20}BeLi_2SeSi$  is a bio-inorganic seed molecule for a biomembrane genesis that defies the current concepts of a protective mantle structure of a cell such as biomenbrane<sup>1-3</sup> to date is promising, challenging. Leaving to the Biochemists their experimental synthesis. Structures of a liquid crystal such as a new membrane may occur, micelles.<sup>1</sup>

A large investigative study in the literature was carried out, but was not found of the structure studied here. There is an absence of a referential of the theme, finding only one work in (Gobato et al.).<sup>1</sup> The present study is an extension of the work already begun in the creation of *ab initio*, computational methods of quantum chemistry, to prove the possible structure of the bio-inorganic nano-molecule genesis,  $C_{13}H_{20}BeLi_2SeSi$ . Their characterization by well-known *ab initio* methods of extreme precision is essential for their future experimental achievement. Therefore, the *ab initio* study using RHF (Restrict Hartree-Fock method)<sup>4-13</sup> and the set of basis used is CC-pVTZ<sup>10-14</sup> and 6-311G\*\* (3df, 3pd).<sup>7,14-37</sup>

The calculation of the Raman spectrum of the nano-molecule Genesis has the purpose of deepening the knowledge and characterization of this. The work characterizes the Raman spectrum of the nano-molecule  $C_{13}H_{20}BeLi_2SeSi$ . Calculations obtained in the *ab initio* RHF (Restrict Hartree-Fock method).<sup>4-13</sup> The set of basis used that have are CC-pVTZ<sup>10-14</sup> and 6-311G\*\* (3df, 3pd).<sup>7,14-18</sup>

## Materials and methods

### Hartree-fock methods

The Hartree-Fock self-consistent method<sup>4-13</sup> is based on the one-electron approximation in which the motion of each electron in the effective field of all the other electrons is governed by a one-particle Schrodinger equation. The Hartree-Fock approximation takes into account of the correlation arising due to the electrons of the same spin, however, the motion of the electrons of the opposite spin remains uncorrelated in this approximation. The methods beyond self-consistent field methods, which treat the phenomenon associated with the many-electron system properly, are known as the electron correlation methods. One of the approaches to electron correlation is the Møller-Plesset (MP)<sup>4-18</sup> perturbation theory in which the Hartree-Fock energy is improved by obtaining a perturbation expansion for the correlation energy.<sup>4,5</sup> However, MP calculations are not variational and can produce an energy value below the true energy.<sup>4,5</sup>

The exchange-correlation energy is expressed, at least formally, as a functional of the resulting electron density distribution, and the electronic states are solved for self-consistently as in the Hartree-Fock approximation.<sup>4-13</sup>

The full Hartree-Fock equations are given by

$$\epsilon_i \sigma_i(\mathbf{r}) = \left( -\frac{1}{2} \nabla^2 + V_{\text{ion}}(\mathbf{r}) \right) \sigma_i(\mathbf{r}) + \sum_j \int d\mathbf{r}' \frac{\sigma_j(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} \sigma_i(\mathbf{r}) - \sum_{\alpha\beta} \int d\mathbf{r}' \frac{\sigma_{\alpha}^*(\mathbf{r}') \sigma_{\beta}(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} \sigma_j(\mathbf{r}). \quad (1)$$

The minimal basis sets are not flexible enough for accurate representation of orbitals. Using multiple functions to represent each orbital, have the double-zeta basis set allows us to treat each orbital separately when we conduct the Hartree-Fock calculation:

$$\Phi_{2s}(\mathbf{r}) = \sum_{2s} \overset{STO}{O}(\mathbf{r}, \mathbf{r}_1) + d \sum_{2s} \overset{STO}{O}(\mathbf{r}, \mathbf{r}_2). \quad (2)$$

In Equation (2)  $2s$  atomic orbital expressed as the sum of two STOs (Slater type orbitals).<sup>4,5</sup> The two equations are the same except for the value of  $\xi$  which accounts for how large the orbital is. The constant  $d$  determines how much each STO will count towards the final orbital.<sup>4,5</sup>

The vast literature associated with these methods suggests that the following is a plausible hierarchy:

$$HF \ll MP2 < CISD < CCSD < CCSD(T) < FCI. \quad (3)$$

The extremes of 'best', FCI, and 'worst', HF, are irrefutable, but the intermediate methods are less clear and depend on the type of chemical problem being addressed. The use of HF<sup>4-13</sup> in the case of FCI was due to the computational cost.

### Hardware and software

**For calculations a computer models was used:** Intel® Core™ i3-3220 CPU @ 3.3 GHz x 4 processors,<sup>38</sup> Memory DDR3 4 GB, HD SATA WDC WD7500 AZEK-00RKA0 750.1GB and DVD-RAM SATA GH24NS9 ATAPI, Graphics Intel® Ivy Bridge.<sup>1</sup>

The *ab initio* calculations have been performed to study the equilibrium configuration of  $C_{13}H_{20}BeLi_2SeSi$  molecule using the GAMESS.<sup>14,15</sup> The set of programs Mercury 3.8,<sup>39</sup> Avogadro<sup>40,41</sup> are the advanced semantic chemical editor, visualization, and analysis platform and GAMESS<sup>14,15</sup> is a computational chemistry software program and stands for General Atomic and Molecular Electronic Structure System<sup>14,15</sup> set of programs. For calculations of computational dynamics, the Ubuntu Linux version 16.10 system was used.<sup>42</sup> The Graphic was edited in Origin software,<sup>43</sup> for comparison of the spectra obtained in the set of bases used.

## Results

Our results take place from the findings of the molecular dynamics of bio-inorganic nano-molecule  $C_{13}H_{20}BeLi_2SeSi$ . The results were summarized and shown in Figures (1) and (2), Tables (1) and (2).


### Nano-molecule Kurumi

The Figure 1 representation of the molecular structure of  $C_{13}H_{20}BeLi_2SeSi / C_{13}H_{19}BeLi_2SeSi$ , Nano-molecule Kurumi, 3-lithio-3-(6-{3-selena-8-beryllatricyclo[3.2.1.0<sup>2,4</sup>]oct-6-en-2-yl}hexyl)-1-sila-2-lithacyclopropane.

### Raman spectrum

In Figure (2) we have representation of the Raman spectrum of  $C_{13}H_{20}BeLi_2SeSi$ -Nano-molecule Kurumi, 3-lithio-3-(6-{3-selena-8-beryllatricyclo[3.2.1.0<sup>2,4</sup>]oct-6-en-2-yl}hexyl)-1-sila-2-lithacyclopropane. Frequency ( $cm^{-1}$ ) for Raman scattering activities (SR,  $\text{\AA}^4/\text{amu}$ ) -using computer programs GAMESS, in the *ab initio* for the RHF method, in sets of basis RHF/CC-pVTZ in left and 6-311G\*\* (3df, 3pd) in right, obtained using computer software GAMESS. The Graphic edited in Origin software, for comparison of the spectra obtained in the set of bases used.

<sup>1</sup>Creative Commons, (CC BY 4.0), <https://creativecommons.org/licenses/by/4.0/>. "Ivy Bridge", [https://pt.wikipedia.org/wiki/Ivy\\_Bridge](https://pt.wikipedia.org/wiki/Ivy_Bridge), Available in: August 31, 2018.



**Figure 1** Representation of the molecular structure of  $C_{13}H_{19}BeLi_2SeSi$ , Nano-molecule Kurumi, 3-lithio-3-(6-{3-selena-8-beryllatricyclo[3.2.1.0<sup>2,4</sup>]oct-6-en-2-yl}hexyl)-1-sila-2-lithacyclopropane, obtained through computer via *ab initio* calculation method RHF/CC-pVTZ.

### RHF/CC-pVTZ

The Table 1 contains the frequencies ( $cm^{-1}$ ) for Raman scattering activities (SR,  $\text{\AA}^4/\text{amu}$ ) of the Nano-molecule Kurumi, 3-lithio-3-(6-{3-selena-8-beryllatricyclo[3.2.1.0<sup>2,4</sup>]oct-6-en-2-yl}hexyl)-1-sila-2-lithacyclopropane via *ab initio* methods, set base RHF/CC-pVTZ for the infrared spectrum.

The highest for Raman scattering activities peaks are in the frequency  $708cm^{-1}$  with  $8.448529382\text{\AA}^4/\text{amu}$ ,  $732cm^{-1}$  with  $6.968083358\text{\AA}^4/\text{amu}$ ,  $1,180cm^{-1}$  with  $8.947657695\text{\AA}^4/\text{amu}$ ,  $2,180cm^{-1}$  with  $7.796950601\text{\AA}^4/\text{amu}$  and  $3,348cm^{-1}$  with  $7.107609729\text{\AA}^4/\text{amu}$  for RHF/CC-pVTZ.

### RHF/6-311G\*\*(3df,3pd)

The Table 2 contains the frequencies ( $cm^{-1}$ ) for Raman scattering activities (SR,  $\text{\AA}^4/\text{amu}$ ) of the  $C_{13}H_{20}BeLi_2SeSi$  molecule, of the Nano-molecule Kurumi, via *ab initio* methods, set base RHF/6-311G\*\* (3df,3pd) for the infrared spectrum.

The highest for Raman scattering activities peaks are in the frequency  $602cm^{-1}$  with  $7.279393384\text{\AA}^4/\text{amu}$ ,  $707cm^{-1}$  with  $7.719121469\text{\AA}^4/\text{amu}$ ,  $2,079cm^{-1}$  with  $6.915053417\text{\AA}^4/\text{amu}$ ,  $2,163cm^{-1}$  with  $8.902805583\text{\AA}^4/\text{amu}$  and  $3,094cm^{-1}$  with  $6.213759835\text{\AA}^4/\text{amu}$  for RHF/6-311G\*\* (3df, 3pd),

## Discussion

The calculations made so far admit a seed molecule at this stage of the quantum calculations of the arrangement of the elements we have chosen, obtaining a highly reactive molecule with the shape polar-apolar-polar. Its structure has polarity at its ends, having the characteristic polar-apolar-polar. Even using a simple base set the polar-apolar-polar characteristic is predominant. The set of bases used that have the best compatible, more precise results are CC-pVTZ and 6-311G(3df, 3pd). In the CC-pVTZ base set, the charge density in relation to 6-311G(3df, 3pd) is 50% lower. The structure of the bio-inorganic seed molecule for a biomembrane genesis that challenge the current concepts of a protective mantle structure of a cell such as biomembrane to date is promising, challenging. Leaving to the biochemists their experimental synthesis.

The study evolves to construct a biomembrane from the seed molecule. The calculations default already performed admit a hydrophobic and hydrophilic molecule, the basis of the formation of a micelle and or biomembrane, as the default template. But kurumi molecule is polar-apolar-polar characteristic is predominant, ie. a hydrophilic-hydrophobic-hydrophilic molecule, thus presenting another standard for the construction of a biomembrane, which is even more innovative because a bio-inorganic molecule. Going


beyond imagination, the most innovative and challenging proposal of the work advances the construction of a structure compatible with the formation of a “new DNA”, based now on the seed molecule.

Our discussion takes place from the findings of the molecular dynamics of bio-inorganic nano-molecule  $C_{13}H_{20}BeLi_2SeSi$ . The results were summarized and shown in Figure 1 & Figure 2, Table 1 & Table 2.

The Figure (1) representation of the molecular structure of  $C_{13}H_{19}BeLi_2SeSi$ , Nano-molecule Kurumi, 3-lithio-3-(6-{3-

seleno-8-beryllatricyclo[3.2.1.0<sup>2,4</sup>]oct-6-en-2-yl}hexyl)-1-sila-2-lithacyclopropane, obtained through computer via *ab initio* calculation method RHF/CC-pVTZ.

The Figure (2) represents of the Raman spectrum of  $C_{13}H_{20}BeLi_2SeSi$  - Frequency ( $cm^{-1}$ ) for Raman scattering activities (SR,  $\text{\AA}^4/\text{amu}$ ) - using computer programs GAMESS, in the *ab initio* for the Restrict Hartree-Fock method, in sets of basis CC-pVTZ and 6-311G\*\* (3df, 3pd), obtained using computer software GAMESS. Graphic edited in origin software, for comparison of the spectra obtained in the set of bases used.



**Figure 2** Representation of the Raman spectrum of  $C_{13}H_{20}BeLi_2SeSi$  - Frequency ( $cm^{-1}$ ) for Raman scattering activities (SR,  $\text{\AA}^4/\text{amu}$ ) - using computer programs GAMESS, in the *ab initio* for the RHF method, in sets of basis RHF/CC-pVTZ in left and 6-311G\*\* (3df, 3pd) in right.

**Table 1** Table contains the frequencies ( $cm^{-1}$ ) for Raman scattering activities (SR,  $\text{\AA}^4/\text{amu}$ ) of the  $C_{13}H_{20}BeLi_2SeSi$  molecule via *ab initio* methods, set base RHF/CC-pVTZ for the infrared spectrum

| $\nu(cm^{-1})$ | SR( $\text{\AA}^4/\text{amu}$ ) | $\nu(cm^{-1})$ | SR( $\text{\AA}^4/\text{amu}$ ) | $\nu(cm^{-1})$ | SR( $\text{\AA}^4/\text{amu}$ ) | $\nu(cm^{-1})$ | SR( $\text{\AA}^4/\text{amu}$ ) |
|----------------|---------------------------------|----------------|---------------------------------|----------------|---------------------------------|----------------|---------------------------------|
| 490            | 2.182596                        | 1050           | 1.851075                        | 1932           | 1.885078                        | 3094           | 6.21376                         |
| 595            | 3.420969                        | 1057           | 1.707548                        | 1939           | 3.535241                        | 3101           | 3.519756                        |
| 602            | 7.279393                        | 1092           | 1.388367                        | 2072           | 1.463697                        | 3108           | 1.427854                        |
| 609            | 1.764814                        | 1099           | 1.118466                        | 2079           | 6.915053                        | 3115           | 1.154649                        |
| 630            | 1.192582                        | 1106           | 1.109801                        | 2086           | 4.240445                        | 3122           | 1.346301                        |
| 637            | 1.038282                        | 1113           | 3.899045                        | 2093           | 1.062849                        | 3129           | 2.315457                        |
| 693            | 1.755585                        | 1120           | 1.974696                        | 2156           | 2.115886                        | 3136           | 1.096041                        |
| 700            | 2.135401                        | 1148           | 1.021344                        | 2163           | 8.902806                        | 3150           | 1.711815                        |
| 707            | 7.719121                        | 1155           | 3.520772                        | 2170           | 2.324113                        | 3262           | 3.23569                         |
| 714            | 2.723003                        | 1162           | 6.018009                        | 2583           | 1.551622                        | 3269           | 3.008913                        |
| 721            | 1.36844                         | 1169           | 3.541951                        | 2590           | 5.920065                        | 3276           | 5.031301                        |
| 728            | 2.834444                        | 1176           | 2.474677                        | 2597           | 3.47486                         | 3283           | 3.414571                        |
| 735            | 5.297166                        | 1183           | 1.335813                        | 2604           | 2.774907                        | 3290           | 2.106915                        |
| 742            | 1.255979                        | 1218           | 1.504666                        | 3045           | 3.454342                        | 3297           | 4.656712                        |
| 756            | 1.287419                        | 1225           | 1.55933                         | 3052           | 1.894141                        | 3304           | 5.694054                        |
| 931            | 1.12492                         | 1603           | 1.285718                        | 3073           | 1.020319                        | 3311           | 1.308788                        |
| 1015           | 2.295633                        | 1610           | 1.928284                        | 3080           | 2.52559                         | -              | -                               |
| 1043           | 1.507847                        | 1617           | 1.257005                        | 3087           | 2.277247                        | -              | -                               |

**Table 2** Table contains the frequencies ( $cm^{-1}$ ) for Raman scattering activities (SR,  $\text{\AA}^4/\text{amu}$ ) of the  $C_{13}H_{20}BeLi_2SeSi$  molecule via *ab initio* methods, set base RHF/6-311G\*\* (3df, 3pd) for the infrared spectrum

| $\nu(\text{cm}^{-1})$ | SR( $\text{\AA}^4/\text{amu}$ ) | $\nu(\text{cm}^{-1})$ | SR( $\text{\AA}^4/\text{amu}$ ) | $\nu(\text{cm}^{-1})$ | SR( $\text{\AA}^4/\text{amu}$ ) | $\nu(\text{cm}^{-1})$ | SR( $\text{\AA}^4/\text{amu}$ ) |
|-----------------------|---------------------------------|-----------------------|---------------------------------|-----------------------|---------------------------------|-----------------------|---------------------------------|
| 490                   | 2.182596                        | 1050                  | 1.851075                        | 1932                  | 1.885078                        | 3094                  | 6.21376                         |
| 595                   | 3.420969                        | 1057                  | 1.707548                        | 1939                  | 3.535241                        | 3101                  | 3.519756                        |
| 602                   | 7.279393                        | 1092                  | 1.388367                        | 2072                  | 1.463697                        | 3108                  | 1.427854                        |
| 609                   | 1.764814                        | 1099                  | 1.118466                        | 2079                  | 6.915053                        | 3115                  | 1.154649                        |
| 630                   | 1.192582                        | 1106                  | 1.109801                        | 2086                  | 4.240445                        | 3122                  | 1.346301                        |
| 637                   | 1.038282                        | 1113                  | 3.899045                        | 2093                  | 1.062849                        | 3129                  | 2.315457                        |
| 693                   | 1.755585                        | 1120                  | 1.974696                        | 2156                  | 2.115886                        | 3136                  | 1.096041                        |
| 700                   | 2.135401                        | 1148                  | 1.021344                        | 2163                  | 8.902806                        | 3150                  | 1.711815                        |
| 707                   | 7.719121                        | 1155                  | 3.520772                        | 2170                  | 2.324113                        | 3262                  | 3.23569                         |
| 714                   | 2.723003                        | 1162                  | 6.018009                        | 2583                  | 1.551622                        | 3269                  | 3.008913                        |
| 721                   | 1.36844                         | 1169                  | 3.541951                        | 2590                  | 5.920065                        | 3276                  | 5.031301                        |
| 728                   | 2.834444                        | 1176                  | 2.474677                        | 2597                  | 3.47486                         | 3283                  | 3.414571                        |
| 735                   | 5.297166                        | 1183                  | 1.335813                        | 2604                  | 2.774907                        | 3290                  | 2.106915                        |
| 742                   | 1.255979                        | 1218                  | 1.504666                        | 3045                  | 3.454342                        | 3297                  | 4.656712                        |
| 756                   | 1.287419                        | 1225                  | 1.55933                         | 3052                  | 1.894141                        | 3304                  | 5.694054                        |
| 931                   | 1.12492                         | 1603                  | 1.285718                        | 3073                  | 1.020319                        | 3311                  | 1.308788                        |
| 1015                  | 2.295633                        | 1610                  | 1.928284                        | 3080                  | 2.52559                         | -                     | -                               |
| 1043                  | 1.507847                        | 1617                  | 1.257005                        | 3087                  | 2.277247                        | -                     | -                               |

The Table 1 & Table 2 present the Raman spectrum frequencies for CC-pVTZ and 6-311G\*\* (3df, 3pd), respectively, for Raman scattering activities (SR,  $\text{\AA}^4/\text{amu}$ ),  $SR > 1$ .

The highest for Raman scattering activities peaks are in the frequency  $3,348\text{cm}^{-1}$  with  $7.107609729\text{\AA}^4/\text{amu}$  and  $2,163\text{cm}^{-1}$  with  $8.902805583\text{\AA}^4/\text{amu}$ , for CC-pVTZ and 6-311G\*\* (3df, 3pd), respectively. It presents “fingerprint” between the intermediate frequency intervals presented in Table 1 & Table 2.

Calculations obtained in the *ab initio* RHF method, on the set of basis used, indicate that the simulated molecule,  $C_{13}H_{20}BeLi_2SeSi$ , is acceptable by quantum chemistry. Its structure has polarity at its ends, having the characteristic polar-apolar-polar.

The 6-311G\*\* (3df, 3pd) set of basis exhibits the characteristic of the central chain, with a small density of negative charges, near the ends of the Carbons of this. In the CC-pVTZ base set, the charge density in relation to 6-311G (3df, 3pd) is 50% lower. It is characterized infrared spectrum of the molecule  $C_{13}H_{20}BeLi_2SeSi$ , for absorbance and transmittance, in Hartree-Fock method in the set of bases CC-pVTZ and 6-311G (3df, 3pd). The dipole moments CC-pTZV are 3.69% bigger than 6-311G\*\* (3df, 3pd).

The new nano-molecule Kurumi is well characterize computationally. As its scientific name 3-lithio-3-(6-{3-selena-8-beryllatricyclo[3.2.1.0<sup>2,4</sup>]oct-6-en-2-yl} hexyl)-1-sila-2-lithacyclopropane. Getting Kurumi name, which means boy in Tupi-Guarani language, which are indigenous inhabitants of southern Brazil.

## Limitations

Our study has so far been limited to computational simulation via quantum chemistry, an applied theory. Our results and calculations are compatible and with the theory of quantum chemistry, but their physical experimental verification depends on advanced techniques for their synthesis, obtaining laboratory for experimental biochemical.

## Conclusion

The highest for Raman scattering activities peaks are in the frequency  $3,348\text{cm}^{-1}$  with  $7.107609729\text{\AA}^4/\text{amu}$  and  $2,163\text{cm}^{-1}$  with  $8.902805583\text{\AA}^4/\text{amu}$ , for CC-pVTZ and 6-311G\*\* (3df, 3pd), respectively. The Raman spectrum was calculated, indicating the characteristic of the bioinorganic nano-molecule genesis. Now the challenge is to build the basic structure of the bio-inorganic membrane. From the unimaginable, going where our mind can take us and build a new DNA, that nanomolecule. Characterized its infrared spectrum and Raman. Quantically calculated, accepted by quantum chemistry parameters, with *ab initio* methods, in the bases CC-pVTZ and 6-311G\*\* (3df, 3pd). An experimental challenge to chemists. The new bio-inorganic nano-molecule  $C_{13}H_{20}BeLi_2SeSi$  /  $C_{13}H_{19}BeLi_2SeSi$ , is well characterize computationally. As its scientific name 3-lithio-3-(6-{3-selena-8-beryllatricyclo[3.2.1.0<sup>2,4</sup>]oct-6-en-2-yl} hexyl)-1-sila-2-lithacyclopropane, getting Kurumi name, which means boy in Tupi-Guarani language, which are indigenous inhabitants of southern Brazil. As the bio-inorganic nano-molecule Kurumi is the basis for a new creation of a biomembrane, later calculations that challenge the current concepts of biomembrane should advance to such a purpose.

## Acknowledgments

Prof. Dr. Lauro Figueroa Valverde, University Autonomous of Campeche (Faculty of Chemical-Biological Sciences), Calle Avenida Agustín Melgar s/n, Buenavista, 24039 Campeche, Camp., Mexico, for their contribution.

## Conflicts of interest

The author declares that there is no conflicts of interest.

## References

1. R Gobato, A Heidari, A Mitra. The Creation of  $C_{13}H_{20}BeLi_2SeSi$ . The Proposal of a Bio-Inorganic Molecule, Using *Ab Initio* Methods for The Genesis of a Nano Membrane. *Arc Org Inorg Chem Sci*. 2018;3(4):377–386.
2. AL Frischknecht, LJD Frink. *Biomembrane frontiers: nanostructures, models, and the design of life*. Humana Press: USA; 2009.
3. Dennis Chapman. *Biomembrane Structure and Function*. Topics in Molecular and Structural Biology. 1983.
4. IN Levine. *Quantum Chemistry*. Pearson Education (Singapore) Pte. Ltd., India. 2003.
5. A Szabo, NS Ostlund. *Modern Quantum Chemistry*. Dover Publications: New York; 1989.
6. P Hohenberg, W Kohn. Inhomogeneous electron gas. *Phys Rev*. 1964;(136):B864–B871.
7. W Kohn, LJ Sham. Self-consistent equations including exchange and correlation effects. *Phys Rev*. 1965;(140):A1133.
8. JP Perdew, M Ernzerhof, K Burke. Rationale for mixing exact exchange with density functional approximations. *J Chem Phys*. 1996;105(22):9982–9985.
9. TH Dunning Jr. Gaussian basis sets for use in correlated molecular calculations, The atoms boron through neon and hydrogen. *J Chem Phys*. 1989;(90):1007–1023.
10. RA Kendall, TH Dunning Jr, RJ Harrison. Electron affinities of the first-row atoms revisited. Systematic basis sets and wave function. *J Chem Phys*. 1992;(96):6796–6806.
11. DE Woon, TH Dunning Jr. Gaussian-basis sets for use in correlated molecular calculations. The atoms aluminum through argon. *J Chem Phys*. 1993;(98):1358–1371.
12. KA Peterson, DE Woon, TH Dunning Jr. Benchmark calculations with correlated molecular wave functions. The classical barrier height of the  $H+H_2 \rightarrow H_2+H$  reaction. *J Chem Phys*. 1994;100(100).
13. AK Wilson, T van Mourik, TH Dunning Jr. Gaussian basis sets for use in Correlated Molecular Calculations. Sextuple zeta correlation consistent basis sets for boron through neon. *J Mol Struct (Theochem)*. 1996;(388):339–349.
14. MS Gordon, Kim K Baldrige, Jerry A Boatz, et al. General atomic and molecular electronic structure system (GAMESS). *J Comput Chem*. 1993;14(11):1347–1363.
15. MS Gordon, MW Schmidt. *Advances in electronic structure theory: GAMESS a decade later. Theory and Applications of Computational Chemistry: the first forty years*. CE Dykstra, et al. editors. Elsevier: USA; 2005:1167–1189.
16. A Schaefer, H Horn, R Ahlrichs. Fully optimized contracted Gaussian-basis sets for atoms Li to Kr. *J Chem Phys*. 1992;97:2571–2577.
17. A Schaefer, C Huber, R Ahlrichs. Fully optimized contracted Gaussian-basis sets of triple zeta valence quality for atoms Li to Kr. *J Chem Phys*. 1994;100:5829.
18. F Weigend, R Ahlrichs. Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy. *Phys Chem Chem Phys*. 2005;7(18):3297–3305.
19. R Gobato. Study of the molecular geometry of Caramboxin toxin found in star flower (*Averrhoa carambola* L.). *Parana J Sci Edu*. 2017;3(1):1–9.
20. R Gobato, A Gobato, DFG Fedrigo. Molecular electrostatic potential of the main monoterpenoids compounds found in oil Lemon Tahiti - (*Citrus Latifolia* Var Tahiti). *Parana J Sci Edu*. 2015;1(1):1–10.
21. R Gobato, DFG Fedrigo, A Gobato. Allocryptopine, Berberine, Chelerythrine, Copsitine, Dihydrosanguinarine, Protopine and Sanguinarine. Molecular geometry of the main alkaloids found in the seeds of *Argemone Mexicana* Linn. *Parana J Sci Edu*. 2015;1(2):7–16.
22. R Gobato, A Heidari. Infrared Spectrum and Sites of Action of Sanguinarine by Molecular Mechanics and *ab initio* Methods. *International Journal of Atmospheric and Oceanic Sciences*. 2018;2(1):1–9.
23. R Gobato, DFG Fedrigo, A Gobato. Molecular geometry of alkaloids present in seeds of mexican prickly poppy. *Quantitative Biology*. 2015.
24. R Gobato, A Gobato, DFG Fedrigo. Study of the molecular electrostatic potential of D-Pinitol an active hypoglycemic principle found in Spring flower Three Marys (*Bougainvillea species*) in the Mm+ method. *Parana J Sci Edu*. 2016;2(4):1–9.
25. R Gobato, A Heidari. Calculations Using Quantum Chemistry for Inorganic Molecule Simulation  $BeLi_2SeSi$ . *Science Journal of Analytical Chemistry*. 2017;5(5):76–85.
26. R Gobato, A Heidari. Calculations Using Quantum Chemistry for Inorganic Molecule Simulation  $BeLi_2SeSi$ . *Science Journal of Analytical Chemistry*. 2017;5(5):76–85. 25
27. R Gobato, A Heidari. Molecular Mechanics and Quantum Chemical Study on Sites of Action of Sanguinarine Using Vibrational Spectroscopy Based on Molecular Mechanics and Quantum Chemical Calculations. *Malaysian Journal of Chemistry*. 2018;20(1):1–23.
28. A Heidari, R Gobato. A Novel Approach to Reduce Toxicities and to Improve Bioavailabilities of DNA/RNA of Human Cancer Cells—Containing Cocaine (Coke), Lysergide (Lysergic Acid Diethyl Amide or LSD),  $\Delta^9$ -Tetrahydrocannabinol (THC) [(–)-trans- $\Delta^9$ -Tetrahydrocannabinol], Theobromine (Xanthose), Caffèine, Aspartame (APM) (NutraSweet) and Zidovudine (ZDV) [Azidothymidine (AZT)] as Anti-Cancer Nano Drugs by Coassembly of Dual Anti-Cancer Nano Drugs to Inhibit DNA/RNA of Human Cancer Cells Drug Resistance. *Parana Journal of Science and Education*. 2018;4(6):1–17.
29. A Heidari, R Gobato. Ultraviolet Photoelectron Spectroscopy (UPS) and Ultraviolet-Visible (UV-Vis) Spectroscopy Comparative Study on Malignant and Benign Human Cancer Cells and Tissues with the Passage of Time under Synchrotron Radiation. *Parana Journal of Science and Education*. 2018;4(6):18–33.
30. R Gobato, A Heidari. Using the Quantum Chemistry for Genesis of a Nano Biomembrane with a Combination of the Elements Be, Li, Se, Si, C and H. *J Nanomed Res*. 2018;7(4):241–252.
31. R Gobato. Study of the molecular geometry of Caramboxin toxin found in star flower (*Averrhoa carambola* L.). *Parana J Sci Edu*. 2017;3(1):1–9. 19

32. R Gobato, A Gobato, DFG Fedrigo. Molecular electrostatic potential of the main monoterpenoids compounds found in oil Lemon Tahiti – (*Citrus Latifolia* Var Tahiti). *Parana J Sci Educ.* 2015;1(1):1–10.
33. R Gobato, DFG Fedrigo, A Gobato. Allocryptopine, Berberine, Chelerythrine, Copsitine, Dihydrosanguinarine, Protopine and Sanguinarine. Molecular geometry of the main alkaloids found in the seeds of *Argemone Mexicana* Linn. *Parana J Sci Edu.* 2015;1(2):7–16. 21
34. R Gobato, A Heidari. Infrared Spectrum and Sites of Action of Sanguinarine by Molecular Mechanics and *ab initio* Methods. *International Journal of Atmospheric and Oceanic Sciences.* 2018;2(1):1–9. 22
35. R Gobato, DFG Fedrigo, A Gobato. Molecular geometry of alkaloids present in seeds of mexican prickly poppy. *Quantitative Biology.* 2015. 23
36. R Gobato, A Gobato, DFG Fedrigo. Study of the molecular electrostatic potential of D–Pinitol an active hypoglycemic principle found in Spring flower Three Marys (*Bougainvillea species*) in the Mm+ method”. *Parana J Sci Educ.* 2016;2(4):1–9.
37. DFG Fedrigo, R Gobato, A Gobato. Avrocar: a real flying saucer. *Cornell University Library.* 2015.
38. [https://en.wikipedia.org/wiki/List\\_of\\_Intel\\_Core\\_i3\\_microprocessors](https://en.wikipedia.org/wiki/List_of_Intel_Core_i3_microprocessors)
39. *Mercury - crystal structure visualisation, exploration and analysis made easy.* The Cambridge Crystallographic Data Centre. 2012.
40. <http://avogadro.cc/>
41. D Marcus, DE Hanwell, DC Curtis, et al. Avogadro: An advanced semantic chemical editor, visualization, and analysis platform. *Journal of Cheminformatics.* 2012;4:17.
42. [https://en.wikipedia.org/wiki/Ubuntu\\_\(operating\\_system\)](https://en.wikipedia.org/wiki/Ubuntu_(operating_system))
43. [https://www.originlab.com/index.aspx?go=Products/Origin/2018b&pid=3289.](https://www.originlab.com/index.aspx?go=Products/Origin/2018b&pid=3289)