

Quality evaluation of noodles produced from wheat (*Triticum spp*) and almond seed (*Prunus dulcis*) flour blends

Abstract

In this study, the quality evaluation of noodles produced from wheat and almond seed composite flour was studied. A preliminary was carried out to ascertain the optimum acceptable level of almond seed flour in wheat flour noodle processing using 0-50%w/w, samples were subjected to sensory evaluation and the most acceptable samples was chosen. Hence, in the main study, level of almond seed flour was varied within the most acceptable level using 5, 10, 15 and 20% w/w inclusion giving rise to four samples. Functional properties, anti-nutrient content, proximate composition, essential amino acid profile, selected mineral content, textural and cooking analyses and sensory evaluation were done using standard methods. The bulk densities of the blend ranged from 0.801 to 0.884 g/ml, WAC, ranged from 2.20g/g to 3.60g/g and swelling index ranged from 6.27 to 2.27 as the proportion of almond seed flour increased from 0-20 %. The values of phytate, oxalate, and tannin range from 0.12 to 0.75g/100g, 0.03 to 0.92g/100g and 0.08 to 1.23g/100g respectively. An increase (9.99 % - 14.55 %) in protein content was observed in the flour blends with increased addition of almond flour, there were significant difference ($P < 0.05$) in mean samples. Lysine value ranged from 3.44 to 5.09%, Valine value ranged from 3.71 to 5.01% and leucine value ranged from 7.06 to 7.24% as the level of almond flour increases from 0 to 20% in the flour blends. Potassium (412.40 mg/100g), magnesium (167.00mg/100g) and calcium (66.55mg/100g) were predominant minerals in the formulated noodles while Iron and sodium were found in low concentration. The values of B₁, B₂, B₃, B₆, B₉, C, D and E ranged from 1.95 to 2.12mg/100g, 0.45 to 1.12 mg/100g, 0.72 to 0.68mg/100g, 1.25 to 2.12mg/100g, 12.09 to 22.02mg/100g, 19.55 to 22.12 and 0.21 to 2.12 mg/100g respectively as the proportion of almond seed increases from 0 to 20% in the blends used in noodle preparation. The results for length, porosity apparent density and weight for noodle sample from wheat flour (control) were 3.34cm, 0.52mm, 0.87 and 1.94 g respectively. The corresponding values for noodle from 95% wheat flour and 5% almond seed flour were 3.25cm, 0.50mm, 0.89 and 1.93g respectively. . Cooking time and cooking lose increased on addition of almond seed flour while percent rehydration reduced as the quantity of almond seed increased in the noodle samples. Blend formulation 80:20 W : A was mostly acceptable. This study therefore has presented a way of increasing consumption and utilization of almond seed with high nutritional content yet are underutilized increasing food security, providing more affordable noodle from local available food sources and further converts waste to wealth.

Keywords: noodle, almond seed and wheat

Volume 9 Issue 3 - 2021

Ogah, Richard Abah, Diana Ahure, Momoh Clement Owoicho, Mohammed Ikagu Yusufu
Federal University of Agriculture, Nigeria

Correspondence: Momoh Clement Owoicho, Department of Food Science and Technology, Federal University of Agriculture, Makurdi, Nigeria, Email Momohclement86@gmail.com

Received: May 01, 2021 | **Published:** August 10, 2021

Introduction

Noodles are quick cooking foods that are increasingly gaining acceptance in Nigeria and gradually displacing most traditional diets that serve as breakfasts and snacks due to convenience, improved income, nutritional values, status symbol and job demands especially among urban dwellers.¹ The popularity of noodles has been attributed to its sensory appeal, versatility of form and ease of preparation at consumers end, nutritional content and long shelf life.² Pasta extrusion is known to result into products where the starch is slowly digested and absorbed.³ The main ingredient in noodles is wheat and water mixed at appropriate concentrations.⁴ Wheat is not only deficient in protein and other micronutrients as with cereals in general; it is also a foreign exchange depleter as a result of importation from scarce resources due to unfavorable climate for large scale production in Nigeria.⁵ It will be therefore thought worthwhile to produce noodle from wheat and almond seeds which are abundant in Nigeria. Almond

is a cereal that is predominantly grown in all the ecological and dietary zones.⁶ Almond seeds are high in protein and other micronutrient.⁷ Almonds are 4% water, 22% carbohydrate, 21 % protein and 50% fats. It is a good source of minerals (Faiyazet al., 2007). One ounce of almond seeds is an excellent source of vitamin E, magnesium, copper, potassium, calcium, iron and B vitamins. Almond seeds are currently underutilized; it is used as agricultural will bete and increases postharvest loss in spite of its high potential as a raw material for the preparation of functional foods and nutraceuticals.⁸⁻¹⁰ Apart from the nutritional benefits of almond seeds, it has anti nutritional factors. Tannins and phenols are enzymes present naturally in almond which attacks nutrient absorption. Extrusion and thermal processing denatures the naturally occurring anti-nutritional factors and thus has been widely used in reduction of anti-nutritional factors in plant based food products.¹¹ In Nigeria where there is an increasing demand for ready-to-eat, convenient and quick cooking food as well as

malnutrition due to deficiency in protein calories contributing to more than half a million deaths of newborn and growing children,¹² Almond seeds can be utilized as a supplementary protein in wheat-based extruded foods to improve their nutritional quality. However, this will achieve the desired result as long as the sensory characteristics of the end product are not discernibly changed and the product is affordable to people at the bottom of the income pyramid. Incorporating almond seeds with wheat in noodle processing could result in low cost noodles with enhanced protein-energy, in addition almond will add reasonable amount of minerals which would serve as a potential solution to the consequences of hidden hunger in Nigeria. Although information on the use of wheat in pasta production has been reported, there is little information on incorporating almond seed flours for production of pasta despite the fact that these are indigenous food sources. They are produced in large quantities in Nigeria and are major sources of protein and micronutrients. This study thus, seeks to produce high protein-energy and more affordable noodles from wheat and almond see composite flours and to evaluate the chemical composition, cooking, textural characteristics and sensory attributes of the noodles with the aim of promoting utilization of almond seeds.

Materials and methods

Procurement of materials

Imported whole wheat flour and Almond seeds flour were purchased from Makurdi Modern Market, Benue State.

Table 1 Mixing proportions of wheat and almond seed flours

	Samples				
	A (Control)	B	C	D	E
Wheat (%)	100	95	90	85	80
Almond seed flour (%)	0	5	10	15	20




Figure 1 Flow chart for preparation of noodle from wheat and almond seed composite flour.

Source: Momoh et al.¹³

Equipment

A laboratory scale LSG65 twin screw extruder (Jianansaibaibainuo Technology LMT) with 15.5cm screw length, 10.01cm screw diameter and 0.8mm die diameter was used for extrusion cooking. Extruder, tray drier (M/s Balaji enterprise, Shaharanpur, India) and moisture analyser (GRD 60H₂O China) was used.

Sample preparation

Wheat flour and Almond seeds flour were purchased from the market and sieved through a 0.5mm size mesh and were packaged in Low density polyethylene bags prior to analyses. African yam bean flour was prepared as shown in Figure 1. The flour blend formulation is captured in Table 1. The flour blends were conditioned to 35% moisture content, mixed manually and extruded at 120°C through 0.8 mm die diameter of a laboratory scale twin screw extruder (Jianansaibaibainuo Technology LMT) with 15.5cm screw length, 10.01cm screw diameter. Extrusion of flour blends was done following the method described by Momoh et al.¹³ for noodle production . Two (2) kilograms of each composite flours were conditioned to 35% moisture content; and kept for 30mins. The composite flours were separately extruded at 120°C barrel temperature. The extruder was stabilized for each run with 2kg whole rice flour at steady state operation; samples were collected in open pans. The extrudate was allowed to cool and dried for 3hours in a tray drier at 60°C to a moisture content of 6-8%, sealed in low density polyethylene bags and store in a refrigerator for subsequent analysis. Samples for analyses were milled into flour.

Analytical methods

The proximate analysis, amino acid analysis and mineral elements (Na, Ca, K, Mg, and Fe) to determine how much of major or macro components which include Moisture, Ash, Fiber, Fat and Protein were determined according to standard methods AOAC, (2012). The carbohydrates were determined by difference using the formula: % carbohydrate = 100 -% protein + % ash + % crude fiber + % crude fat + % moisture.

Statistical analysis

Statistical Package for Social Sciences (SPSS) V23 computer software was used to analyze the data. Means and Standard deviation were calculated where appropriate. One way Analysis of variance (ANOVA) was used to determine the treatment that was different from others in the various parameters tested; differences were considered significant at 95% (p<0.05) significant level and 99% (p<0.01) significant level where mentioned.

Result and discussion

Functional properties of wheat and almond seed flour

The values for bulk density, foam capacity, water absorption capacity, oil absorption capacity, Swelling index and gelation temperature for wheat flour and almond seed composite flours are similar although with different ingredients to that reported by Ekop

(2006) and Eneche (2005) for composite flours produced from African yam beans and wheat flour. Wheat flour has the lowest value for foam capacity (10.50%), water absorption capacity (1.00g/g) and oil absorption capacity (0.98g/g) while swelling characteristic (7.25) was highest in wheat flour. This is probably attributed to the fact that protein characteristic of wheat and almond which are same at room temperature are believed to be largely responsible for functional properties such as foaming, emulsification, oil and water absorption capacity, while swelling characteristics and gelation temperatures are starch related. Foam capacity of composite flours ranged from 18.00 to 32.00%. There was a gradual increase in foam capacity with increasing addition of almond seed flour, this value is higher than those obtained for flour from boiled dates (1.98%). Physiochemical changes in proteins such as denaturation at high temperature may affect functional properties such as water and fat absorption capacities, protein solubility and product characteristics.¹⁴ The bulk density of the composite flours ranged from 0.80 to 0.88g/ml, with the highest value found in sample 80:20:W:A formulation, flour samples were significantly different ($P<0.05$) from each other. Higher value of bulk density (2.45 and 2.45g/ml) were reported by Agunbiade & Ojezele¹⁵ for breakfast cereals made from maize, sorghum, African yam beans and soybeans. Severtz & Desrositer reported bulk density ranges of 0.31 g/ml and 0.20 g/ml for roasted ground coffee and instant coffee powder respectively. The bulk densities suggest that the samples may require different package space and material. The less the bulk density, the more packaging space is required.¹⁵ Water absorption capacity (WAC) of the flour blends ranges from 1.00 to 2.20 g/g. The water absorption capacity increased with increase in almond seed flour inclusion. This may be due to the hygroscopic properties of almond seed flour, thus absorbing moisture on exposure to water. Similar values were recorded for breakfast cereal from blends of maize; African yam beans, defatted coconut cake and sorghum extract.¹ Values obtained varied significantly ($P<0.05$). Oil absorption capacity (OAC) of the flour blends vary in trend as those obtained for water absorption capacity. The values ranged from 0.91 to 1.4 g/g with the highest recorded for sample 80:20W: A formulation. There was significant difference ($P<0.05$) amongst samples mean. Oil absorption capacity also increased with increasing addition of almond seed flour. There is a gradual decrease in the values for swelling index with increasing almond seed flour. Mean values ranged from 7.25 to 6.21, with the highest value recorded for wheat flour. Sample means with respect to oil absorption capacity vary significantly ($P<0.05$). The Gelation Temperature of the flour blends ranged from 69.30 to 73.11°C. Gelation Temperature of the blends generally increased with increasing addition of almond seed flour. Extrusion cooking however reduces the energy required since it becomes pre-cooked. Gelation temperature might be associated with the relative ratio of amylase and amylopectin in the composite flour. Case et al. reported that waxy and regular maize flour gelatinize at 62-72°C, whereas high-amylose starches begin to swell below 100°C; temperatures of 120°C and above are required to fully disperse these starches.

Anti-nutrient content of wheat and almond seed composite flour

The anti-nutrient composition shows that wheat flour contained 0.11g/100g of phytic acid, 0.03g/100g Oxalate and 0.09g/100g tannin. The values of phytic oxalet, tannin and affloxaxin range from 0.11 to 0.75g/100g, 0.03 to 0.92g/100g, 0.08 to 1.23g/100g and 0.00 to 0.14g/100g respectively as the proportion of almond seed increases from 0 to 20% in the blends used in noodle preparation. The increase in these anti-nutrient proportion with respect to increasing levels of almond seed flour is probably due to higher levels of these anti-nutrients in almond seeds flour. The values obtained are in range to the recommended by FAO for anti-nutrient in flour samples and are

similar to those reported by pyakas, 2016 for anti-nutrient factors in selected plants.

Proximate composition of noodles prepared from wheat and almond composite flours

The results of proximate composition of noodles prepared from wheat (W), and almond seed composite flours showed that the protein contents ranged from 8.95 to 13.00%. The increase in the protein content could be attributed to high protein content for almond seed flour, these values are in agreement with those reported by Nwosu (2014), Ameh (2007), Eneche (2005), Obatolu *et al.* (2003) and Oshodi *et al.* (1997) are 23.2%, 21.0%, 22.3%, 20.5% and 23.0% respectively. A similar trend was observed by Igbabul *et al.* (2014) and Ekop (2006) with cookies produced from wheat, cocoyam and African yam beans composite flours and extruded cakes from wheat and African yam beans flour blends respectively. All noodle samples have dissimilar protein content and are significantly different ($P<0.05$). Sample 80:20:W:A had the highest protein content greater than that of wheat noodles (13.00%). The result obtained for ash content showed significant difference ($P<0.05$) among noodle samples. Similar values have been recorded by other researchers with 'Kokoro' a Nigerian snack made from Maize and African yam bean flour blends: 1.87, 2.17 and 2.42%. The relative increase in ash values of noodle samples may be attributed to the presence of almond seed used as part of the ingredients with relative high ash content. The crude fat content of noodle samples ranged from 1.24 to 5.06%, these values agrees with that reported by Priyanka *et al.* (2012) with extruded ready-to-eat snack produced using egg albumin cheese powder and African yam beans flour. Significant differences ($P<0.05$) were observed among samples. The increase in value obtained for fat content of noodle samples may be attributed to almond seed having high fat content. Low fat food products are less susceptible to rancidity and hence more shelf stable. Dietary fats that provide essential fatty acids (EFA) have been shown to enhance the taste and acceptability of foods, slow gastric emptying and intestinal motility, thereby prolonging satiety and facilitate the absorption of lipid-soluble vitamins. The lipid component also helps to determine the texture, flavor and aroma of foods. The crude fiber content of noodle samples ranged from 3.25 to 3.43%. Samples had dissimilar crude fiber and were significantly different ($P<0.05$) from each other. Crude fiber is very important in adding bulkiness to food and for prevention of some disease of the colon. Fiber is important for the removal of waste from the body thereby preventing constipation and many health disorders. The noodles therefore have a great potential for application as diabetic food because of the fiber source in their formulation that could be modified to generate a near perfect recipe for production of diabetic food/snacks. The carbohydrate content of noodle samples ranged from 75.10% to 66.53%. This implies that the noodles are good sources of energy needed for normal body metabolism. However, carbohydrate content of noodle samples decreased with addition of almond seed flour, Wheat noodle had the highest value. This could be because wheat flour used as the principal ingredient in the formulation has higher starch content than almond seed flour. All noodle samples were significantly different ($P<0.05$) from each other in carbohydrate content. The values for carbohydrate are in agreement with values reported for flakes produced by substitution of wheat flour with rice flour and rice bran.

Essential amino acid (%) composition of noodles prepared from wheat and almond seed composite flours

The results of essential amino acid content of noodles produced from wheat (W), and almond seed composite flours showed that the

predominant amino acid in the noodle are Lysine (5.09%), Leucine (7.85%), phenylalanine (3.30%), threonine (3.54%) and Valine (5.00%). The least in the noodle samples was tryptophan (1.00%). This probably is due to its low content in the raw materials used. The values of all the essential amino acids in the noodle samples compared favorably with other extruded legume based food products from soya beans and peas. The observed value of 5.09% for lysine content of 80:20:W:A formulation is high when compared with the range of values (1.20 to 3.90%) reported for most cereal based extruded food products. The values for essential amino acids observed for all noodle samples also compare favorably with Recommended Daily Intake for children (2-5years) which imply that these noodle samples are rich in amino acid and are nutritive. The values obtained compares favorably to the recommended (37.6%) and considered to be adequate for idea protein for children and 24% for adults.

Mineral (mg/100g) composition of noodle prepared from wheat and almond seed composite flours

The mineral content of noodles produced from wheat (W) and almond seed (A) composite flours was found to vary significantly ($P<0.05$) with increasing almond seed flour. This increase could be attributed to the relative high mineral content of almond flour. 80:20:W:A had the highest values for all minerals analysed. Potassium (411.10mg/100g), magnesium (166.50mg/100g) and calcium (66.50mg/100g) were predominant minerals in the formulated noodles while Iron and sodium were found in low concentration, this is similar to the report of Edem et al. (1990) that low concentration was found of iron and zinc with extruded food from African yam bean flour and fonio flour. Higher mineral content were found in sample containing higher quantity of almond seed flour which contributed to the relative high ash content in noodle samples. Uwaegbute et al. (2012) reported similar values for calcium content of African yam beans flour.

Vitamins composition of noodles prepared from wheat and almond seed composite flours .

The vitamin composition shows that wheat noodle contained 1.95mg/100g of B₁, 0.46mg/100g B₂, 0.73 mg/100g B₃, 1.26mg/100g B₆, 12.56mg/100g vitamin C, 19.00mg/100g vitamin D, 0.22 vitamin E, 0.02 vitamin B₉ and 0.14 mg/100g B₁₂. The values of B₁, B₂, B₃, B₆, C, D and E ranged from 1.95 to 2.16 mg/100g, 0.46 to 0.55mg/100g, 0.73 to 0.66mg/100g, 1.26 to 1.48mg/100g, 12.56 to 23.56mg/100g and 0.21 to 2.23 mg/100g respectively as the proportion of almond seed increases from 0 to 20% in the blends used in noodle preparation. These values are in agreement with that reported by Singapore, 2015: for chemical composition of extruded food from almond seed flours and maize. Momoh et al., 2018 further reported similar values for vitamin content of noodles produced from broken rice, rice bran and African yam beans composite flours. All noodle samples had dissimilar ($P<0.05$) minerals content and were significantly different at 5% significant level. The noodle prepared from 20% level of almond seed flour was significantly ($P<0.05$) higher in all minerals analyzed.

Textural properties and cooking quality of noodles prepared from wheat and almond composite flours

Wheat noodle (Control) had the lowest apparent density(1.94) and weight (0.87) while the highest was recorded for sample formulation 80:20: W:A (0.93 and 1.94g respectively), these results were similar to the study by Sajilata et al. (2006), who found that weight and porosity value of noodle sample added with banana flour were higher than

those of wheat noodles. It is highly vital that the structural integrity of noodles need to be maintained throughout the cooking process. High porosity is unacceptable as there can be high amount of solubilized starch present, which leads to cloudy boiling water and “sticky” mouth feel with lower tolerance. There was significant difference ($P<0.05$) in means with respect to porosity and weight of noodles. The decrease in length and porosity on addition of almond seed flour in this study is in agreement with previous findings on quality evaluation of ‘flat’ noodle noodles. Textural characteristics of noodles are also a critical characteristic, which determines consumer acceptance of the product.

Sensory properties of noodles prepared from wheat and almond composite flours

Sensory perception varied significantly ($P<0.05$) among samples. Values obtained for taste showed a low level of acceptance of the noodles with increasing level of almond seed flour. However, no significant difference ($P>0.05$) was obtained for appearance (color) of the samples with change in quantity (%) of almond seed flour. Higher sensory scores for flavor with increasing almond seed flour in the sample shows the undesirable choky flavor usually associated with almond seeds is negligible in the noodle. This may be attributed to the method used during almond flour production and extrusion, which might have off-set the choky flavor. There was no significant difference ($P>0.05$) for flavor of the samples with change in quantity (%) of almond seed flour. All noodle samples were generally accepted for all attributes evaluated as none scored below the minimum acceptable rating of 5 on a 9 point hedonic scale. This indicates a high level of acceptance of the noodle prepared from wheat and almond seed composite flours, thereby improving the nutritional contents and as well as increasing the utilization of almond seed which was added as part of the ingredient.

Conclusion

Extrusion of wheat, and almond seed composite flour produced acceptable noodles with significantly increased protein and ash contents in blend ratio 80:20:W:A. Crude fat and crude fibre were similarly increased. The noodles could contribute significant amounts of mineral with respect to FAO/WHO/UNU recommended daily intakes for Mg (130mg/day), Na (1000mg/day), Ca (500mg/day), K (3000mg/day) and Fe (10mg/day). Functional properties of wheat and almond seed flour composite compared favorably with other legumes and cereals, hence it could find application in food industries. Length and porosity reduced as apparent density and weight increased on addition of almond seed flour. The sensory attributes showed that acceptable noodle can be produced from wheat supplemented with almond seed flour. Blend formulation 80:20 W:A was mostly acceptable. This study therefore presents a way of increasing consumption and utilization of almond seed: with low economic value, which are underutilized. Almond seeds could therefore serve as ingredients in food formulation to reduce malnutrition and hidden hunger among children and vulnerable groups in Nigeria, increase food security and further convert waste to wealth.¹⁶⁻¹⁰¹

Acknowledgments

The authors acknowledges the department of Food Science and Technology of the Federal Polytechnic Mubi, Adamawa State for making their extruder available.

Conflicts of interest

The authors declare that there is no conflict of interest.

References

1. Okafor GI, Usman GO. Production and evaluation of breakfast cereal from blends of African Yam beans (*sphenostylis stenocarpa*), Maize(*Zea mays*), and deffated coconut (*cocos Nucifera*). *Journal of food processing and preservation (in pres)*. 2015.
2. Anounye JC, Jigam AA, Ndaako GM. Effects of extrusion cooking on the nutrients and anitnutrients composition of pigeon pea and unripe plantain blends. *Journal of Applied Pharmaceutical Science*. 2012;2(5):58–162.
3. Wolever T, Whentit P, Dedoba WT. Function of Resistant starch in extruded foods produced from plantain and banana specie. *Journal of food and applied Bioscience*. 1986;2(1):1–5.
4. Chen Z, Sagis L, Legger A, et al. Evaluation of starch noodles made from three typical chinese sweet potato starches. *Journal of Food science*. 2003;67:3342–3347.
5. FAO. African experience in the improvement of post-harvest techniques. Synthesis based on the workshop held in Accra, Ghana, 4. 1994.
6. Erhabor P, Ojogho O. Demand analysis for rice in Nigeria. *Journal of food and Agric. Technology*. 2011;9(2):66–74.
7. Nnam NM. Nutrient composition and sensory evaluation of snacks produced from flour blends of some Nigerian indigenous food crops. *Nigerian Journal of Nutrittonal Sciences*. 2003;24(2):52–56.
8. Pitchaporn W, Channarong C, Sirithon S. Substitution of wheat flour with rice flour and rice bran in flake products: Effects on physical, chemical and antioxidant properties. *World applied science journal*. 2009;7(1):50.
9. Sudarat J, Srijesdaruk V, Harper WJ. Extraction of rice bran protein concentrate and its application in bread. *Journal of Science and Technology*. 2005;27(1):55–64.
10. Tahira R, Ata-ur-Rehman A, Butt NA. Characterization of rice bran oil. *Journal of Agricultural Resources*. 2007;45(3):225–230.
11. Olapade AA, Aworh OC. Chemical and nutritional evaluation of extruded complementary foods from blends of fonio (*DigitariaExilisStapf*) and cowpea (*VignaUnguiculata L. Walp*) flours. *International Journal of Food and Nutrition Science*. 2012;3(1):4–5.
12. Onyezilli P. Protein calorie deficiency: Control efforts. Abia. New times Purblisher; 1999:656.
13. Momoh CO, Abu JO, Yusufu MI. Quality Evaluation of Noodles Produced from Broken Rice (*Oryza sativa*), African Yam Bean (*Sphenostylis stenocarpa*) and Rice bran composite flours: A tool for Ameliorating PEM in Nigeria. *International Journal of Food Engineer- ing and Science*. 2020;10(4):31–38.
14. Potter D, Eriksson T, Evans, et al. Phylogeny and classification of Rosaceae. *Plant System Evolution*. 2007;266:5–43.
15. Agunbiade JO, Olankolun JO. Evaluation of some nutritional characteristics of the Indian almond (*Prunus amygdalus*) nut. *Pakistan Journal of Nutrition*. 2006;5:316–318.
16. AACC. American Association of Cereal Chemists, Approved methods of the AACC.10thedn. St. Paul, MN, AACC USA. 1995:1165.
17. Ahrens S, Venkatachalam M, Mistry AM, et al. Almond (*Prunus dulcis L.*) protein quality. *Plant Foods in Human Nutrition*. 2005;60:123–128.
18. Albert CM, Gaziano J, Willett WC, et al. Nut consumption and decreased risk of sudden cardiac death in the Physicians' Health Study. *Archives of Internal Medicine*. 2002;162:1382–1387.
19. Amarowicz R, Agnieszka T, Troszynska F, et al. Antioxidant activity of almond seed extract and its fractions. *Journal of Food Lipids*. 2005;12:344–358.
20. AOAC. Official Methods of Analysis. 20th edn. Association of Official Analytic Chemists. Washinton D.C. 2012:7765.
21. Arena A, Bisignano C. Immunomodulatory and the antiviral activities of almonds. *J I m let*. 2010;132(1-2):18–23.
22. Bailey LH, Bailey EZ. The staff of the Liberty Hyde Bailey Horotorium. Hortus third: A concise dictionary of Plants Cultivated in the United States and Canada. Macmillan, New York. 1976:234–241
23. Bansal P, Sannd R, Srikanth N, et al. Effect of a traditionally designed nutraceutical on the stress induced immunoglobulin changes at Antarctica. *African Journal of Biochemistry Res*. 2009;3:1084–1088.
24. Berry EM, Eisenberg S, Friedlander Y. Effects of diets which are rich in mono-unsaturated fatty acids on the plasma lipoproteins – the Jerusalem Nutrition Study II. Mono-unsaturated fatty acids Vs Carbohydrates. *American Journal of Clinical Nutrition*. 1992;56:394–403.
25. Berry SEE, Tydeman EA, Lewis HB, et al. Manipulation of lipid bioaccessibility of almond seeds influences postprandial lipemia in healthy human subjects. *American Journal*. 2008;23(7):2–24.
26. Berryman CE, Preston AG, Karmally W, et al. Effects of almond consumption on the reduction of LDL- Cholesterol: a discussion of potential mechanisms and future research directions. *Journal of Nutrition Review*. 2011;69:171–185.
27. Bes-Rastrollo M, Wedik NM, Martinez-Gonzalez MA, et al. Prospective study of nut consumption, long-term weight change, and obesity risk in women. *American Journal of Clinical Nutrition*. 2009;89:1–7.
28. Blomhoff R, Carlsen MH, Andersen LF, et al. Health benefits of nuts: potential role of antioxidants. *British Journal of Nutrition*. 2006;96:S52–S60.
29. Boca R, Trichopolou A, Lagiou P. Healthy traditional Mediterranean diet: an expression of culture, history and lifestyle. *Nutrition Reviews*. 1997;55:383–389.
30. Cassady BA, Hollis JH, Fulford AD, et al. Mastication of almonds: effects of lipid bioaccessibility, appetite and hormone response. *American Journal of Clinical Nutrition*. 2009;62:1231–1238.
31. Chen CY, Blumberg JB. *In vitro* activity of almond skin polyphenols for scavenging free radicals and inducing quinine reductase. *Journal of Agricultural and Food Chemistry*. 2008;56:4427–4434.
32. Chen CY, Milbury PE, Chung SK, et al. Effect of almond skin polyphenols and quercetin on human LDL and apolipoprotein B-100 oxidation and conformation. *Journal of Nutritional Biochemistry*. 2007;18:785–794.
33. Chen CY, Lapsley K, Blumberg J. A nutrition and health perspective on almonds. *Journal of the Science of Food and Agriculture*. 2006;86:2245–2250.
34. Chen CY, Milbury PE, Lapsley K, et al. Flavonoids from almond skins are bioavailable and actsynergistically with vitamins C and E to enhance hamster and human LDL metabolism to oxidation. *J Nutr*. 2005;135(6):1366–1373.
35. Daniel ZH. Domestication of plants in the old world: the origin and spread of cultivated plants in West Asia, Europe, and the Nile Valley. Oxford University Press; 2000:186.
36. Due A, Larsen TM, Hermansen K, et al. Role of cell walls in the bioaccessibility of lipids in almond seeds. *American Journal of Clinical Nutrition*. 2004;80:604–613.
37. Estruch R, Martinez-Gonzalez AM, Corella D, et al. Effects of a Mediterranean-style diet on cardiovascular risk factors. *Annals of Internal Medicine*. 2006;145:1–11.
38. European Parliament and Council. Corrigendum to Regulation (EC) No. 1924/2006 of the European Parliament and of the Council of 20th December 2006 on nutrition and health claims made on foods. Official Journal of the European Union: L404 30 December 2006, published 18 January 2007. 2007.

39. Fraser GE, Bennett HW, Jaceldo KB, et al. Effect on bodyweight of a free 76 kilojoule (320 calorie) daily supplement of almonds for six months. *Journal of the American College of Nutrition*. 2002;21:275–283.
40. Fraser GE, Sabate J, Beeson WL, et al. A possible protective effect of nut consumption on risk of coronary heart disease. *Archives of Internal Medicine*. 2008;152:1416–1424.
41. Frecka JM, Hollis JH, Mattes RD. Effects of appetite, BMI, food form and flavour on mastication: almonds as a test food. *European Journal of Clinical Nutrition*. 2008;62:1231–1238.
42. Frison-Norrie S, Sporns P. Identification and quantification of flavonol glycosides in almond seed coats by using MALDI-TOF.MS (2002). *Journal of Agricultural Food Chemistry*. 2002;50:2782–2787.
43. Gibson GR, Roberfroid MB. Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *Journal of Nutrition*. 1995;125(6):1401–1412.
44. Gopumadhavan S, Rafiz M, Venkataranganna MV, et al. Assessment of “Tentex royal” for sexual activity in an experimental model. *Indian Journal of Clinical Practice*. 2003;(13):10:23–26.
45. Griel AE, Kris-Etherton PM. Nuts: source of energy and macronutrients. *British Journal of Nutrition*. 2006;96:S29–S35.
46. Harper J, Janson R. Production of nutritious pre-cooked food in developing countries by low-cost extrusion technology. *Food Rev Int*. 2011;1:27–97.
47. Hollis J, Mattes R. Effect of chronic consumption of almonds on body weight in healthy humans. *British Journal of Nutrition*. 2007;98:651–656.
48. Holt SH, Miller JC, Petocz P, et al. A satiety index of common foods. *European Journal of Clinical Nutrition*. 1995;49:675–690.
49. Hu FB, Stampfer MJ. Nut consumption and risk of coronary heart disease: a review of epidemiological evidence. *Current Atherosclerosis Reports*. 1999;1:204–1209.
50. Hu FB, Stampfer MJ, Manson JE. Frequent nut consumption and risk of coronary heart disease in women: pro- spective cohort study. *British Medical Journal*. 1998;317:1341–1315.
51. Hyson DA, Schneeman BO, Davis PA. Almonds and almond oil have similar effects on plasma lipids and LDL oxidation in healthy men and women. *Journal of Nutrition*. 2002;132:703–707.
52. Isfahlan AJ, Mahmoodzadeh A, Hassanzadch A, et al. Anti-oxidant and anti radical activities of the phenolic extracts of the hulls and shells of the Iranian almond (*Prunus Amygdalus*). *Turkish Journal of Biology*. 2010;34:165–173.
53. Jaceldo-Siegl K, Sabate J, Rajaram S, et al. Long term almond supplementation without advice on food replacement induces favourable nutrient modification to the habitual diets of free-living individuals. *British Journal of Nutrition*. 2004;92:533–540.
54. Jenkins DJ, Kendall CW, Marchie A, et al. Almonds decrease postprandial glycemia, insulinemia and oxidative damage in healthy individuals. *Journal of Nutrition*. 2006;136(12):2987–2992.
55. Jenkins DJ, Kendall CW, Marchie A, et al. The effect of combining plant sterols, soy protein, viscous fibres and almonds in treating hypercholesterolemia. *Metabolism*. 2003;52:1478–1483.
56. Jenkins DJ, Kendall CW, Marchie A, et al. Direct comparison of a dietary portfolio of cholesterol-lowering foods with a statin in hypercholesterolemic participants. *American Journal of Clinical Nutrition*. 2005;81:380–387.
57. Jenkins DJ, Kendall CW, Marchie A, et al. Dose response of almonds on coronary heart disease risk factors: blood lipids, oxidised low density lipoproteins. 2002;100(1):34–36.
58. Jenkins DJA, Kendall CWC, Marchie A, et al. Almonds reduce the biomarkers of lipid per oxidation in older hyperlipidaemic subjects. *Journal of Nutrition*. 2008;138:908–913.
59. Jia X, Li N, Zhang W, et al. A pilot study on the effects of almond consumption on DNA damage and oxidative stress in smokers. *Nutrition and Cancer*. 2006;54:179–183.
60. Keith R. Collins wildlife trust guide trees: A photographic guide to the trees of Britan and Europe. London. Harper Collins; 1999:13–19.
61. Kelly JH, Sabate J. Nuts and coronary heart disease: an epidemiological perspective. *British Medical Journal*. 2006;96:S61–S67.
62. Knuops KT, de Groot LC, Krombout D, et al. Mediterranean diet, lifestyle factors and 10-year mortality in elderly European men and women: the HALE project. *Journal of the American Medical Association*. 2004;292:1433.
63. Knuops KT, Groot LC, Fidanza F, et al. Comparison of three different dietary scores in relation to 10-year mortality in elderly European subjects: the HALE project. *European Journal of Clinical Nutrition*. 2006;60:746–755.
64. Kris-Etherton PM, Zhao G, Binkoski AE, et al. The effects of nuts on coronary heart disease risk. *Nutrition Reviews*. 2001;59:103–111.
65. Kulkarni KS, Kastura SB, Mengi SA. Efficacy of the *Prunus amygdalus* (almonds) nuts in scopolamine induced amnesia in rats. *Indian Journal Pharmacol*. 2010;42:168–173.
66. Kushi LH, Folsom AR, Prineas RJ, et al. Dietary antioxidant vitamins and death from coronary heart disease in post-menopausal women. *New England Journal of Medicine*. 1996;334:1156–1162.
67. Liegner KB, Beck EM, Rosenberg A. Laetrile-induced agranulocytosis. *JAMA*. 1981;246(24):41–42.
68. Mandalari G, Neuno-palop C, Bigignano G, et al. Potential prebiotic properties of almond seeds. *J Applied and Environmental Microbiology*. 2008;74(14):4264–4270.
69. Mandalari G, Faulks RM, Rich GT, et al. Release of protein, lipidand vitamin E from almond seeds during digestion. *Journal of Agricultural and Food Chemistry*. 2008a;56:3409–3416.
70. Mandalari G, Nueno-Palop C, Bisignano G, et al. Potential prebiotic properties of almond (*Amygdalus communis* L.) seeds. *Applied and Environmental Microbiology*. 2008b;74:4264–4270.
71. Mark GD, Huxlen AJ. The New Roval Horticultural Society dictionary of gardening. London. Macmillan Press;1992.
72. Mattes RD. The energetics of nut consumption. *Asia Pacific Journal of Clinical Nutrition*. 2008;17:337–339.
73. Mattes RD, Kris-Etherton PM, Foster GD. Impact of peanuts and tree-nuts on bodyweight and healthy weight loss in adults. *Journal of Nutrition*. 2007;138:1741S–1745S.
74. Nuts, almonds. USDA National Nutrient Database for Standard Reference, Release 17. Agricultural Research Service, U.S. Department of Agriculture. *Journal of Clinical Nutrition*. 2004;88:922–929.
75. Phillips KM, Ruggio DM, Ashraf-khorassani M. The phytosterol composition of the nuts and seeds which are commonly consumed in the United States. *Journal of Agriculture and Food Chemistry*. 2005;53:9436–9445.
76. Phillips KM, Ruggio DM, Ashraf-Khorassani M. Phytosterol composition of nuts and seeds commonly consumed in the United States. *Journal of Agricultural and Food Chemistry*. 2005;53:9436–9445.
77. Phung OJ, Makanji SS, White CM, et al. Almonds have a neutral effect on the serum lipid profile. A meta analysis of random-ized trails. *Journal of the American Dietetic Association*. 2009;109(5):865–873.
78. Pinelo M, Rubilar M, Sineiro J, et al. Extraction of anti-oxidant phenolics from almond hulls (*Prunus amygdlaus*) and pine sawdust (*Pinus pinaster*). *Food Chemistry*. 2004;85:267–273.
79. Prasad K, Singh Y, Anil A. Effects of grinding methods on the quality of pusa 1121 rice flour. *Journal of Trop Agric and Food Sci*. 2012;40(2):195.

80. Puri A, Sahai R, Singh KL, et al. Immunostimulant activity of dry fruits and plant materials which are used in the Indian traditional medical system for mothers after child birth and invalids. *Journal Ethnopharmacol.* 2000;71:89–92.
81. Rajaram S, Sabate J. Nuts bodyweight and insulin resistance. *British Journal of Nutrition.* 2006;96:79S–86S.
82. Ros E, Mataix J. Fatty acid composition of nuts implications for cardiovascular health. *British Journal of Nutrition.* 2006;96(Suppl. 2):S29–S35.
83. Sabate J, Ang Y. Nuts and health outcomes: new epidemiological evidence. *American Journal of Clinical Nutrition.* 2009;89:1643S–1648S.
84. Sabate J, Haddadn E, Tanzman JS, et al. Serum lipid response to the graduated enrichment of a Step 1 diet with almonds: a randomized feeding trial. *American Journal of Clinical Nutrition.* 2003;77:1379–1381.
85. Salas-Salvado J, Fernandez-Ballart J, Ros E, et al. Effects of a Mediterranean-style diet supplemented with nuts on metabolic syndrome status. *Archives of Internal Medicine.* 2008a;168:2449–2458.
86. Salas-Salvado J, Garcia-Arellano A, Estruch R, et al. For the PREDIMED investigators Components of the Mediterranean-type food pattern and serum inflammatory markers among patients at high risk for cardiovascular disease. *European Journal of Clinical Nutrition.* 2008;62:651–659.
87. Sathe SK, Wolf WJ, Roux KH, et al. Biochemical characterisation of amandin, the major storage protein in almond (*Prunus dulcis* L.). *Journal of Agricultural and Food Chemistry.* 2002;9(8):23–34.
88. Scientific Advisory Committee on Nutrition. Nutrition well-being of the British population. TSO, London, UK. 2008.
89. Segura R, Javierre C, Lizarraga MA, et al. Other relevant components of nuts: phytosterols, folate and minerals. *British Journal of Nutrition.* 2006;96:S36–S44.
90. Shah KH, Patel JB, Shrma VJ, et al. Evaluation of the anti diabetic activity of *Prunus amygdalus* in streptozocin induced diabetic mice. *RJPBCS.* 2011;2(2):429–434.
91. Shargg TA, Albertson TE, Fisher C J Jr. Cyanide poisoning after bitter almond ingestion. *West Journal of Medicine.* 1982;136(1):65–69.
92. Soni M, Mohanthy PK, Jaliwala YA. Hepato protective activity of the fruits of *Prunus*. *International Journal of Pharma and Biosciences.* 2011;2(2):439–452.
93. Spiller GA, Jenkins DA, Bosello O, et al. Bruce nuts and plasma lipids: An almond –based diet lowers the LDL-C while it preserves the HDL-C. *J Am Coll Nutr.* 1998;17:285–290.
94. Spiller GA, Jenkins DJ, Cragen LN, et al. Effect of a diet high in monounsaturated fat from almonds on plasma cholesterol and lipoproteins. *Journal of the American College of Nutrition.* 1992;11:126–130.
95. Ternus ME, Lapsley K, Geiger CJ. Health benefits of tree nuts. In: Alasalver C, Shahidi F, editors. *Tree nuts: composition, phytochemicals and health effects.* 2009:37–64.
96. Toubro S, Martinussen T, Astrup A. Comparison of the effects on insulin resistance and glucose tolerance of 6-mo high-monounsaturated-fat, low-fat and control diets. *American Journal of Clinical Nutrition.* 2008;87:855–862.
97. Trichopolou A, Costacou T, Bamia C, et al. Adherence to a Mediterranean diet and survival in a Greek population. *New England Journal of Medicine.* 2003;348:2599–2608.
98. U.S. Department of Agriculture, Agricultural Research Service. USDA National Nutrient Database for Standard Reference, Release 21. Nutrient Data Laboratory Home Page.
99. U.S. Food and Drug Administration Centre for Food Safety and Applied Nutrition. Qualified health claims subject to enforcement discretion. Nuts & heart disease. Docket No. 02P- 0505. 2003.
100. Venkatachalam M, Teuber SS, Roux KHm et al. Effects of roasting, blanching, autoclaving and microwave heating on antigenicity of almond (*Prunus dulcis* L.) proteins. *Journal of Agricultural and Food Chemistry.* 2002;50:3544–3548.
101. Wien MA, Sabate JM, Ikl DN, et al. Almonds vs complex carbohydrates in a weight reduction program. *International Journal of Obesity and Related Metabolic Disorders.* 2003;27:1365–1372.