

Pattern of Childhood Cancers Presenting to the Paediatric Cancer Unit of a Tertiary Hospital in Kumasi, Ghana

Abstract

Introduction: Childhood cancers are increasing in incidence worldwide with geographic differences having an impact on the types of cancers seen. The main aim of this study was to assess the trends in yearly proportions and number of cases seen, to determine age and gender proportions of the types of cancers and types of childhood cancers diagnosed at the cancer unit from 2012 through 2014. The purpose of this study is to set the baseline framework for future research to improve knowledge, clinical care and advocacy programmes for the newly established paediatric cancer unit in KATH in Kumasi, Ghana.

Methods: A retrospective study was conducted to review secondary data from the paediatric cancer registry of all patients aged 0-15 years diagnosed with cancer by fine needle aspiration cytology and tissue biopsy at the paediatric cancer unit from 2012-2014. Demographic, anthropometric and clinical data stored in excel was exported to Stata and analysed. Simple descriptive statistics was used and presented in tables.

Results: Out of 309 cases, 187(60.5%, 95%CI: 54.8-66.0) were males. Burkitt's lymphoma (126, 40.8%) was the most common cancer seen followed by Acute Lymphoblastic Leukaemia (31, 10%), Wilm's Tumour (28, 9.1%), other Non-Hodgkin's Lymphoma (28, 9.1%), and retinoblastoma (22, 7.1%). Hodgkin's Lymphoma was the commonest cancer in the age group 10-15 years, Burkitt's lymphoma in the 5-10 year age group and Acute Lymphoblastic Leukaemia in the under 5 years group.

Conclusion and Recommendations: There is an observable trend in the progressive increment in the number of cancer cases presenting to the unit over the three years period. Except for the lymphomas, CNS cancers and osteosarcomas which were commoner in the 5-9 years age group, most of the other childhood cancers; leukaemias, renal cancers, soft tissue sarcomas and neuroblastomas were predominant in the 0-4 year age group. Except for a few rare cancers, majority of the common cancers affected more males than females in the paediatric population studied. The Male: Female ratio of all the cases was 1.5:1. The lymphomas were the commonest cancers diagnosed with Burkitt's lymphoma being the highest among the 5-9 years group. The acute lymphoblastic leukaemias were next to the lymphomas in KATH. Based on these findings, further studies need to be conducted into the specific clinical patterns particularly, the lymphomas and leukaemias for better case management.

Keywords: Pattern; Lymphoma; Leukaemia; Paediatric Cancer Unit; Ghana

Research Article

Volume 3 Issue 3 - 2015

Paintsil V¹, Dogbe J^{1,2*}, Blay Nguah S¹, Osei-Akoto A^{1,2}, Osei-Tutu L¹ and Hammond C¹

1 Directorate of Child Health- Komfo Anokye Teaching Hospital, Ghana

2 Kwame Nkrumah University of Science and Technology, Ghana

***Corresponding author:** Joslin Dogbe, Department of Child Health, Kwame Nkrumah University of Science and Technology and the Komfo Anokye Teaching Hospital, PO. Box: 1934, Kumasi, Ghana, Tel: +233244271701; Email: slimdogbe@gmail.com

Received: October 25, 2015 | **Published:** November 5, 2015

Abbreviations: CHRPE: Committee on Human Research Publications and Ethics; KATH: Komfo Anokye Teaching Hospital; KNUST: Kwame Nkrumah University of Science and Technology

Introduction

Cancers form one of the major causes of death in children aged 15 years or less. They differ markedly from adult cancers in their nature, distribution and prognosis. Worldwide, childhood cancer burden is increasing [1] and over 80% of children who develop cancers each year live in low and middle income countries [2]. While the outcome for cancers in the developed countries is good, the opposite is the case for patients in low and middle income countries. Early diagnosis of cancer is a fundamental

goal because it allows an opportunity for timely treatment while disease burden is in its earliest stages [3]. Consequently, prognosis may improve and a cure can be attained with minimal side or late effects [3]. Geographic differences in childhood cancer incidence rates suggest a genetic and environmental influence on disease susceptibility [2]. Socioeconomic status, malnutrition, exposure to viral infections and malaria are thought to play important roles in cancer pathology and clinical characteristics [4].

The patterns of childhood cancers in the United States of America and Europe are very similar, with leukaemia and central nervous system tumours accounting for over one-half of the new cases. In contrast, lymphoma is the most common prevailing cancer of this age group in Africa [5]. Although many studies have

been published on this in some African countries [6-8], reports on the patterns and incidence of childhood cancer in Ghana are very few [4,9]. There is paucity of data regarding childhood cancers in Ghana and no such publication exists on the cancer cases seen and treated at the Komfo Anokye Teaching Hospital (KATH) in Ghana. It is for this reason that this study was conducted with these objectives in mind, to assess the trends in yearly proportions of cancer cases being captured, to determine age and gender proportions of the types of cancers and types of childhood cancers diagnosed at the cancer unit from 2012 through 2014. The purpose of this study was to set the baseline framework for future research to improve knowledge, clinical care and advocacy programmes for the newly established paediatric cancer unit in KATH.

Materials and Methods

Study site

This study was conducted at the Paediatric Oncology Unit of KATH. KATH is a tertiary health institution accredited for training medical students and residents. It is located at Bantama which is near the central business centre of Kumasi. Kumasi is the regional capital of the Ashanti Region of Ghana and the second largest city in Ghana. The city has a population of 1,889,934 by 2009 according to the Ghana Statistical Service in 2010. KATH is a 1200 bed capacity hospital and operates as a referral centre for the whole northern sector of Ghana. The catchment area stretches beyond Ashanti region to the Western, Brong-Ahafo, Eastern, Central and the 3 Northern regions. The Paediatric Cancer Unit is one of the two paediatric cancer treatment centres in the country. It is located in the Child Health Directorate at KATH. Children with cancer up to 15 years are seen here.

Study type

The study involved a retrospective review of secondary data from the cancer registry of all patients from birth to 15 years, who were diagnosed with cancer between 1st January, 2012 and 31st December, 2014.

Study procedure

The cancer registry at the Paediatric Cancer Unit is in excel format. The registry has been in existence since 2008. Every patient who has a confirmed diagnosis of a cancer is entered on the excel database at the unit. Diagnosis for the various malignancies was confirmed using fine needle aspiration cytology or tissue biopsies for solid tumours. Bone marrow aspiration morphology was done for the diagnosis of haematological malignancies. Data entered in the registry includes demographic elements such as the name, age, gender, and address. Anthropometric data includes weight, height or length as well as laboratory, type of confirmed cancer, stage of cancer were entered daily, cleaned, stored and backed up. Data extracted from the cancer registry for this study included the demographic characteristics (age and gender) and the type of cancers which were classified based on the international classification of childhood cancer.

Statistical analysis

The data stored in excel format in the cancer registry was

exported to Stata and analysed. Simple descriptive statistics using, frequency distribution, percentages, means and standard deviation were used to describe the various variables and presented in tables.

Ethical considerations

Ethical approval was obtained from the Committee on Human Research Publication Ethics - KATH/KNUST.

Results

Trends in yearly proportions of cancer cases

A total of 309 patient data was extracted from the cancer registry to determine the proportion and trend of the number of cases being captured annually from 2012-2014. The yearly distribution is captured in Table 1 as follows.

Table1: Trends in Yearly Proportions of Cancer Cases.

Year	No. of Cases	Yearly Proportions (%)
2012	84	27.2
2013	92	29.8
2014	133	43
Total	309	100

Types of cancers by age distribution

Table 2 shows the types of cancer by age distribution from 0-15 years. The lymphomas; 166 were the commonest cancers seen across the age groups. The Leukaemias; 21, Wilm's tumour; 20, Soft tissue sarcomas; 11, Rhabdomyosarcoma; 9 and Neuroblastoma; 9 were documented more among the 0-4 year group while, the lymphomas; 96, Central Nervous System Cancers; 3, and Osteosarcomas; 2 were commoner in the 5-9 year group. Hepatoblastoma, though appears rare in the cancer registry among a few others, was seen with equal proportions in both 0-4 and 5-9 years age groups. The 10-15 year groups recorded mainly Lymphomas; 42, but less of the other cancers.

Types of cancers by gender distribution

There were a total of 187 (60.5%, 95%CI: 54.8-66.0) Males and 122 (39.5%, 95%CI: 34.0-45.2) Females, with a Male to Female ratio of 1.5:1. Most of the cancers are commoner in the males than the females by the proportions shown in the Table 3.

Types of childhood cancers diagnosed

Lymphomas were the commonest childhood malignancies seen with Burkitt's lymphoma being the commonest among them (Table 4). Burkitt's lymphoma accounted for 126 (40.8%) whiles Hodgkin's Lymphoma accounted for 12(3.9%) of the cases seen. Acute Lymphoblastic Leukaemias were the second commonest malignancies; 31(10.0%) Embryonal tumours such as Wilm's tumour and retinoblastoma were the 3rd and 4th commonest tumour with 28 (9.1%) and 22 (7.1%) respectively. The least common cancers seen were hepatoblastoma, nasopharyngeal carcinoma and germ cell tumours in the proportions of; 0.6%, 0.3% and 0.3% respectively.

Table 2: Types of Cancers by Age Distribution.

Type of Cancers	No. Cases	Age Groups (years)		
		0 to 4	5 to 9	10 to 15
Lymphoma				
Burkitt's Lymphoma	126	21	80	25
Non-Hodgkin's Lymphoma(NHL)	28	7	13	8
Hodgkin's Lymphoma	12	0	3	9
Leukaemia				
Acute Lymphoblastic Leukaemia (ALL)	31	14	13	4
Acute Myeloid Leukaemia (AML)	14	7	6	1
Wilms Tumours	28	20	7	1
Retinoblastoma	22	18	4	0
Soft Tissue Sarcomas				
Rhabdomyosarcoma	17	9	4	4
Other soft tissue sarcoma	4	2	1	1
Neuroblastoma	14	9	3	2
CNS Tumour	3	0	3	0
Osteosarcoma	3	1	2	0
Hepatoblastoma	2	1	1	0
Nasopharyngeal Carcinoma	1	0	1	0
Germ cell Tumours	1	1	0	0
Others	3	2	1	0
Total	309	112	142	55

Table 3: Types of Cancers by Gender Distribution.

Type of Cancer	Sex (n)	
	Male	Female
Lymphoma		
Burkitt's Lymphoma	84	42
Non-Hodgkin's Lymphoma(NHL)	15	13
Hodgkin's Lymphoma	8	4
Leukaemia		
Acute Lymphoblastic Leukaemia(ALL)	17	14
Acute Myeloid Leukaemia(AML)	8	6
Wilm's Tumour	19	9
Retinoblastoma	13	9
Soft Tissue Sarcomas		
Rhabdomyosarcoma	9	8
Other soft tissue sarcoma	1	3
Neuroblastoma	8	6
CNS Tumour	2	1
Osteosarcoma	1	2
Hepatoblastoma	1	1
Nasopharyngeal Carcinoma	0	1
Germ Cell Tumours	0	1
Others	1	2
TOTAL	187	122

Discussion

Trends in yearly proportions of cancer cases

The emerging threat of cancers in developing countries especially in the paediatric age group has received little attention. There is pre-occupation with infectious diseases such as malaria, tuberculosis and HIV which contributes to about 25% of the deaths in children under the age of one year. With successes in immunizations coverage, there will be a reduction in infectious diseases with attendant increases in prevalence of non-communicable diseases like cancers. For the three years under review, a total of 309 new cases of cancers were diagnosed in the unit and this could even be the tip of the iceberg. As shown in Table 1, the trend reveals an increment in the number of cases yearly from 27.2% to 29.8% then to 43.0%. It can be estimated from Table 1, that the average number of new cases seen per year over the three years under study, was approximately 103, which may not be too different from the 124 cases per year reported at Korle-Bu Teaching Hospital in Accra, the largest teaching hospital in Ghana [4].

Types of cancers by age distribution

Age is a palpable demographic indicator in the causation and the types of cancers. This study looked at children from 0-15 years with some exciting revelations. More specifically, the lymphomas were more dominant in the 5-9 year group. The peak age for patients presenting with Burkitt's lymphoma was similar to other studies [8,10].

Even more revealing is the finding that far more different cancers like the leukaemias, Wilm's tumour, soft tissue sarcomas, Rhabdomyosarcoma, and Neuroblastomas were all commoner in the 0-4 year group. In terms of prevalence of diagnosed cancers, the 5-9 year age group were at a higher risk but the 0-4 year group were at a higher risk for embryonal cancers and leukaemias. In our study most patients with acute lymphoblastic leukaemia were in the 0-4 year age group, similar to literature from the industrialized countries [11]. Patients with Hodgkin's Lymphoma seen in our study were mostly in the group of 10-15 years. This is comparable in the literature where adolescents are commonly affected [12].

Types of cancers by gender distribution

The study showed a male preponderance over the females with a male to female ratio of 1.5:1 (Table 3). This result is comparable with other studies found in other developing countries [5,10,13]. In the study by Segbefia et al at Korle-Bu Teaching Hospital, there was a male preponderance with a male to female ratio of 1.3:1. In resource poor countries, gender disparity in the diagnosis of paediatric cancers may be exaggerated mainly because of our cultural practices and economic factors. Males are normally given more medical attention than females [14]. The ratio of childhood cancer in boys as compared to girls increases with decrease in gross domestic product and with increasing infant mortality [15].

Types of childhood cancers diagnosed

Burkitt's lymphoma was the most prevalent tumour seen in this study as shown in Table 4. This is in agreement with other studies in developing countries which showed that Burkitt's lymphoma was the commonest cancer in the paediatric age group [8,10,16]. The high incidence of Burkitt's lymphoma in Equatorial

Africa is related to endemic malaria and Epstein Barr virus infections [14]. Older studies such as that by Welbeck et al. [9] have shown a higher frequency of around 67% for Burkitt's Lymphoma, while newer studies in the same hospital; Korle-Bu Teaching Hospital have lower results with 30.7% [4]. The prevalence found in KATH; 40.8% compares more favourably with the more recent Korle-Bu study. At this point we can only question whether the prevalence is indeed reducing? Could improve malaria control in Ghana with better diagnostic facilities be the reason for the perceived reduction? In patients with Burkitt's lymphoma that were seen, there was a shift from the mainly jaw masses classically seen in the tropics to those with both Jaw and abdominal masses. Acute Lymphoblastic Leukaemias were the second most common cancer seen in our study. With improved clinical and laboratory diagnosis of the leukaemias, there is a chance that the incidence might even go higher. Clinicians need to have a high index of suspicion and be aware of the early warning signs for leukaemias to diagnose it as it often mimics other common childhood illnesses like sickle cell disease and malaria [4]. The commonest embryonal tumours seen were Wilm's Tumour and Retinoblastoma, the third and fourth most commonly seen cancers in our study. Comparing our data with what is seen in Accra, Retinoblastoma was more common than Wilm's tumour. Central Nervous System tumours and neuroblastomas which are common in the industrialized countries were relatively uncommon here with 3 (1.0%) and 14 (4.5%) respectively. Under-diagnosis might be a reason for our relatively low numbers as the cost and availability of investigative imaging, often a pre-requisite for diagnosis, is too expensive for families to bear. Nasopharyngeal carcinoma which is seen in older children was seen in 1 (0.3%) patient with metastatic disease at presentation.

Conclusion and Recommendation

There is an observable trend in the progressive increment in the number of cancer cases presenting to the unit over the three years period. Except for the lymphomas, CNS cancers and osteosarcomas which were commoner in the 5-9 years age group, most of the other childhood cancers; leukaemias, renal cancers, soft tissue sarcomas and neuroblastomas were predominant in the 0-4 year age group. Except for a few rare cancers, majority of the common cancers affected more males than females in the paediatric population studied. The Male: Female ratio of all the cases was 1.5:1. The lymphomas were the commonest cancers diagnosed with Burkitt's lymphoma being the highest among the 5-9 years group. The acute lymphoblastic leukaemias were next to the lymphomas in KATH. Based on these findings, further studies need to be conducted into the specific clinical patterns of the individual cancers particularly the lymphomas and leukaemias for better case management.

References

1. Mellstedt H (2006) Cancer initiatives in developing countries. *Ann Oncol* 17(Suppl 8): viii24-viii31.
2. Howard S, Metzger M, Wilimas J, Quintana Y, Pui C, Robinson L, et al. (2008) Childhood cancer epidemiology in low-income countries. *Cancer* 112: 461-472.
3. Dang-Tan T, Franco EL (2007) Diagnosis delays in childhood cancer: Review. *Cancer* 110(4): 703-713.
4. Segbefia C, Renner L, Dei-Adomakoh Y, Welbeck J (2013) Changing

- pattern of childhood cancers at Korle Bu Teaching Hospital, Accra, Ghana. *Postgraduate Medical Journal of Ghana* 2(2): 65-67.
5. Haroun H, Mahfouz M, Elhaj A (2006) Patterns of childhood Cancer in children admitted to the institute of nuclear medicine, molecular biology and oncology (INMO), Wad Medani, Gezira State. *J Family Community Med* 13(2): 71-74.
 6. Ojesina A, Akang E, Ojemakinde K (2002) Decline in the frequency of Burkitt's lymphoma relative to other childhood malignancies in Ibadan, Nigeria. *Ann Trop Paediatr* 22(2): 159-163.
 7. Obioha F, Kaine W, Ikerionwu S, Obi G, Ulasi T (1989) The pattern of childhood malignancy in eastern Nigeria. *Ann Trop Paediatr* 9(4): 261-265.
 8. Shija J (1984) Paediatric malignant tumours seen in Dar es Salaam, Tanzania. *East Afr Med J* 61: 212-219.
 9. Welbeck J, Hesse A (1998) Pattern of childhood malignancy in Korle bu Teaching Hospital, Ghana. *West African Journal of Medicine* 17(2): 81-84.
 10. Agboola A, Adekanmbi F, Musa A, Sotimehin A, Deji-Agoola A, Shonubi A, et al. () Pattern of childhood malignant tumours in a teaching hospital in south-western Nigeria. *MJA* 190(1): 12-14.
 11. Li J, Thompson T, Miller J, Pollack L, Stewart S (2008) Cancer incidence among children and adolescents in the United States, 2001-2003. *Paediatrics* 121: e1470-e1477.
 12. Olu-Eddo AN, Omoti CE (2011) Hodgkin lymphoma: Clinicopathologic features in Benin City, Nigeria and update on its biology and classification. *Niger J Clin Pract* 14: 440-444.
 13. Ocheni S, Okafor CO, Emodi IJ, Ibegbulam OG, Olusina DB, et al. (2005) Spectrum of childhood malignancies in Enugu, Nigeria (1999-2004). *Afr J Med Sci* 34(4): 371-375.
 14. Yaris N, Mandiracioglu A, Buyukpamukcu M (2004) Childhood cancer in developing countries. *Pediatr Hematol Oncol* 21: 237-253.
 15. Pearce M, Parker L (2001) Childhood Cancer registrations in the developing world: still more boys than girls. *Int J Cancer* 91(3): 402-406.
 16. Mohammed A, Aliyu H (2009) Childhood cancers in a referral hospital in northern Nigeria. *Indian Journal of Medical and Paediatric Oncology* 30(3): 95-98.