

A Comparative Evaluation of Antibacterial Efficacy of 'Calcium Hydroxide Plus Points', 'Activ Points' and 'Combi Points' Against *Enterococcus Faecalis* in Endodontic Therapy: An *In Vitro* Analysis

Abstract

The success of endodontic therapy relies on complete eradication of microorganisms from the root canal system. However, there are some resistant strains such as *Enterococcus faecalis* which persist in the root canals, leading to endodontic failures. Various antimicrobial agents have been tested time to time for their efficiency against this pathogen. The purpose of this study was to compare the antimicrobial efficacy of calcium hydroxide releasing 'Calcium hydroxide plus points'; chlorhexidine releasing 'Activ points', chlorhexidine and calcium hydroxide releasing 'Combi points' against *Enterococcus faecalis*. By measuring the zone of inhibition, it was concluded that 'Calcium hydroxide plus points' were not effective, while both Activ points and Combi points showed results significantly better than Calcium hydroxide plus points. The results indicate the calcium hydroxide alone may not be effective when expecting *Enterococcus faecalis* in particular endodontic cases and must be replaced or adjuncted by the use of chlorhexidine.

Keywords: Activ points; Calcium hydroxide; Chlorhexidine; Combi points; *Enterococcus faecalis*

Research Article

Volume 2 Issue 5 - 2015

Harpreet Singh*, Mandeep Kaur, Jaidev Dhillon, Meenu Saini and Ashish Kaushal

Department of Conservative dentistry & Endodontics, Gian Sagar Dental College and Hospital, India

***Corresponding author:** Harpreet Singh, Associate Professor, Department of Conservative dentistry & Endodontics, Gian Sagar Dental College and Hospital, Ramnagar, Patiala-140401, Punjab, India, Tel: 91-9815493618; Email: drharpreetsingh.hs@gmail.com, hsgentledental@gmail.com

Received: May 14, 2015 | **Published:** June 25, 2015

Introduction

Bacteria and their products play a primary etiological role in the initiation and perpetuation of pulpo-periapical pathosis. The biological objective of root canal treatment is eradication of bacterial load from the root canal system. Despite the fact that all measures are being taken to achieve this goal, failures are still being reported [1]. The ability of certain microbes to form biofilms is one of the key factors in their pathogenesis of endodontic infection. These biofilms not only adhere strongly to the dentin but also penetrate deep into the dentinal tubules [2]. Also, there are certain bacteria which are, in general, resistant to many chemical reagents used in endodontic therapy.

Amongst these resistant bacteria, *Enterococcus faecalis* has been found to be a major isolate [3]. Its prevalence in failure root canal cases ranges from 24-77% [4] probably owing to its ability to establish monoinfection. It can also get converted into viable but non-cultivable state and can survive low pH, high salinity and high temperatures. It has also been observed that it encompasses Gene encoded antibiotic resistance. A number of anti-microbial irrigants and medicaments have been tried in endodontics since years, expecting to completely eliminate this bacteria from the root canals system such as calcium hydroxide, formocresol, camphorated paramonochlorophenol, chlorhexidine, etc [5-8]. Calcium hydroxide has bactericidal properties since it inhibits bacterial enzymes by means of hydroxyl ions of the bacteria's cytoplasmic membrane, generating the antibacterial effect [9]. Its high pH (around 12.5) has a damaging effect on cell membranes and protein structure [10].

Chlorhexidine (CHX) is a broad-spectrum antimicrobial agent that has been used as an effective intracanal medicament. It is a synthetic cationic bis-guanide, which consists of two symmetric 4-chlorophenyl rings and two biguanide groups, connected by a central hexamethylene chain [11]. Its effectiveness is because of altering the microbial cells osmotic equilibrium due to the interaction of the positive charge of the molecule and the negatively charged phosphate groups on microbial cell walls. CHX molecule penetrates into the bacteria due to increase in permeability of the cell wall [12]. It is a well-established fact that chlorhexidine alone or a combination of chlorhexidine and calcium hydroxide has very high efficiency against *E. faecalis* [13]. The evaluation of chlorhexidine as intracanal medicaments in the form of sustained releasing points (Activ points) alone or in combination with calcium hydroxide (Combi Points) has already been done [13,14]. The present *in vitro* study was undertaken to evaluate the antimicrobial efficacy of calcium hydroxide releasing 'Calcium hydroxide plus points' as compared 'Activ Points' and 'Combi points' against *E. faecalis* so as to analyse, if at all the sustained release of calcium hydroxide from the points has any better advantage.

Materials and Methods

The standard protocol of maintaining *E. faecalis* (ATCC 47077) on agar plates was used [13]. Thirty McConkey agar plates were inoculated with the bacteria. These agar plates were divided into three groups, depending on the test specimen used. 'Calcium Hydroxide plus points' (Roeko, Germany) were designated as Group I, chlorhexidine releasing 'Activ points' (Roeko,

Germany) designated as Group II ; and both calcium hydroxide and chlorhexidine releasing 'Combi points'(Roeko, Germany) were designated as Group III. The two test agents of same group were placed in each plate at a distance (Figure 1). Then the McKonkey agar plates were incubated for 24 hours at 37°C aerobically. Zones of inhibition indicating efficacy of the drug were measured after 48 hours using digital vernier caliper (Figure 2).

compared to group II, it was found that no significant difference was observed between the two groups.

Table I: Zone of inhibition (in mm) measured for all the three groups after 48 hours.

S. No	Group I	Group II	Group III
1	0	6.17	6.10
2	0	6.96	6.58
3	0	5.87	4.75
4	0	5.71	4.90
5	0	6.67	6.55
6	0	6.86	6.11
7	0	6.29	5.90
8	0	6.70	6.50
9	0	5.19	4.74
10	0	5.30	6.00
11	0	6.25	5.90
12	0	6.40	6.26
13	0	6.28	5.56
14	0	6.60	6.00
15	0	6.88	6.50
16	0	6.12	6.00
17	0	6.50	6.25
18	0	7.18	7.19
19	0	7.90	7.72
20	0	7.26	7.10




Figure 1: Test specimens placed in McKonkey agar plate.



Figure 2: Zone of inhibition measured by digital vernier caliper.

Observations and Results

Table 1 represents the zones of inhibition measured after 48 hours. While, no zone of inhibition was seen in samples for Group I (Figure 3). Zone of inhibition was seen in all the samples of Group II (Figure 4) and III (Figure 5). The mean zone of inhibition of each group was calculated and subjected to statistical analysis (paired T-test) for the mean zones of inhibition between the different groups. The significance value was set at $p < 0.05$ = significant. When compared, a statistically significant difference was found between Group I and II, with latter performing better than former ($p < 0.05$ = significant). Also, when Group III was compared to Group I, it performed and this was found to be statistically significant ($p < 0.05$ = significant). When group III was




Figure 3: No zone of inhibition seen in Group I sample.




Figure 4: Zone of inhibition seen in Group II sample.



Figure 5: Zone of inhibition seen in Group III sample.

Discussion

For the successful performance of the endodontic therapy, both the mechanical and biological principles of root canal preparation must be followed. While the mechanical principles are often strictly adhered to, the biological ones are, at times, overlooked. It is the presence of these remnant bacteria in the root canal system, which is commonly associated with the endodontic failure. Complex microbial flora consisting of cocci, rods, spirochetes, filaments and sometimes fungi are present in root canals of infected teeth [15]. One of the most resistant microbes found in root canal system is *E faecalis*. It is a Gram-positive, catalase negative, fermentative and non-spore-forming facultative anaerobic coccus. Its cells are ovoid with diameter ranging from 0.5 to 1µm. *Enterococci* possess several virulence factors that assist adherence to host cells and extracellular matrix and assist in tissue invasion. These factors include: aggregation substance; enterococcal surface proteins such as gelatinase; cytolysin toxin;

extracellular superoxide production; capsular polysaccharides and antibiotic resistance determinant [16,17]. One significant feature contributing to pathogenic potential of *E. faecalis* is its ability to adhere and infiltrate into dentine.

The analysis of the fact that which antimicrobial agents are effective against *E. faecalis* is of utmost importance. The use of calcium hydroxide in elimination of *E. faecalis* has been tested previously and results have not been very favorable. It has been postulated that *E. faecalis* is resistant to calcium hydroxide [18,19]. The results obtained in our study match with these previous tests. Siren et al. [20] in their study confirmed the effectiveness of chlorhexidine as an intracanal medicament in cases where *E. faecalis* was suspected. Similar conclusion has also been reported by Lui et al. [21] and Singh et al. [22]. Also, there have been studies earlier which suggest that combined use of chlorhexidine and calcium hydroxide is also effective against *E. faecalis* [13,23].

References

1. Soares JA, Leonardo MR, da Silva LA, Tanomaru Filho M, Ito IY (2006) Elimination of intracanal infection in dog's teeth with induced periapical lesions after rotary instrumentation: Influence of different calcium hydroxide pastes. *J Appl Oral Sci* 14(3): 172-177.
2. Delgado RJR, Gasparoto TH, Sipert CR, Pinheiro CR, Moraes IG, et al. (2010) Antimicrobial effects of Calcium Hydroxide and Chlorhexidine on *Enterococcus faecalis*. *J Endod* 36(8): 1389-1393.
3. Molander A, Reit C, Dahlen G, Kvist T (1998) Microbiological status of root-filled teeth with apical periodontitis. *Int Endod J* 38(1): 1-7.
4. Stuart CH, Schwrtz SA, Beeson TJ, Owatz CB (2006) *Enterococcus faecalis*: its role in root treatment failure and current concepts in retreatment. *J Endod* 32(2): 93-98.
5. Almyroudi A, Mackenzie D, McHugh S, Saunders WP (2002) The effectiveness of various disinfectants used as endodontic intra canal medicaments: an *in vitro* study. *J Endod* 28(3): 163-167.
6. Law A, Messer H (2004) An evidence based analysis of the antibacterial effectiveness of intracanal medicaments. *J Endod* 30(10): 689-694.
7. Meneges MM, Valera MC, Jorge AO, Koga-ito CY, Camargo CH, et al. (2004) *In vitro* evaluation of the effectiveness of irrigants and intracanal medicaments on microorganisms. *Int Endod J* 37(5): 311-319.
8. Shur AL, Sedgley CM, Fenno JC (2003) The antimicrobial efficacy of 'MGP' gutta percha *in vitro*. *Int Endod J* 36(9): 616-621.
9. Sinha N, Patil S, Dodwad PK, Patil AC, Singh B (2013) Evaluation of antimicrobial efficacy of calcium hydroxide paste, chlorhexidine gel, and a combination of both as intracanal medicament: An *in vivo* comparative study. *J Conserv Dent* 16(1): 65-70.
10. Gomes BP, Vianna ME, Sena NT, Zaia AA, Ferraz CC, et al. (2006) *In vitro* evaluation of the antimicrobial activity of calcium hydroxide combined with chlorhexidine gel used as intracanal medicament. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 102(4): 544-550.
11. Basrani B, Tjaderhane L, Santos JM, Pascon E, Grad H, et al. (2003) Efficacy of chlorhexidine- and calcium hydroxide-containing medicaments against *Enterococcus faecalis* *in vitro*. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 96(5): 618-624.

12. Mohammadi Z, Abbott PV (2009) The properties and applications of chlorhexidine in endodontics. *Int Endod J* 42(4): 288-302.
13. Singh H, Kapoor P (2014) A Comparative Evaluation of Antibacterial Efficacy of 'Activ Points' and 'Combi Points' as Intra-Canal Medicaments against *Enterococcus faecalis*: An *Ex Vivo* Study. *Oral Health Dent Manag* 13(3): 610-613.
14. Ebert J, Roggendorf MJ, Frank K, Petschelt A (2008) Antimicrobial activity of various 'active' gutta percha points against *Enterococcus faecalis* in simulated root canals. *Int Endod J* 41(3): 249-257.
15. Sen BH, Piskin B, Demirci T (1995) Observation of bacteria and fungi in infected root canals and dentinal tubules by SEM. *Endod Dent Traumatol* 11(1): 6-9.
16. Portenier I, Waltimo TMT, Haapasalo M (2003) *Enterococcus faecalis* -the root canal survivor and 'star' in post treatment disease. *Endodontic Topics* 6(1): 135-159.
17. Sundqvist G, Figdor D (2003) Life as an endodontic pathogen- Ecological differences between the untreated and root-filled root canals. *Endodontic Topics* 6(1): 3-28.
18. Delgado RJ, Gasparoto TH, Sipert CR, Pinheiro CR, Moraes IG, et al. (2010) Antimicrobial effects of Calcium Hydroxide and Chlorhexidine on *Enterococcus faecalis*. *J Endod* 36(8): 1389-1393.
19. Evans M, Davies JK, Sundqvist G, Figdor D (2002) Mechanisms involved in the resistance of *Enterococcus faecalis* to calcium hydroxide. *Int Endod J* 35(3): 221-228.
20. Siren EK, Haapasalo MP, Waltimo TM, Orstavik D (1997) *In vitro* antibacterial effect of calcium hydroxide combined with chlorhexidine or iodide potassium iodine on *Enterococcus faecalis*. *Eur J Oral Sci* 112(4): 326-331.
21. Lui JN, Lim VS, Song KP, Chen NN (2004) *In vitro* antimicrobial effect of chlorhexidine- impregnated gutta percha points on *Enterococcus Faecalis*. *Int Endod J* 37(2): 105-113.
22. Singh H, Gupta NP, Singh G, Kapoor P (2010) A comparison of antibacterial efficacy of "Activ Points" and 2% chlorhexidine solution against *Enterococcus faecalis*: an *in vitro* study. *Indian J Dent* 1(2): 49-54
23. Evans MD, Baumgartner JC, Khemaleelakul SU, Xia T (2003) Efficacy of calcium hydroxide: chlorhexidine paste as an intracanal medication in bovine dentin. *J Endod* 29(5): 338-339.
24. Korkmaz FM, Tuzuner T, Baygin O, Buruk CK, Durkan R, et al. (2013) Antibacterial activity, surface roughness, flexural strength, and solubility of conventional luting cements containing chlorhexidine diacetate/cetrimide mixtures. *J Prosthet Dent* 110(2): 107-115.