

Colostrum as the speed up key for ruminant newborn: what do we know and should further characterize

Abstract

Feeding and management programs are essential to determine neonatal health and survival. Therefore, an adequate and early intake of high quality colostrum, rich in antibodies and nutrients, is crucial. Some factors such as i) season; ii) length of the dry period and nutrition; iii) number of births; iv) litter size; v) body condition; vi) milk production; vii) fat and protein content; viii) somatic cell count and ix) breed/genetics have been associated with the quality of colostrum. Colostral nutritional and protective characteristics may be assessed through the protein concentration, which reflects the amount of albumin and globulins, indicating the success or failure in passive transference of immunity. Failure to passive immunoglobulin transfer results in an increased risk of morbidity and mortality of the newborn. This review addresses factors with potential to interfere with the quality of colostrum, prospecting future research in sheep.

Volume 9 Issue 3 - 2020

Luis Martins, Luiza da Costa Correa Oliveira
Department of Veterinary Medicine, University of Evora, Portugal

Correspondence: Luis Martins, Department of Veterinary Medicine, School of Science and Technology, and Mediterranean Institute for Agriculture, Environment and Development, University of Evora, Polo da Mitra, 7000-094 Evora, Portugal, Tel +351919498570, Email Imm@uevora.pt

Received: June 12, 2020 | **Published:** June 23, 2020

Introduction

In extensive sheep production systems current neonatal mortality can reach 15-20%, with a possible increase in case of multiparous deliveries. The amount of colostrum available at birth, to which mother's nutrition stands as a relevant factor, plays a major role in neonate survival.¹ Failure in passive transference of immunity increases the risk of newborn mortality, turning essential to provide an adequate amount of high-quality colostrum, early in the first hours of life, in order to establish that life-saving passive immunity.^{2,3}

Several factors have been described with potential to interfere with the quality of colostrum, such as the i) season; ii) length of the dry period and nutrition; iii) number of births; iv) litter size; v) body condition; vi) milk production; vii) fat and protein content; viii) somatic cell count and ix) breed/genetics.⁴⁻⁶ Further identification and characterization of parameters with possible implication in the quality of ovine colostrum, such as the season, the environmental temperature, the number of births is becoming more and more perceived. In addition, the length of the dry period, as well as the age at first delivery, should also be considered.

The frame for the colostrum quality

Colostrum characterization: Colostrum is the first liquid that provides food to the newborn. It is rich in proteins, vitamins, minerals, lactoferrin, lysozyme, antimicrobial peptides and growth factors, which are nutritionally important, but composition also evolves during the first days after delivery as it is shown in Table 1, regarding bovine.^{7,8} Colostrum is definitely different from milk, mainly because it contains a high level of antibodies against a diversity of potentially infectious agents playing a vital role in immunity and thus contributing to the development of the gastrointestinal tract, conditioning the endocrine system and metabolism, and providing an essential source of energy for young animals to generate heat, preventing hypothermia.^{9,10} While normal sheep's milk contains 12% solids, the same parameter in colostrum has a rate of 22%. This difference is mainly associated

to the concentration of immunoglobulins, but furthermore the solid matter is richer in casein, fat, protein and vitamins A, B12, D, and E, although lower in lactose.^{11,12} Approximately 92% of the total immunoglobulins belong to the IgG class, 6% to IgA and 2% to IgM.¹³

Table 1 Composition of colostrum⁸

	Peripartum	24h	72h	Milk
Solids (%)	23.9	17.9	14.1	12.9
Proteins (%)	14.0	8.4	5.1	3.1
Fat (%)	6.7	5.4	3.9	3.7
Lactose (%)	2.7	3.9	4.4	5.0
Minerals (%)	1.1	0.95	0.87	0.74
Immunoglobulins (%)	6.0	4.2	2.4	0.09

Passive transference of immunity

Passive transference of immunity is defined as the transfer of immunoglobulins from the mother to the newborn, through colostrum.¹⁴ Colostrum must be of high quality, with an IgG concentration ≥ 15 mg/mL.^{15,16} Content in immunoglobulins may be roughly perceived from total protein concentration in serum, which reflects the amount of albumin and globulins, but this technique lacks enough specificity.¹⁷ Confirmatory tests considered as the golden standard for IgG measurement include single radial immunodiffusion and ELISA.¹⁸ As passive immunity is essential for the newborn ruminant and failure in passive transfer increases the risk of morbidity and mortality, studying the factors that affect the quality of colostrum is an essential issue. According to Bartier et al.,¹⁵ and Tyler et al.,²¹ an incorrect management can lead to the death of the newborn and 39 to 50% of mortality may be associated to the failure of passive transfer of colostrum antibodies.^{20,21}

Parameters associated with colostrum quality

In ruminants, colostrum is the only way to acquire immunity soon after birth because of the epithelial placenta, which prevents *in utero* diffusion of immunoglobulins.^{2,22} Therefore, colostrum should be provided as early as possible after birth to ensure the appropriate passive transfer of antibodies from the mother to the neonate.¹⁸ It should make up to 10% of the birth weight in the first 24 hours postpartum.²³ Its supply must occur early, within this period, as the intestinal process of passive immunoglobulin absorption ceases (“intestinal closure”) approximately 24 hours after birth. Therefore, any delay in breastfeeding the newborn or low quality/amount of colostrum will reduce the level of immunoglobulins absorbed, increasing the susceptibility to pathogens, with a higher morbidity/mortality.²² A colostrum that has a minimum IgG concentration of 50 g/L, a total bacterial count <100,000 cfu/mL and a total coliform count <10.000 cfu/mL can be considered of good quality.^{24–28} The concentration of immunoglobulins may vary between breed, number of lactations, duration of dry periods, volume of colostrum, milking in the prepartum period and number/frequency of milking after delivery.^{29–31} Therefore, the success of the immune transfer depends on factors related to the sheep, the lamb and to the handling of these animals.¹⁶

a. Body condition

The body condition is based on a scoring system designed to evaluate the corporal reserves of animals and consists on the evaluation of several parameters of animal external appearance. It is associated with the energy balance, which correlates with body fat, milk production and reproduction, and should contribute to the maximum production with the best health status.³²

To maintain the desirable balance of body condition, meeting sheep’s nutritional needs, in order to guarantee a good quantity and quality of colostrum for the newborn is crucial.¹ At the end of pregnancy, nutrition strongly influences gland development and mammary cell differentiation, demanding a good nutritional management that promotes proper udder development with enough production of colostrum for the neonates.^{12,33–36} Preferably, optimum productivity is obtained in sheep presenting a medium body condition (e.g. 3) requiring an evaluation one month before the mating begins.³⁷

b. Season

First hours after birth are critical for newborn survival as it must neutralize the enormous heat loss in result of the environmental transition from the warm uterus to the outside. During the first 15 minutes after birth the internal temperature of a lamb decreases between 1 and 2 °C, regarding the 39 °C of the intrauterine environment. The colder the external environment is, the faster lamb’s metabolism must be, to maintain the homeothermic status. Other environmental conditions must also be considered, like wind speed and humidity, which will influence the rate of heat loss associated to the evaporation of amniotic fluids. Nardone et al.,²² reported heat stress related to high environmental temperature in association with impaired transference of IgG from plasma to the udder, resulting in lower colostrum concentration of IgG as well as IgA, but not IgM or lactoglobulin.³⁸ Therefore, good environmental management is crucial to allow the consumption of sufficient amount of good quality colostrum which in turn will meet the high energy thermoregulation requirements of small newborn ruminants.⁵

c. Duration of the dry period

The dry period is very important in a dairy farm because it allows the udder to rest between two successive lactations.³⁹ The length of the dry period can influence the concentration of immunoglobulins, since its secretion in the mammary gland starts between four to six weeks before delivery (Silva, 2019).¹⁵ In cattle, the maximum production of antibodies is reached with dry periods of at least 40 days.⁴⁰

d. Number of deliveries

The colostrum of first calving is less rich in immunoglobulins when compared to that of subsequent pregnancies and that happens because immunization against pathogens, to which animals become exposed to, with the production of specific antibodies (which will be transferred to the colostrum) is a lifelong process.^{41,42} Muller and Ellinger (1981) demonstrated that third lactation cows present a higher concentration of IgG in colostrum than those from first or second lactation, with a progressive increase in colostrum IgG until the third or fourth calves.⁴³ In addition to the number of births, reported the existence of a significant difference between lambs from simple and twin births. Twin lambs showed passive transfer failure as they born smaller, weaker and did not suck colostrum properly and in sufficient amounts.¹⁷

e. Milk production

The available information on the influence of good quality colostrum on ruminant productivity is somehow limited, however, Godden et al. (2009) described the failure in passive transference of immunity in heifers as associated with reduced milk and fat production at first lactation and delayed age at first calving.⁴⁴ It is also expected that those animals will show a colostrum deficit, considering the existence of a positive correlation between body weight and the concentration of immunoglobulins in colostrum.⁴⁵ Thus, in terms of productive management, the drying phase and nutrition during this period should be decisive for the quality of colostrum.

f. Genetics

Regarding the relevance of genetic factors to the quality of colostrum it is known that the concentration of IgG and nutritional factors in colostrum shows considerable variation in sheep, with special association to breed.⁵ Hence, a characterization of the qualitative variation of colostrum must consider the different management factors, like nutritional, environmental and reproductive, relating them with breed and aptitude. This will make possible the identification of associations between parameters, allowing formulating adjustments for productive improvement. The Merino/Suffolk comparative approach could stand as a good studying model.

Discussion

Some studies have associated the quality of colostrum with the calving season, with results showing that, in the case of cattle, animals that calved in winter presented a higher concentration of immunoglobulins and, in spring, a higher concentration of fat in the colostrum.⁴⁶ Volume is another parameter associated with the season, but not much scientific information is available.⁴ Recent studies in cattle have pointed to a higher concentration of immunoglobulins in colostrum, in association with energy-restricted diet during the dry period, when compared with more energetic diets, where a higher percentage of fat was associated with the supply of

concentrate between 0 and 3 weeks pre-delivery.⁴⁶ On the other hand, the quality of colostrum and the absorption of IgG by the newborn show a decrease in summer as a result of heat stress.⁴⁷ Furthermore, very relevant parameters for colostrum quality as concentration in IgG and nutritional factors have shown marked variation between different sheep breeds.⁵ Hence, aiming a better neonatal health and performance, studying colostrum quality should focus on the above parameters, regarding their influence on a specific breed.

Acknowledgments

None.

Conflicts of interest

Author declares that there are no conflicts of interest.

References

- Banchero G E, Milton JTB, Lindsay DR, et al. Colostrum production in ewes: a review of regulation mechanisms and of energy supply. *Animal*. 2015;9(5):831–837.
- Stelwagen K, Carpenter E, Haigh B, et al. Immune components of bovine colostrum and milk. *J Sci*. 2009;87(1):3–9.
- Morrill KM, Conrad E, Lago A, et al. Nationwide evaluation of quality and composition of colostrum on dairy farms in the United States. *J Dairy Sci*. 2012;95:3997–4005.
- Gavin K, Neiberghs H, Hoffman A, et al. Low colostrum yield in Jersey cattle and potential risk factors. *J Dairy Sci*. 2018;101:6388–6398.
- Kessler EC, Bruckmaier RM, Gross JJ. Immunoglobulin G content and colostrum composition of different goat and sheep breeds in Switzerland and Germany. *J Dairy Sci*. 2019;102(6):5542–5549.
- Zentrich E, Iwersen M, Wiedrich MC, et al. Effect of barn climate and management-related factors on bovine colostrum quality. *J Dairy Sci*. 2019;102(8):7453–7458.
- Kumar H, Goyal A, Kumar S, et al. Chemical and immunological quality of sheep colostrum: effect of breed. Editors: Naveen Kumar, Ravinder Kaushik Prince Chawla, et al; *Human Health and Nutrition*. 2017.
- Campos OL, Lizieire RS. *Dairy calves from birth to weaning*. Brazil: UOV – Online University of Viçosa; 2015.
- Abdel-Salam ZA, Abdel-Salam SA, Abdel-Mageed II, et al. Evaluation of proteins in sheep colostrum via laser-induced breakdown spectroscopy and multivariate analysis. *J Adv Res*. 2019;15:19–25.
- Rauprich ABE, Hammon HM, Blum JW. Influence of feeding different amounts of first colostrum on metabolic, endocrine and health status and on growth performance in neonatal calves. *J Anim Sci*. 2000;78:896–908.
- Selk GE. *Management factors that affect the development of passive immunity in the newborn calf*. Beef Cattle Handbook–2240. Extension Beef Cattle Resource Committee. 1998;1–7.
- Treacher TT. Artificial rearing of lambs: A review. *Vet Rec*. 1973;92:311–315.
- Smith WD, Dawson AM, Wells PW, et al. Immunoglobulin concentration in ovine body fluids. *Res Vet Sci*. 1975;19(2):189.
- McGuirk SM, Collins M. Managing the production, storage, and delivery of colostrum. *Vet Clin Food Anim*. 2004; 20:593–603.
- Silva DFM, Costa JN, Araújo AL et al. Failure of transfer of passive immunity in crossbred lambs (Santa Inês × Dorper): Effect on proteinogram and mortality rate from birth to weaning. *Cien Ani Bras*. 2009;1:158–163.
- Turquino CF, Flaiban KKMC, Lisbôa JAN. Transferência de imunidade passiva em cordeiros de corte manejados extensivamente em clima tropical. *Pesq Vet Bras*. 2011;31(3):199–205.
- Flaiban KKMC, Balarin MRS, Ribeiro ELA, et al. Transferência de imunidade passiva em cordeiros cujas mães receberam dietas com diferentes níveis de energia ou proteína no terço final da gestação. *Ci Anim Bras*. 200;181.
- Alves AC, Alves NG, Ascari IJ, et al. Colostrum composition of Santa Inês sheep and passive transfer of immunity to lambs. *J Dairy Sci*. 2015;98(6):3706–3716.
- Rovira LT, Pacheco JLP, Hernandez F, et al. Identification of factors affecting colostrum quality of dairy Lacaune ewes assessed with the Brix refractometer. *J Dairy Res*. 2017; 84:440–443.
- Bartier AL, Windeyer MC, Doepel L. Evaluation of on-farm tools for colostrum quality measurement. *J Dairy Sci*. 2015;98:1878–1883.
- Tyler JW, Hancock DD, Thorne JG, et al. Partitioning the mortality risk associated with inadequate passive transfer of colostral immunoglobulins in dairy calves. *J Vet Intern Med*. 1999;13:335–337.
- Nowak R, Poindron P. From birth to colostrum: early steps leading to lamb survival. *Reprod Nutr Dev*. 2006;46:431–446.
- Morrill D, Hartwig NR, Youngs C. *Colostrum and health of newborn lambs*. Sheep Management. USA: Iowa State University Press; 1995.
- Patel S, Gibbons J, Wathes DC. Ensuring optimal colostrum transfer to newborn dairy calves. *Cattle Pract*. 2014;22(1):95–104
- Arede MC. *Comparaçãodomaneio de viteloscémnascidosemexploraçõesleiteirasinglesas e americanas*. Dissertação de MestradoIntegradoemMedicinaVeterinária. Lisboa. 2013. Faculdade de MedicinaVeterinária. Portugal: Universidade de Lisboa; 2013.
- Godden S, McMartin S, Feirtag J, et al. Heat-treatment of bovine colostrum. II: Effects of heating duration on pathogen viability and immunoglobulin G. *J Dairy Sci*. 2006;89:3476–3483.
- Techmix calf management catalog. 2014.
- Maunsell F. Cow factors that influence colostrum quality. *Adv Dairy Technol*. 2014;26:113–121.
- Bamn. *A guide to colostrum and colostrum management for dairy calves*. USA: USDA–APHIS–CEAH–BAHM. Fort Collins; 2001.
- Weaver DM, Tyler JW, VanMetre DC, et al. Passive transfer of colostral immunoglobulins in calves. *J Vet Intern Med*. 2000;14:569–577.
- Radostits OM, Gay CC, Hinchcliff KW, et al. *Veterinary medicine: A textbook of the diseases of cattle, horses, sheep, pigs, and goats*. 10th ed. USA: Saunders Elsevier Eds; 2006.
- Gransworthy PC. *The effect of energy reserves at calving on performance of dairy cows*. Nutrition and lactation in the dairy cow. 1st edn. UK: Gransworthy Ed; 1998.
- Wallace JM, Bourke DA, Da Silva P, et al. Nutrient partitioning during adolescent pregnancy. *Reproduction*. 2001;122:347–357.
- Swanson TJ, Hammer CJ, Luther JS, et al. Effects of gestational plane of nutrition and selenium supplementation on mammary development and colostrum quality in pregnant ewe lambs. *J Anim Sci*. 2008;86:2415–2423.
- Tygesen MP, Nielsen MO, Norgaard P, et al. Late gestational nutrient restriction: effects on ewes' metabolic and homeorhetic adaptation, consequences for lamb birth weight and lactation performance. *Arch Anim Nutr*. 2008;62:44–59.
- Meyer AM, Reed JJ, Neville TL, et al. Nutritional plane and selenium supply during gestation affect yield and nutrient composition of colostrum and milk in primiparous ewes. *J Anim Sci*. 2011;89:1627–1639.

37. Moraes JCF, Souza CJH, Jaume CM. O uso da avaliação da condição corporal visando a máxima eficiência produtiva dos ovinos. Brazil: Ministério da Agricultura, Pecuária e Abastecimento; 2005.
38. Nardone A, Lacetera N, Bernabucci U et al. Composition of colostrum from dairy heifers exposed to high air temperatures during late pregnancy and the early postpartum period. *J Dairy Sci.* 1997;80:838–844.
39. Diniz AMMNS. O Maneio do vitelo recém-nascido: Efeito da quantidade ingerida de colostro na vitalidade dos vitelos. Dissertação de Mestrado em Engenharia Zootécnica/produção animal. Lisboa. Portugal: Instituto Superior de Agronomia. Universidade de Lisboa; 2007.
40. McGuirk SM. Colostrum: quality and quantity. *Cattle Pract.* 1998;6:63–66.
41. Schuch LFD. *Diarréia dos bezerros. Doenças de ruminantes e equinos.* Brazil: Varela SP; 2007:408–420.
42. Feitosa FLF, Camargo DG, Yanaka R, et al. Índices de falha de transferência de imunidade passiva (FTIP) em bezerros holandeses e nelores, às 24 e 48 horas de vida: valores de proteína total, de gama globulina, de imunoglobulina G e da atividade sérica de gama glutamiltransferase, para o diagnóstico de FTIP. *Pesq Vet Bras.* 2010;30:696–704.
43. Muller LD, Ellinger DK. Colostral immunoglobulin concentrations among breeds of dairy cattle. *J Dairy Sci.* 1981;64(8):1727–1730.
44. Godden SM, Haines DM, Konkol K, et al. Improving passive transfer of immunoglobulins in calves. II: Interaction between feeding method and volume of colostrum fed. *J Dairy Sci.* 2009;92:1758–1764.
45. Al-Sabbagh TA. Effect of nutritional status, age, breed, litter size and litter sex on colostral IgG levels for ewes delivered in Kuwait. *Elect J Env Agricult Food Chem.* 2008;7:3199–3205.
46. Dunn A, Ashfield A, Earley B, et al. Evaluation of factors associated with immunoglobulin G, fat, protein, and lactose concentrations in bovine colostrum and colostrum management practices in grassland-based dairy systems in Northern Ireland. *J Dairy Sci.* 2017;100:2068–2079.
47. Genc M, Coban O. Effect of Some Environmental Factors on Colostrum Quality and Passive Immunity in Brown Swiss and Holstein Cattle. *Isr J Vet Med.* 2017;72(3):28–34.